

by about 8% a year in low risk men of similar age and severity. Hospital mortality among women remained constant. Improvement in 12 month mortality of hospital survivors was seen at all levels of age and severity in both men and women; this was equivalent to a reduction of about 7% a year, controlling for age and severity. Men showed an overall reduction in 12 month mortality of 15% a year.

Women were older than men and had more severe infarcts. Greater proportions of women had the severest prognostic factors of cardiogenic shock, ventricular standstill, and third degree atrioventricular block, and on average women had higher scores on the Norris index and Killip scale. Nevertheless, a higher overall mortality in women than in men, more so in the later years, was still evident after adjusting for these factors. Statistical modelling predicted that, although the hospital mortality of men was nearly twice that of women of similar age and clinical characteristics in 1969, by 1983 it was less than two thirds. For 12 month mortality of hospital survivors the ratio of male to female mortality was constant at about 0.8. Therefore a higher mortality rate in women with acute myocardial infarction<sup>25,26</sup> cannot be explained by sex differences in the age or in the severity measures considered in this study.

Two predictors of mortality, the Royal Melbourne Hospital indices, were derived using the measured patient characteristics shown to be independently predictive of mortality in this study. These indices might prove to be useful because they are derived from large samples and give direct estimates of mortality. They suggest that ventricular fibrillation has become a less important predictor of mortality in recent years.

Patients who had never smoked were over 50% more likely to die in hospital than patients who had ever smoked, taking account of their ages and their having had more severe infarcts,<sup>27</sup> but the reasons for this are not clear.

There was a decrease of about 30% in the median time between onset of symptoms and admission to the coronary care unit over the study period, with at least a halving of long delays but no improvement in short delays. The decline in the severity of acute myocardial infarction over the study period was consistent, except that cardiogenic shock and ventricular tachycardia showed only a marginal decline in prevalence.

In recent years mortality from coronary heart disease has fallen in Australia,<sup>28,29</sup> while that from acute ischaemic heart disease has fallen in the United States.<sup>30</sup> If the epidemic of clinical coronary heart disease is waning then the residual cases would be expected to occur among older people and might be less severe, as we have observed. Our data are, however, hospital based, and there are other possible explanations such as confounding by variables other than age or sex, a tendency for younger patients with acute myocardial infarction to seek treatment outside public hospitals, or a change in admissions policy within the hospital. Nevertheless, the time trends we have found are consistent with a waning of the epidemic of clinical coronary heart disease in the community.

Our analyses have shown that, despite the fact that more recently admitted patients have included a higher proportion of people with milder infarctions, their better prognosis was greater than could be explained by the lower severity of their infarction. This was particularly true among men, who constituted 80% of patients with acute myocardial infarction.

This paper is dedicated to past and present nurses and doctors of the coronary care unit at the Royal Melbourne Hospital, and to Dr J Graeme Sloman, who started the unit in 1963 and its research data bank. We thank Dr J R E Fraser, Professor J D Mathews, and Mr J Campbell. This work was

supported by the Australian National Health and Medical Research Council.

- 1 Killip T, Kimball JT. Treatment of myocardial infarction in a coronary care unit: a two year experience with 250 patients. *Am J Cardiol* 1967;20:457-64.
- 2 MacMillan RL, Brown KWG, Peckham GB. Changing perspectives in coronary care. A five year study. *Am J Cardiol* 1967;20:451.
- 3 Restieaux N, Bray C, Bullard H, et al. 150 patients with cardiac infarction treated in a coronary care unit. *Lancet* 1967;i:1285-9.
- 4 Norris RM, Brandt PWT, Caughey DE, Lee AJ, Scott PJ. A new coronary prognostic index. *Lancet* 1969;ii:274-8.
- 5 Gillum RF, Folsom SR, Blackburn H. Decline in coronary heart disease mortality. Old questions and new facts. *Am J Med* 1984;76:1055-65.
- 6 Goldman L, Cook E. The decline in ischemic heart disease mortality rates. An analysis of the comparative effects of medical interventions and changes in lifestyle. *Ann Intern Med* 1984;101:825-36.
- 7 Marmot MG. Interpretations of trends in coronary heart disease mortality. *Acta Med Scand [Suppl]* 1985;701:58-65.
- 8 Thompson PL, Vandongen YK, Eccles JL, de Klerk NH. Assessment of the impact of coronary artery surgery on mortality after recovery from myocardial infarction. *Aust NZ J Med* 1984;14:424-9.
- 9 Hofvendahl S. Influence of treatment in a coronary care unit on prognosis in acute myocardial infarction. *Acta Med Scand* 1971;191(suppl):1-78.
- 10 Christiansen I, Iversen K, Skouby AP. Benefits obtained by the introduction of a coronary care unit: a comparative study. *Acta Med Scand* 1971;189:285-91.
- 11 Chapman BL. Effect of coronary care on myocardial infarct mortality. *Br Heart J* 1979;42:386-95.
- 12 Bain C, Siskind V, Neilson G. Site of care and survival after acute myocardial infarction. *Med J Aust* 1981;iii:185-8.
- 13 Mather HG, Morgan DC, Pearson NG, et al. Myocardial infarction: a comparison between home and hospital care for patients. *Br Med J* 1976;ii:925-9.
- 14 Hill JD, Hampton JR, Mitchell JRA. A randomised trial of home-vs-hospital management for patients with suspected myocardial infarction. *Lancet* 1978;ii:837-41.
- 15 Colling A, Dellipiani AW, Donaldson RJ, McCormack R. Teeside Coronary Survey: an epidemiological study of acute attacks of myocardial infarction. *Br Med J* 1976;ii:1169-72.
- 16 Rose G. The contribution of intensive coronary care. *British Journal of Preventive and Social Medicine* 1975;29:147-50.
- 17 Stern MP. The recent decline in ischaemic heart disease mortality. *Ann Intern Med* 1979;91:630-40.
- 18 Hunt D, Sloman G, Christie D, Pennington C. Changing patterns and mortality of acute myocardial infarction in a coronary care unit. *Br Med J* 1977;ii:795-8.
- 19 Hamer A, Hunt D, Sloman JG. The coronary care unit: experience at the Royal Melbourne Hospital. *Patient Management* 1980;Sep:49-57.
- 20 World Health Organisation. *Ischaemic heart disease registers: report of a working group*. Copenhagen: World Health Organisation, 1968.
- 21 World Health Organisation Regional Office for Europe. Myocardial infarction community registers. *Public Health in Europe* 1977;5:1-232.
- 22 McCullagh P, Nelder JA. *Generalized linear models*. London: Chapman and Hall, 1983.
- 23 Baker AJ, Nelder JA. *The GLIM System*. Release 3. Oxford: Numerical Algorithms Group, 1978.
- 24 Pathik B. *Changing patterns and trends in acute myocardial infarction*. [MD thesis.] Melbourne: University of Melbourne, 1986.
- 25 Puletti M, Senseri L, Curione M, Erba S, Borgia CL. Acute myocardial infarction. Sex related differences in prognosis. *Am Heart J* 1984;108:63-6.
- 26 Peter T, Harper RT, Luxton M, Pennington C, Sloman JG. Acute myocardial infarction in women. The influence of age on complications and mortality. *Med J Aust* 1978;ii:189.
- 27 Kelly TL, Gilpin E, Ahnves S, Henning H, Ross J. Smoking status at the time of acute myocardial infarction and subsequent prognosis. *Am Heart J* 1985;110:535-41.
- 28 National Heart Foundation of Australia. *Information Paper 19*. Canberra: National Heart Foundation, 1984.
- 29 Leeder SR, Gibberd RW, Dobson AJ, Lloyd DM. Declining mortality rates from ischaemic heart disease in Australia. *Aust NZ J Med* 1984;14:388-94.
- 30 Smith DP, Slater C. The pattern of ischaemic heart disease 1931-1980. *Stat Med* 1985;4:397.

(Accepted 1 August 1989)

## Correction

### Influence of maternal diet during lactation and use of formula feeds on development of atopic eczema in high risk infants

Two editorial errors occurred in this paper by Ranjit Kumar Chandra and others (22 July, pp 228-30). The conclusions in the abstract should read, "In families with a history of atopic disease [not eczema, as stated] mothers who breast feed should avoid common allergenic foods during lactation." The first sentence in the subjects and methods section should read, "In cases in which either of the baby's parents had a history [not a family history] of atopic disease mothers were asked whether they planned to breast feed exclusively."

### Case-control study of infections with *Salmonella enteritidis* phage type 4 in England

Two authors' errors occurred in this paper by Dr John M Cowden and others (23 September, pp 771-3). In the last sentence of the first paragraph on p 772 "the first three" was omitted, and later in the sentence a percentage of 0.02% was given. The sentence should read, "Of 137 patients, 26 (19%) ate at least one of the first three products specified in the table compared with only four (2%) of the 196 controls."