logical findings. Forty-three cases were not x-rayed, and 10 had negative radiological findings. In contrast with this, Hutchinson's cases all had positive radiological findings.

We have studied 415 infants with respiratory distress since August 1963. Of these, 24 had rarer causes of respiratory distress such as diaphragmatic hernia and pneumothorax, 18 died from other causes, and 59 were not x-rayed. Of the remaining 314 cases 150 had radiographs which were normal or showed only localized areas of atelectasis, and the mortality in these cases was only 2%, the three deaths all being in infants of less than 28 weeks' gestation. The good prognosis in these cases shows the necessity of excluding them from studies of the effects on mortality of different methods of treatment.

In your leading article, in discussing treatment of the respiratory distress syndrome with alkalis, you say that "further studies are required to determine the precise effects of these substances and the best way of giving them." We have carried out a study of the effects of intragastric sodium bicarbonate in the respiratory distress syndrome and our findings are to be published.² We found a mortality of 58% in 31 cases given intragastric sodium bicarbonate, and 39% in 69 cases given early milk feeds. These mortality figures are higher than those reported by Abraham and Brown, but refer only to cases with positive radiological findings, and are therefore more strictly comparable with the mortality figures quoted by Hutchison *et al.*

We would agree with the view expressed in your leading article that measures directed towards the relief of hypoxia have more effect in lowering mortality in the respiratory distress syndrome than correction of metabolic acidosis alone.—We are, etc.,

MARGARET E. R. STONEMAN.

ROSALIND M. OWENS.

Paediatric Research Unit
and Special Care Baby Unit,
Exeter City Hospital,
Exeter, Devon.

REFERENCES

- ¹ Hutchison, J. H., *et al.*, *Lancet*, 1962, 2, 465.
- ² Stoneman, M. E. R., and Owens, R. M., *Arch. Dis. Child.*, in press.

SIR,—I was interested to read Dr. J. M. Abraham's and Dr. R. J. K. Brown's article (9 September, p. 640) dealing with bicarbonate administration in the respiratory distress syndrome.

I have two comments to make regarding the treatment of respiratory distress syndrome from my own experience in the special care nursery, Royal Victoria Hospital, Belfast, over the past two or three years. In my experience oral sodium bicarbonate 8.4% may worsen and increase the difficulties in respiration in the very small premature baby, as it causes marked gastric distension with consequent embarrassment of chest expansion, making the infant's labours even more difficult, and for this reason I have not used this method.

Secondly, I have found that the umbilical vein is simple to use and is the least traumatic method for administering intravenous fluids to the small infant, and from my own experience phlebotrombosis has occurred only once during many dozens of catheterizations. Using this route for intravenous fluids the

infant is not subjected to repeated trauma of scalp vein infusions that may counteract any good effect obtained from the possible correction of the metabolic acidosis with the bicarbonate. Therefore I feel that this factor may account for the increased mortality in Dr. Abraham's and Dr. Brown's group given intravenous scalp vein fluids.—I am, etc.,

Belfast City Hospital,
Belfast 9.

MARK REID.

Problem of Unused Drugs

SIR,—I was interested to read the article "Collection of Unwanted Drugs from Private Homes" by Dr. W. A. Nicholson (16 September, p. 730). It is extremely difficult for each individual practitioner to gain any control over this mounting problem. However, I think it is worth mentioning an extremely simple technique which gives the doctor a small measure of control in some cases and which I am surprised is not used by many of my colleagues. Quite simply, each patient is asked to bring his supply of medicines with him to each consultation. A great number of patients are willing to co-operate in this way and the prescriber is thus enabled to determine the number of doses required until the next visit.

Using this method, it is surprising how often one meets the patient who says he is taking his tablets regularly but who in fact brings stock bottles which indicate the contrary. This discrepancy can be used as a valuable discussion point about the role which the medicaments are playing in the patient's life and the treatment of his illness. In addition, when he sees all the medicines together, the doctor is reminded of the complex regimen he sometimes asks a patient to undertake and can adjust his prescribing habits accordingly.

I realize that a determined saver of tablets for suicidal purposes will not be influenced by such a system, but I think that quite a number of patients can be spared the risk of having large quantities of dangerous chemicals in their households.—I am, etc.,

Institute of Psychiatry,
London S.E.5.

J. C. GUNN.

SIR,—Being involved as a medical officer of health member of a home safety committee which is currently considering a scheme for the recovery of unwanted drugs, I was particularly interested in the paper by Dr. W. A. Nicholson (16 September, p. 730) on the Hartlepool scheme for this service. However, one aspect which is causing my home safety committee much thought is the method of disposal of the drugs once they are collected. They would normally be brought in to pharmacists, who in turn would be required to devise their own means of disposal or to organize central disposal such as at a hospital. It is generally agreed that neither the refuse collection service nor the public sewers ought to be used. Incineration, either by the pharmacists or by a hospital, could involve some danger.

Would Dr. Nicholson or Dr. Milligan, the medical officer of health, please indicate in your correspondence columns what method of ultimate disposal is used in Hartlepool?

This information may be of value to others besides my own committee.—I am, etc.,

Office of the Medical
Officer of Health,
South Wilts County Districts,
Salisbury, Wilts.

JOHN LISHMAN.

Penicillin Encephalopathy

SIR,—Your leading article on this subject (12 August, p. 384) contains the following statement. "And if, as it seems, penicillin G is particularly apt to produce such an encephalopathy, other antibiotics might be considered as first choice in treatment." Since the patients to whom your article refers were all in a state of renal failure, and since the dosage of penicillin G exceeded 25 million units per day, I would like to know what antibiotics you have in mind that can be given more safely, in comparable doses, to uraemic patients. The semi-synthetic penicillin and the cephalosporins are not usually less toxic than penicillin G, while most other antibiotics are highly toxic or lethal in such subjects in much lower doses.

The doses mentioned (25–40 million units daily) correspond to amounts of 14 g. per day or more of penicillin G intravenously. At this level plasma concentrations reach 50 mg./100 ml. even in normal subjects, and are much higher in patients with bilateral nephrectomy, azotaemia, and other conditions referred to in the article. The remarkable fact is not that penicillin G causes temporary encephalopathy, but that it can be given at all to such patients in such dosage.

Penicillin G and its simpler semi-synthetic derivatives are among the least toxic of all drugs, being well-tolerated by mammalian tissues in concentrations far exceeding those of many physiological substances. There is evidence,¹ however, that penicillins in solution can polymerize as well as degrade, and it seems possible that some untoward effect might be attributable to these changes, which are minimized if solutions of the purified drugs for injection are used fresh, well within the stated date of expiry.—I am, etc.,

G. T. STEWART.

School of Public Health,
University of North Carolina,
U.S.A.

REFERENCE

- ¹ Stewart, G. T., *Lancet*, 1967, 1, 1177.

Visual Hallucinations in Cranial Arteritis

SIR,—With reference to the communication of Mr. C. T. Hart (9 September, p. 643) it may be of interest that in my own personal series of some 67 cases of giant-celled arteritis of the aged (which name, despite opposition, I still consider preferable) there have been three in whom formed visual hallucinations were a prominent symptom, and it would seem that this symptom may not be a particularly uncommon characteristic of what is fundamentally an extremely pleomorphic condition.

None of these patients developed signs related to the external visual apparatus, possibly owing to the institution of early treatment. One in particular was most striking, being an 85-year-old retired headmistress in a very good state of mental preservation who was extremely distressed by the continuous appearance of men's