

STRANGE BUT TRUE

Could mannequins menstruate?

Minna Rintala, Pertti Mustajoki

Women should have at least 17% of their weight as fat in order to have menarche and 22% in order to have regular cycles.¹ This fat contains easily mobilisable energy which provides nourishment during pregnancy and lactation.


Mannequins that display clothes in fashion shops may influence women's perception of ideal weight. We investigated the changing shape of display figures over time and determined whether women of their size would have enough fat for menstruation.

Subjects, methods, and results

Six old mannequins (from the 1920s, '30s, '50s, and '60s) in the Helsinki City Museum and modern dummies imported from three countries (Italy, Japan, and Malaysia) were investigated. Height and various circumferences were measured on every mannequin.

Body fat was calculated according to the formula: $\text{body fat (kg)} = 1.176 \text{ arm circumference} + 0.635 \text{ thigh circumference} - 44.255$.² Amount of fat was expressed as kg not as percentage of body weight because formulas for percentages of fat would need body weight or density, both of which would have been inaccurate estimations. Body fat of hypothetical women of same height as the dummies was estimated for two classes of body mass index, 20 kg/m² and 25 kg/m² according to the criteria of Scott and Johnston.³

The same anthropometric measurements and calculations were made on six average sized female medical students aged 20-30 years. In addition their subscapular, abdominal, and tricep skin folds were measured. Amount of body weight as fat was calculated from circumferences as in the dummies² and



Calculated amount of fat of mannequins from five decades and young women are shown as solid circles (●). The vertical lines give estimated amount of fat in hypothetical women of same height and with body mass indexes 20-25 kg/m²

from skin folds, taking the mean result obtained by four different formulas.^{2,4}


Arm, hip, and thigh circumferences of the modern display figures were 2-3 cm, 8 cm, and 4-5 cm less, respectively, than those of figures from before the second world war. Modern display figures and female medical students respectively had a mean height of 169 (range 166-172) cm and 175 (168-182) cm; arm circumference, 23 (21-24) cm and 28 (26-31) cm; hip circumference, 79 (73-82) cm and 93 (87-100) cm; thigh circumference, 43 (41-44) cm and 53 (51-57) cm.

Our figure shows that the calculated amount of fat of the display figures was mostly in the normal range before the 1950s but has been considerably less since

Third Department of
Medicine, University
Central Hospital,
00290 Helsinki, Finland
Minna Rintala, medical
student
Pertti Mustajoki, senior
physician

Correspondence to:
Dr Mustajoki.

BMJ 1992;305:1575-6



Mannequins through the
decades—(left to right) 1920s,
1930s, 1950s, 1960s, 1990s
(Helsinki City Museum)

then. Percentages of fat in the women students varied from 23.0% to 32.2% when calculated from circumferences and were comparable (21.6% to 31.5%) when calculated from skin fold measurements.

Comment

Our results show that display figures have become thinner with time and their proportions now differ considerably from those of normal young women. The structural changes seen in display figures during this century have led to a considerable reduction in the proportion of body weight calculated to be fat. A woman with the shape of a modern mannequin would probably not menstruate.

We often take a smiling or even horrified attitude towards the fashion phenomena of foreign cultures, but in our own society we get accustomed to the whims of fashion from early childhood. A woman with the proportions of a Barbie doll would be even thinner than the modern display figure (unpublished observations). Many of the phenomena of fashion are harmless, but the trend of extreme thinness is not without dangers. Most girls in industrialised countries are concerned with their body shape and practise occasional dieting.⁵

Although most of them survive without major problems, persistent dissatisfaction with one's own body may cause unnecessary concern and lessen happiness. In some girls this may proceed to the development of an eating disorder.

Why is the idealised weight so low? From the history of fashion we can see that during times of scarcity wide skirts with plenty of material were fashionable. When plenty of material was available the skirts were short and narrow. Similarly, being fat was socially desirable in times when there was a shortage of food. Now, in societies with excess food the ideal body shape is extremely thin. It seems that things difficult to achieve are pursued.

- 1 Frisch RE. Fatness, menarche and female fertility. *Perspect Biol Med* 1985;28:611-33.
- 2 Steinkamp RC, Cohen NL, Gaffey WR, McKey T, Bron G, Siri WE, *et al.* Measures of body fat and related factors in normal adults. II. A simple clinical method to estimate body fat and lean body mass. *J Chron Dis* 1965;18:1291-307.
- 3 Scott EC, Johnston FE. Critical fat, menarche, and the maintenance of menstrual cycles. *J Adolesc Health Care* 1982;2:249-60.
- 4 Smith DP, Boyce RW. Prediction of body density and lean body weight in females 25 to 37 years old. *Am J Clin Nutr* 1977;30:560-4.
- 5 Patton GC, Johnson-Sabine E, Wood K, Mann AH, Wakeling A. Abnormal eating attitudes in London schoolgirls—a prospective epidemiological study: outcome at twelve months. *Psychol Med* 1990;20:383-94.

How to gain weight by looking up

A J Newens, S N Jarvis, R George

Division of Epidemiology and Public Health, University of Newcastle upon Tyne, Newcastle upon Tyne NE2 4HH
A J Newens, research associate
R George, guest researcher

Gateshead Health Authority, Gateshead NE8 3EP

S N Jarvis, consultant in public health medicine

Correspondence to: Dr Jarvis.

BMJ 1992;305:1576-7

Weights recorded on birth certificates

Low birth weight is generally defined as births "less than 2500 g (up to and including 2499 g)"¹ and is associated with increased perinatal mortality risk.² An alternative criterion for low birth weight (less than 2800 g) was used by Townsend *et al* as part of a health index for districts in the Northern health region of England.³

During a study of perinatal mortality in Gateshead we examined the recording of birth weight over time to explain a fall from 7.8% in 1982 to 5.9% (below the average in Northern region) in 1990 in the proportion of babies weighing less than 2500 g at birth. There was less change (14.8-14.1%) when 2800 g was used as a criterion.

Method and results

Data on the weight of babies born to mothers resident in Gateshead were obtained from the Office of Population Censuses and Surveys for the years 1982-

91. These data were derived from birth notification documents completed at the time of birth, in grams, by a midwife and subsequently transferred to the registrar of births. In the Gateshead maternity unit, which accounted for two thirds of local births, the babies were weighed on the same analogue scales calibrated in 20 g and 1 ounce intervals throughout the study period. In the two Newcastle hospitals that deliver most other Gateshead babies both analogue and digital scales were used. Of the 26 131 birth registrations, 0.3% had birth weight missing.

Rounding of most weights, in 20 g or 30 g increments, was found throughout the study period. The pattern of rounding changed, however, over the period 1983 to 1985, and there was a tendency for weights to be 10 g or 20 g higher after 1985. As an example, the distribution of birth weights between 2770 g and 2850 g is shown in the table. The weight 2780 g remained over the study period, whereas 2800 g increased by 10 g to 2810 g, and 2820 g increased by 20 g to 2840 g. Although rounding was noted in at least two hospitals, individual maternity units were identifiable only from 1989 onwards.

Although it was not as clear cut as at higher birth weights, a degree of readjustment was apparent at 2500 g. In 1982-3 six births were recorded at exactly this weight, whereas in 1985-6 the number was 18. By 1989-90 the number had risen further to 30.

Discussion

Any cut off point for the definition of low birth weight calls for accurate recording of such information if useful comparisons are to be made between districts. The increase in the number of babies recorded as weighing exactly 2500 g after 1982-3, although small in comparison to the overall number of births, may nevertheless account for a part of the apparent fall in the proportion of low birthweight babies born to Gateshead mothers. As there was no upward revision

Weight (grams)	No of births recorded by year										Total
	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	
2780	22	21	18	26	22	11	22	23	27	23	215
2785		1		1							2
2788			1								1
2790	2	5	3		1			1	1		13
2792					1						1
2795	1			2							3
2800	22	20	15	2	1	4	8	5	1	1	79
2805		1		1	1						3
2807	3	1	1		1	1	1	1	2		11
2810	3	1	11	25	24	21	16	21	24	20	166
2815	2					1				1	4
2820	23	15	20	2	1	3	4	3	2	2	75
2824			1								1
2825		1	1							1	4
2826		1									1
2830	1		1	2	1	1	1	1	1	3	12
2835	4	3	3	4	2	3	2	1	3	6	31
2840	2	4	8	43	16	26	31	35	22	20	207
2845						1		1			2