

Health promotion for adolescents in primary care: randomised controlled trial

Zoe Walker, Joy Townsend, Laura Oakley, Chris Donovan, Hilary Smith, Zunia Hurst, Janet Bell, Sally Marshall

Abstract

Objectives To evaluate the effectiveness of inviting teenagers to general practice consultations to discuss health behaviour concerns and appropriate follow up care.

Design Randomised controlled trial, with participants randomised to a consultation (intervention) or usual care (control). Questionnaires completed at baseline, 3 months, and 12 months.

Setting Eight general practices in Hertfordshire, England.

Participants 1516 teenagers aged 14-15 years.

Intervention Consultations with practice nurses to discuss health concerns and develop plans for healthier lifestyles.

Main outcome measures Mental and physical health, "stage of change" for health related behaviour, and use of health services.

Results At baseline 970 teenagers completed questionnaires; 23% smoked, 35% had been drunk in the previous three months, 64% considered they ate unhealthily, 39% took little exercise, and 36% had possible depression. 41% (304) of teenagers invited attended for a consultation; over one third (112) were offered follow up care. More intervention group teenagers reported positive movement in stage of change for diet and exercise and in at least one of four behaviours (diet, exercise, smoking, drinking alcohol) at 3 months (41% *v* 31%, $P < 0.01$), but this did not persist at 12 months. There was marginally more positive change in actual behaviour by intervention teenagers at 3 months (16% *v* 12%, $P = 0.06$). Recognition of possible depression resulted in improved mental health outcomes at 3 and 12 months. 97% of attenders said they would recommend the intervention to a friend.

Conclusions Change in behaviour was slight but encouraging, and the intervention was well received and relatively cheap.

Introduction

Government policy in Britain is focused on preventing heart disease and stroke, accidents, cancer, and mental illness.¹ Adolescents have a high prevalence of risk factors associated with each of these priority areas,² but few receive any health promotion from general

practice.³⁻⁵ Consultations with teenagers are shorter than for other age groups.⁶ Very few reports of health promotion or screening for teenagers in the general practice setting have been published.⁷⁻¹¹ Any such initiatives have been welcomed by both health professionals and teenagers, but no controlled evaluations have been conducted to determine outcomes. Studies of primary care interventions with adults have reported small but significant effects on health behaviour.^{12, 13} Health professionals are known to have credibility with adolescents, and the advice they give may be important for teenage behaviour.³

This paper reports the first UK randomised controlled trial to evaluate the effectiveness of inviting teenagers to general practice for a health consultation and appropriate follow up care. Our hypotheses were that the intervention would provide a useful service, enable mental and physical health problems to be identified and addressed with appropriate information, and encourage healthy lifestyles. The intervention was based on models of self efficacy and behaviour change¹⁴ and teenagers' attitudes to general practice services.¹⁵⁻¹⁷ The intervention followed the structure suggested by the American Medical Association for consultations promoting self efficacy for healthy lifestyles with adolescents¹⁸ and was informed by the views of teenagers elicited from local surveys and focus groups.⁵ The study was approved by the local research ethics committees of East and North Hertfordshire and West Hertfordshire and was first piloted in one practice.¹⁴

Methods

Participants and randomisation

We identified teenagers aged 14 or 15 years on 1 January 1999 from eight general practice registers in Hertfordshire. The general practitioners sent letters to parents asking for consent. We then gave the teenagers a study number, stratified them according to sex, and randomised them by household with SPSS "random number seed," within the practice, to the intervention or control group.

Intervention

After parental consent had been obtained, teenagers in the intervention group received an appointment for a 20 minute consultation with the practice nurse to

Centre for Research in Primary and Community Care, University of Hertfordshire, Hatfield AL10 9AB

Zoe Walker
senior lecturer

Joy Townsend
director

Laura Oakley
research fellow

Royal College of General Practitioners Working Party on Adolescent Health, 14 Princes Gate, London SW7 1PU
Chris Donovan
chair of working party on adolescent health

Maltings Surgery, 8 Victoria Street, St Albans AL1 3JB
Hilary Smith
general practitioner
Janet Bell
nurse practitioner

Bennetts End Surgery, Gatecroft, Hemel Hempstead HP3 9LY

Zunia Hurst
general practitioner
Sally Marshall
practice nurse

Correspondence to: J Townsend, Health Promotion Research Unit, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT
joy.townsend@lshtm.ac.uk

(Dr Walker died on 10 September 2001)

bmj.com 2002;325:524

discuss their health and health related behaviour, focusing on topics of their choice. Twelve nurses received training in the study protocol, which aimed to improve adolescent self efficacy for behaviour change.¹⁴ Researchers (ZW and LO) observed two consultations for each nurse, to see how this was interpreted in practice. Teenagers who attended completed baseline and satisfaction questionnaires and provided saliva samples for measurement of cotinine to validate self reported smoking status. Teenagers who did not attend after two invitations were sent health promotion leaflets and baseline questionnaires at home. The control group received usual care and were sent baseline questionnaires at home. Both intervention and control teenagers were asked to complete follow up questionnaires at three months and 12 months and were invited for a consultation at 12 months to provide saliva samples for cotinine and have physical measurements taken. Up to three reminders were sent for all questionnaires. Mental and physical health, “stage of change” for four health related behaviours (diet, exercise, smoking, drinking alcohol), and use of general practice services were compared between groups. We used the Center for Epidemiological Studies depression scale for children to assess mental health.¹⁹

Masking

Because of the nature of the intervention, teenagers could not be blinded to their intervention status. However, both groups completed identical questionnaires, and the coding and entry of data were blinded to intervention status.

Analysis

We calculated the sample size needed to detect an effect size of a 10% reduction in the prevalence of

smoking (defined as smoking at least one cigarette a week). We aimed to recruit 1200 teenagers, assuming a statistical power of 80%, a 60% response rate, and 30% loss to follow up, resulting in a minimum of 254 teenagers in each of the two groups at follow up. We analysed the data with SPSS (version 10.0) by using Student's *t* test, Mann-Whitney U test, and χ^2 analysis as appropriate. Analysis was by intention to treat, and we assumed that teenagers who completed baseline questionnaires but did not return follow up questionnaires had made no changes. The main endpoints were health behaviour and stage of change for health related behaviours at 12 months; secondary endpoints were these outcomes at three months.

Results

Participant flow

We identified 1516 teenagers from the practice registers. All teenagers not withdrawn from the study were invited to complete questionnaires at baseline, three months, and 12 months, irrespective of whether they had completed earlier questionnaires. The figure illustrates the progress of participants through the trial. Of the 739 intervention group teenagers invited to a baseline consultation, 304 attended—173 (49%) of the girls and 131 (35%) of the boys. One was withdrawn at this stage because of learning difficulties. A further 200 intervention teenagers and 466 control teenagers completed baseline questionnaires at home. At three months 378 of the intervention group and 357 of the control group completed questionnaires.

At one year the 1358 teenagers remaining in the study were invited to attend a consultation (158 of the original sample had withdrawn from the study; 67 teenagers withdrew consent, 28 parents refused consent, 60 had moved house, and three withdrew for other reasons). Three hundred and ninety three (29%) attended, and non-attenders were sent questionnaires at home. Questionnaires were completed by 322 (49%) of the intervention group and 337 (48%) of the control group.

Baseline results


Demographic data

Respondents were aged 14-16 years (mean 14.8 years). Approximately half (478, 49%) were male. Most (868, 89%) were white, and 466 (48%) were in professional, managerial, or technical socioeconomic groups. Six hundred and forty eight (67%) lived in privately owned accommodation. Three quarters (706, 73%) lived with both natural parents.

Physical and mental health problems

One hundred teenagers (10.3%) reported a “major health problem”; asthma was the most common problem. Problems varied from sexual abuse to acne. More girls than boys reported problems—13% (64) *v* 8% (36); $\chi^2=5.23$, *df*=1, *P*=0.02).

Girls scored significantly higher than boys on the depression scale, indicating poorer mental health (17.1 *v* 11.9; 95% confidence interval for mean difference -6.49 to -4.01; *t*=-8.32, *P*<0.01). (A score of 16 indicates probable depression.)



Flow of participants in study

Table 1 Prevalence of self reported health related behaviour at baseline. Values are numbers (percentages)

Behaviour	Prevalence (n=970)
Smoking:	
Current smoker	225 (23)
Never smoked	613 (63)
Alcohol:	
Drinks alcohol more than once a week	98 (10)
Never drunk alcohol	231 (24)
Been drunk in past three months	342 (35)
Drugs:	
Has taken drugs	145 (15)
Knows someone who takes drugs	617 (64)
Has been offered drugs or encouraged to try drugs	299 (31)
Thinks will be tempted to take drugs in the future	122 (13)
Diet and exercise:	
Does not exercise regularly	378 (39)
Does not eat healthily	617 (64)
Believes is overweight	264 (27)
Believes is underweight	99 (10)

Health related behaviour

Table 1 shows the prevalence of health damaging behaviour at baseline. No significant differences existed between intervention and control groups. Self reported smoking behaviour was validated by a saliva cotinine test for attenders. Four samples (2%) provided by the 234 self reported non-smokers contained cotinine levels greater than 15 ng/ml, which suggested probable smoking or high exposure to passive smoking.²

Seven hundred and twenty six (78%) of the 930 teenagers who answered reported at least one of the health damaging behaviours measured (current smoking, drinking alcohol more than once a week, not eating healthily, not exercising regularly), and 128 (14%) reported at least three. No significant differences existed in the number of behaviours reported between intervention and control groups or between attenders and non-attenders. Girls reported significantly more health damaging behaviours than boys (mean 1.5 for girls, 1.2 for boys; 95% confidence interval for mean difference 0.25 to 0.51; $t=5.8$, $P<0.01$). Increased numbers of these behaviours were associated with poorer mental health on the depression scale ($r=0.31$, $P<0.01$).

Sexual health knowledge

One quarter (236, 24%) of the respondents did not know where to go for contraception without their parents knowing, and 30% (293) did not know where to get confidential advice. Almost one third (292, 30%) did not know that emergency contraception is effective for 72 hours, and two thirds (645) did not know that chlamydia is a sexually transmitted infection.

Use of health services

In the year before baseline, 709 (73%) of the teenagers had visited their general practitioner and 226 (23%) had visited their practice nurse at least once. Girls were significantly more likely to have visited, but there was no significant difference by age, ethnic group, or socio-economic group. Forty one per cent (398) had visited their school nurse, and one fifth (202) had visited another health professional. Teenagers who had seen

their general practitioner were more likely to report health damaging behaviour and poorer mental and physical health. This was most marked for current smokers and for teenagers who reported having been drunk in the previous three months, taking drugs or knowing someone who takes drugs, considering themselves overweight, or having a depression scale score indicating probable depression.

Consultation

Of the 290 attenders who answered the question, three quarters (230) wished to discuss at least one health behaviour topic with a practice nurse (table 2). A quarter (77) wished to discuss body shape and diet; 12% (35—half the smokers) wished to discuss smoking. Three quarters (225) indicated at least one behaviour they would like to work on changing; the most common were diet (144, 50%), exercise (104, 36%), dealing with stress (68, 23%), and smoking (39, 13%).

Three quarters (204) of the teenagers set a health related behaviour target. Sixty three planned to tackle more than one area; the most common topics were diet or nutrition (101), exercise (95), and smoking (29). Two hundred and sixty (85%) teenagers completed satisfaction questionnaires, and all but one were satisfied or fairly satisfied with their consultations. Most (246, 95%) said they had felt able to talk about all the issues they wanted to, 116 (45%) found the consultation better than they had expected, 242 (94%) thought the consultation was very useful or fairly useful, and 252 (97%) would recommend the service to a friend.

The nurses identified 48 (16%) teenagers as having some mental health problems, such as anxieties related to family, school, or friends. They identified 12 (4%) as likely to be depressed and a further 17 (6%) as possibly depressed; 21 teenagers were offered mental health follow up care.

Overall, more than one third (112) of the teenagers were offered follow up care. Thirty two were referred to the general practitioner, mostly (26, 81%) for physical health problems. Twenty seven teenagers were offered follow up appointments with the nurse for 45 different problems; these appointments tended to be to discuss health related behaviour (25, 56%) or mental health (15, 33%). In addition, four teenagers were referred to the asthma clinic, eight were offered counselling, two were referred to the family planning clinic, and a

Table 2 Topics teenagers most wanted to discuss. Values are numbers (percentages)

Topic	All (n=290)	Boys (n=122)	Girls (n=168)
Body size or shape	77 (27)	26 (21)	51 (30)
Acne	77 (27)	38 (31)	39 (23)
Diet	75 (26)	28 (23)	47 (28)
Exercise	62 (21)	28 (23)	34 (20)
Stress	59 (20)	15 (12)	44 (26)
General health	58 (20)	21 (17)	37 (22)
Smoking	35 (12)	17 (14)	18 (11)
Periods	34 (12)	NA	34 (20)
Contraception	32 (11)	6 (5)	26 (15)
Depression	24 (8)	9 (7)	15 (9)
Cancer	23 (8)	9 (7)	14 (8)
Alcohol	22 (8)	16 (13)	6 (4)
Sex	22 (8)	6 (5)	16 (10)
School	19 (7)	9 (7)	10 (6)
Nothing	60 (21)	33 (27)	27 (16)

NA=not applicable.

Table 3 Number (percentage) of teenagers in intervention and control groups who reported positive behaviour change at three month follow up

Positive change at three months	Intervention group (n=504)	Control group (n=466)	χ^2	df	P
Diet	21 (4.2)	10 (2.1)	3.20	1	0.07
Exercise	18 (3.6)	12 (2.6)	0.80	1	0.37
Smoking	33 (6.5)	23 (4.9)	1.16	1	0.28
Drinking	32 (6.3)	22 (4.7)	1.22	1	0.27
Any of four areas	82 (16.3)	56 (12.0)	3.59	1	0.06

Table 4 Comparison at three month follow up between intervention and control groups in number (percentage) of teenagers reporting positive movement along the stage of change continuum

Positive movement at three months	Intervention group (n=504)	Control group (n=466)	χ^2	df	P
Diet	89 (17.6)	55 (11.8)	6.57	1	0.01
Exercise	86 (17.1)	54 (11.6)	5.88	1	0.01
Smoking	41 (8.1)	25 (5.4)	2.93	1	0.09
Drinking	65 (12.9)	61 (13.1)	0.01	1	0.93
Any of four areas	206 (40.9)	143 (30.6)	10.91	1	<0.01

further 16 were encouraged to return if they wanted to follow up on any of the issues discussed.

Observation of a sample of clinics confirmed that the nurses were following the protocol, explaining the code of confidentiality, and trying to build a rapport with the teenagers. Mean length of consultations was 22.6 (SD 8.7) minutes. Most teenagers (288/304, 95%) saw the nurse alone.

Follow up

Health related behaviour

At three months marginally more teenagers in the intervention group than in the control group reported positive change in at least one of the four areas of health related behaviour (16% *v* 12%; $\chi^2=3.59$, *df*=1, *P*=0.06) (table 3). We found no significant difference between intervention and control group teenagers in the proportion who reported unhealthy lifestyles for the four areas measured individually at three month follow up, although the intervention group reported more positive movement in each case, except for drinking alcohol where change was similar for both groups. No significant difference existed at 12 months.

Stage of change for health related behaviour

Positive movement along the stage of change continuum indicates increased intention to improve the behaviour and assesses attempted as well as maintained change. At three months significantly more of the teenagers in the intervention group (41%) than in the control group (31%) indicated some positive movement in stage of change for at least one of the four key behaviours ($\chi^2=10.91$, *df*=1, *P*<0.01) (table 4). For individual behaviours, significantly more teenagers in the intervention group than in the control group made positive movement in stage of change for diet and for exercise. For smoking, the difference between the groups was of marginal significance (8% *v* 5%; $\chi^2=2.93$, *df*=1, *P*=0.09). No difference in drinking alcohol was found (table 4). At 12 months, no significant differences existed in the proportion showing positive movement in stage of change in at least one area.

Change in mental health score

We found no significant difference in change in mental health score overall between intervention and control group teenagers at three or 12 months. However, attending teenagers who scored 16 or more on the depression scale and who were identified by a nurse as having possible depression reduced their mental health score significantly more than did controls with these scores. The difference was significant both at three months (-8.14 intervention *v* -1.35 control; 95% confidence interval for mean difference -0.32 to -13.26 ; *t* = -2.10 , *P*=0.04) and at one year (-1.58 *v* 4.42 ; -0.53 to -11.48 ; *t* = -2.20 , *P*=0.03), indicating significantly improved mental health relative to controls.

Use of health services

Significantly more of the teenagers who were offered follow up had returned to the practice within three months compared with those offered no specific follow up—41% (46/112) *v* 28% (54/192); $\chi^2=4.82$, *df*=1, *P*=0.03). In the follow up year intervention group teenagers reported fewer visits to their general practitioner than did controls (1.74 *v* 2.05; -0.64 to 0.02 ; *P*=0.06). No difference existed between groups in the number of visits to practice nurses or school nurses.

At 12 months significantly more intervention group teenagers than controls knew where to go for confidential advice (83% (257/311) *v* 77% (249/325); $\chi^2=4.30$, *df*=1, *P*=0.04) and for information on contraception (88% (273/311) *v* 80% (259/325); $\chi^2=5.65$, *P*=0.02).

Discussion

The results of the trial are disappointing in that benefits (even where significant) were small. In this the results confirm the findings of trials of health promotion among adults and suggest that more needs to be offered to make a real difference.^{12 13} However, although the sample size was met, the analysis by intention to treat and assumptions of no change for the considerable number of non-responders is likely to have resulted in a considerable underestimate of the real effects of the intervention, especially at 12 months, when non-response was highest. Considering these factors, and that changing behaviour is notoriously difficult, the results provide an encouraging start, in terms of both actual behaviour change and stage of change. They suggest the need for a larger study with greater power as well as a more sustained intervention to help to maintain the short term gains and the positive intentions.

The trial confirmed the broad range of topics that adolescents would like the opportunity to discuss with a health professional. Nearly three quarters of teenagers visit their general practitioner each year—over half attend by themselves⁵—and these visits provide an opportunity to develop an adult relationship with health professionals and to discover the range of services available through the NHS. Confidentiality is a major issue that may stop this age group seeking advice, and teenagers in the intervention group were better informed at one year about access to confidential information, especially concerning sexual health. In consultations, neither teenagers nor health

professionals raise many of the topics of concern to teenagers,²⁰ but if topics have been previously discussed teenagers are more willing to discuss them again.²¹ If, as in this intervention, health professionals initiate discussion on sensitive issues and health promotion, teenagers may feel more able to raise such concerns on future occasions when they need to.

Most teenagers who attended did make plans to improve their health related behaviour. After three months intervention group teenagers reported positive movement in stage of change in significantly more lifestyle areas than did teenagers in the control group, and marginally more intervention group teenagers reported positive change in at least one of the four behaviours measured. Significant differences were not sustained at one year, which suggests that further research is needed into means of reinforcement to keep the teenagers motivated. Encouraging a written stepped plan might have increased levels of actual change.²²

Attendance did not vary by risk behaviour, age, or ethnic group. The practices included large and small practices in town based and rural locations. Lower attendance rates were recorded in the more deprived areas, but a higher proportion of teenagers who attended at these practices reported serious health concerns. Our findings may not be generalisable to inner city or very rural settings with particular problems of access but would be likely to apply to a wide range of practice settings and could be considered as part of a broad approach to providing health promotion and advice to this age group.

Recruitment to behaviours with health risks tends to be high at this age, so any reduction in uptake is important. Nurses expected to discuss sexual and mental health, drugs, and alcohol, and the teenagers did raise these concerns. But teenagers' greatest concern was about diet and exercise, both important factors in heart disease and stroke, while half the smokers wished to discuss smoking.

Interest in training about teenage health issues is increasing,²³ but no specific financial incentives for adolescent care are available, and a recent survey reported that only one in 10 health authorities had a policy on adolescents.²⁴ The intervention reported in this study was brief and inexpensive and possibly offset by reduced consultations in the following year. With minor financial outlay and some publicising of policies on confidentiality, practices can at least create an atmosphere that welcomes teenagers.

Conclusion

The consultations enabled the mental and physical health concerns of adolescents to be identified and addressed, were well received, and helped the teenagers to develop healthier lifestyles. A larger study with more substantial intervention is needed.

We thank all the teenagers and nurses for their essential contribution to this project.

Contributors: ZW and JT developed the research question. ZW, JT, CD, HS, JB, ZH, and SM developed the study design. ZW, LO, CD, HS, JB, ZH, and SM developed and led the training. SM and JB were among the 12 nurses who held clinics for teenagers in their practices. ZW undertook the day to day running of the trial, with the help of LO. All authors attended regular meetings to oversee the development and management of the trial. ZW and LO carried out data collection, coding, and cleaning. ZW carried out data analysis in discussion with all authors. ZW wrote

What is already known on this topic

Teenagers have a high prevalence of health damaging behaviour and have expressed a wish to discuss a broad range of health related issues with a health professional

Few teenagers receive health promotion advice or information from their general practice teams

What this study adds

General practice based health promotion consultations are welcomed by teenagers who attend

Such consultations provide an effective opportunity to identify and tackle mental and physical health problems and encourage healthy lifestyles

The effect on teenagers' actual lifestyles is modest

the first draft of the paper, and all authors have reviewed successive drafts. JT is guarantor for the study.

Funding: NHS Executive—Eastern Region, and HertNet (Hertfordshire Primary Care Research Network).

Competing interests: None declared.

- 1 Department of Health. *Our healthier nation*. London: Department of Health, 1998.
- 2 Department of Health. *Health survey for England. The health of young people '95-97: findings*. London: Stationery Office, 1998.
- 3 Walker ZA, Townsend J. The role of general practice in promoting teenage health: a review of the literature. *Fam Pract* 1999;16:164-72.
- 4 Walker Z, Townsend J. Promoting adolescent mental health in primary care: a review of the literature. *J Adolesc* 1998;21:621-34.
- 5 Walker ZAK, Townsend JL, Bell J, Marshall S. An opportunity for teenage health promotion in general practice: an assessment of current provision and needs. *Health Educ J* 1999;58:218-27.
- 6 Jacobson LD, Wilkinson C, Owen PA. Is the potential of teenage consultations being missed? A study of consultation times in primary care. *Fam Pract* 1994;11:296-9.
- 7 Donovan CF, McCarthy S. Is there a place for adolescent screening in general practice? *Health Trends* 1988;20:64-6.
- 8 Townsend J, Wilkes H, Haines A, Jarvis M. Adolescent smokers in general practice: health, lifestyle, physical measurements, and response to antismoking advice. *BMJ* 1991;303:947-50.
- 9 Smith A, Melville E. Targeting teenagers in the primary care setting. *Health Visitor* 1996;69:228-30.
- 10 Campbell A, Edgar S. Teenage screening in a general practice setting. *Health Visitor* 1993;66:365-6.
- 11 Cowap N. GPs need to be more proactive in providing health care to teenagers. *BMJ* 1996;313:941.
- 12 Family Heart Study Group. Randomised controlled trial evaluating cardiovascular screening and intervention in general practice: principal results of British family heart study. *BMJ* 1994;308:313-20.
- 13 Imperial Cancer Research Fund OXCHECK Study Group. Effectiveness of health checks conducted by nurses in primary care: final results of the OXCHECK study. *BMJ* 1995;310:1099-104.
- 14 Walker ZAK, Oakley LL, Townsend JL. Evaluating the impact of primary care consultations on teenage lifestyle: a pilot study. *Methods Inf Med* 2000;39:260-6.
- 15 Jacobson LD, Mellanby AR, Donovan C, Taylor B, Tripp JH, members of the Adolescent Working Group RCGP. Teenagers' views on general practice consultations and other medical advice. *Fam Pract* 2000;17:156-8.
- 16 McPherson A, Macfarlane A, Allen J. What do young people want from their GP? *Br J Gen Pract* 1996;46:627.
- 17 Churchill R, Allen J, Denman S, Fielding K, Williams D, Hollis C, et al. *Factors influencing the use of general practice based health services by teenagers: a descriptive study in the East Midlands*. Nottingham: University of Nottingham, 1999:109.
- 18 American Medical Association. *AMA guidelines for adolescent preventive services (GAPS): recommendations and rationale*. Baltimore, MD: Williams and Wilkins, 1994.
- 19 Olsson G, von Knorring AL. Depression among Swedish adolescents measured by the self-rating scale Center for Epidemiology Studies-depression child (CES-DC). *Eur Child Adolesc Psychiatry* 1997;6:81-7.
- 20 Maturing dangers. *Lancet* 1995;345:997-8.
- 21 Boekeloo BO, Schamus LA, Cheng TL, Simmens SJ. Young adolescents' comfort with discussion about sexual problems with their physician. *Arch Pediatr Adolesc Med* 1996;150:1146-52.
- 22 Gollwitzer PM. Goal achievement: the role of intentions. *Eur Rev Soc Psychol* 1993;4:141-85.

- 23 Richardson G, Parry-Langdon M, Jacobson L, Donovan C. *Bridging the gap: how do teenagers and primary care health care providers view each other?* Cardiff: Health Promotion Wales, 2000.
- 24 *Improving children's health: a survey of 1999-2000 health improvement programmes*. London: NSPCC, Children's Society, National Children's Bureau, 2000.

(Accepted 9 July 2002)