

Health, Kings Fund, and Imperial College for their help in identifying grey literature.

Contributors: see bmj.com

Funding: The project was funded by a grant from London Region NHS Executive, Organisation and Management Research and Development Programme.

Competing interests: None declared.

1 NHS Executive. *The NHS plan: a plan for investment, a plan for reform*. London: Department of Health, 2000.

2 Beresford P, Croft S. *Citizen involvement: a practical guide for change*. Basingstoke: Macmillan, 1993.

3 Barker J, Bullen M, de Ville J. *Reference manual for public involvement*. Bromley, West Kent, Lambeth, Southwark, and Lewisham Health Authorities, 1997.

4 NHS Executive. *Patient and public involvement in the new NHS*. Leeds: Department of Health, 1999.

5 McIver S. *Obtaining the views of users of health services*. London: King's Fund, 1991.

6 Kelson M. *User involvement: A guide to developing effective user involvement strategies in the NHS*. London: College of Health, 1997.

(Accepted 30 May 2002)

Involving users in the delivery and evaluation of mental health services: systematic review

Emma L Simpson, Allan O House

Abstract

Objectives To identify evidence from comparative studies on the effects of involving users in the delivery and evaluation of mental health services.

Data sources English language articles published between January 1966 and October 2001 found by searching electronic databases.

Study selection Systematic review of randomised controlled trials and other comparative studies of involving users in the delivery or evaluation of mental health services.

Data extraction Patterns of delivery of services by employees who were current or former users of services and professional employees and the effects on trainees, research, or clients of mental health services.

Results Five randomised controlled trials and seven other comparative studies were identified. Half of the studies considered involving users in managing cases. Involving users as employees of mental health services led to clients having greater satisfaction with personal circumstances and less hospitalisation. Providers of services who had been trained by users had more positive attitudes toward users. Clients reported being less satisfied with services when interviewed by users.

Conclusions Users can be involved as employees, trainers, or researchers without detrimental effect. Involving users with severe mental disorders in the delivery and evaluation of services is feasible.

Introduction

The Department of Health in the United Kingdom is committed to involving patients in the NHS; it is establishing the Commission for Patient and Public Involvement in Health. Users and carers have been involved in delivering and evaluating mental health services, but the effects of this involvement have not been rigorously assessed.¹⁻³

We sought evidence on involving users and the outcomes of involvement on clients (those receiving services).

Methods

We searched Medline, Embase, CINAHL, PsycINFO, HealthSTAR, Cochrane Controlled Trials Register, Web of Science, HMIC, and BIDS for references in English between January 1966 and October 2001 (see bmj.com).

We wrote to experts and organisations who had an interest in involving healthcare users asking whether they were aware of additional studies. We searched the references in all papers for additional studies, whether we included them or not. We searched collections by hand in the Health Sciences Library of the University of Leeds.

Inclusion and exclusion criteria

We included evaluations of the impact of research on services if users had an active role in the design or in collecting data. We also included studies about users who delivered services by training mental health professionals.

We included studies about delivery involving users in partnership with others if services were integrated by health professionals and users working together in a team; cross-consultation; or recruitment, training, supervision, or payment of users by healthcare providers.

Exclusion criteria

We excluded studies if they dealt with only

- Learning disabilities
- Involvement in decisions about a user's own treatment
- Providing information to users
- User satisfaction surveys that were researched by the provider (which do not require users' partnership)
- General health services not specifically aimed at mentally ill people
- Forensic services
- Services for mentally ill people which are not health related, such as housing or vocational rehabilitation
- Services with no contact with professionals or which could not be run by professionals which operate outside the mental health system—for example, self help groups



This is an abridged version; the full version is on bmj.com

Academic Unit of Psychiatry and Behavioural Sciences, University of Leeds, Leeds LS2 9LT

Emma L Simpson
research fellow
Allan O House
professor of liaison psychiatry

Correspondence to: E L Simpson
medelsi@south-01.novell.leeds.ac.uk

BMJ 2002;325:1265-8

We excluded studies which dealt only with the criteria in the box.

To assess the quality of the data, we sought the method of randomisation, evidence of blinding during data collection, and an intention to treat analysis.⁴ Meta-analysis was unacceptable because of heterogeneity in the study design and outcome measures so we summarised these qualitatively.⁴

Results

We identified five randomised controlled trials and seven other comparative studies.^{5–16}

The nature of users' involvement

Eight studies focused on involving users as service providers, mainly working as case managers in services for

clients with severe mental illness (table 1). Case managers need to engage clients, coordinate agencies, and help maintain effective delivery; the necessary skills are organisational and interpersonal rather than therapeutic. Two studies looked at the effects of involving users as trainers (table 2), and two studies considered involving users as interviewers (table 3).

The users who were involved were current or former users of mental health services who had had serious psychiatric illness—most commonly schizophrenia or bipolar disorder; many had been hospitalised. Employees who were or who had been users of mental health care services and interviewers had similar disorders to their clients.

Interviewers and employees who had been users all received training. Where applicable, this training was

Table 1 Involving current or former users of mental health services as providers in mental health services

Study	Involvement of	No of users involved and inclusion criteria	Study design (n=No of clients)	Measures of client* outcomes or service delivery patterns	Differences between groups
Solomon and Draine, 1994-6, USA ⁷	Case managers in community mental health service	4 in team (population changed over time); recent use of psychiatric services	Randomised controlled trial; 2 case management team conditions: employing users (n=48) and employing non-users (n=48)	Delivery: dates, locations, and manner of contact with clients Outcomes: income, level of functioning, quality of life, attitude to drugs compliance, social contacts, symptoms, inpatient days, treatment satisfaction	User employees: more face to face, fewer telephone or office based contacts 1 year: clients of user employees less satisfied with treatment, less family contact; 2 years: none
Paulson et al, 1997-2000, USA ⁸	Case managers in assertive community treatment programme	5 in team (population changed over time)	Randomised controlled trial; 3 conditions: assertive community treatment employing users (n=58), employing non-users (n=59), and usual care (n=61)	Delivery: time spent on categories of case manager activities Outcomes: time until first hospitalisation, arrest, emergency hospital care, or homelessness	User employees (compared with non-user ACT employees): longer in supervision, more flexible scheduling Clients of user employees: longer before hospital admission, fewer hospitalised, or had emergency care
O'Donnell et al, 1998-9, Australia ⁵	Client advocates attached to case management service	Number not stated	Randomised controlled trial; 3 case management conditions: clients focused with advocacy (n=45), clients focused (n=39), and standard care (n=35)	Outcomes: satisfaction with service, quality of life, functioning, family burden, inpatient days, use of crisis services	Family burden lower for client focused (2 groups combined) than for standard case management
Klein et al, 1998, USA ¹⁰	Peer counsellors alongside case management service	Number not stated; recovering from addiction	Comparative study; 2 case management conditions: with peer support (n=10) and standard (n=51)	Outcomes: hospital admissions, crisis events, social support, functioning, quality of life, drug use, satisfaction with service	Clients of peer support: fewer inpatient days, better social functioning, some quality of life improvements
Felton et al, 1995, USA ¹¹	Peer specialists on case management teams	3	Comparative study; 3 case management conditions: additional employees who were users (n=125), additional non-user employees (n=118), and no additional employees (n=68)	Outcomes: self esteem, engagement in programme, attitude to recovery, social support, quality of life, inpatient days, life problems, symptoms	Clients of user employees (compared with other 2 groups combined): more satisfied with living situations and finances, fewer reported life problems, less decline in contact with case managers
Chinman et al, 2000, USA ¹²	Case managers in outreach service	Number not stated; prior psychiatric treatment	Descriptive study; case management service sites separated into 2 conditions: sites with ≥10 clients of user employees (n=113) and sites with all or most services from non-user employees (n=630)	Outcomes: symptoms, quality of life, days of homelessness, social support, employment, relationship between client and case manager	None
Chinman et al, 2001, USA ¹³	Service providers in community outreach service	3 in team (population changed over time)	Comparative study; 2 conditions: programme with user employees (n=92) and matched sample of clients receiving usual care (n=79)	Outcomes: number of readmissions to hospital, inpatient days	None
Lyons et al, 1996, USA ¹⁴	Users as service providers in mobile crisis assessment service	8; prior psychiatric hospitalisation and medication or prior outpatient treatment	Descriptive study; compared working pairs in which: 1 or both of the pair had history of hospitalisation and neither user employee had a history of hospitalisation	Delivery: time spent on categories of duties, pattern of hospitalising clients	Working pairs in which at least 1 user employee had previous hospitalisation: more mobile outreach, fewer emergency responses, more hospitalising of clients involuntarily during routine dispatch

*Clients are recipients of services in which users are employed.

†These studies are also described in other publications cited elsewhere.²⁰

Table 2 Involving current or former users of mental health services as trainers of mental health service providers

Study	Users	Users involved	Study design	Outcome measures	Differences between groups
Cook et al, 1995, USA ⁶	Training mental health professionals	One person with bipolar disorder	Randomised controlled trial of 57 trainees trained by the user trainer or a non-user trainer	Trainee attitudes toward user employees; stigmatising factors of mental illness; likelihood of recovery	Trainees in the user trainer group had significantly more positive attitudes toward user employees and stigmatising factors of mental illness
Wood and Wilson-Barnet, 1999, UK ¹⁵	Student nurse classroom education	Not stated	Comparative study of 2 groups of students (n=15; n=14) differing in exposure to involving users in training	Student approach to mental health assessment; qualitative themes; empathy; individualised approach	Students with more and earlier exposure to user involvement, less jargon, more empathy, more individualised approach

similar to that received by employees who had not been users of mental health services. Payment was mentioned in most studies, and support workers were available to nearly all of the employees who were or who had been users of services.

Effects of users' involvement

The process of service delivery of employees who were or who had been users of mental health services differed from that of employees who had not. Users spent longer in supervision, in face-to-face contact with clients, or doing outreach work, and they spent less time on telephone or office work. Employees who were or who had been users had a higher turnover rate and had less distinct professional boundaries.

Employing users in, or alongside, case management services did not have any detrimental effect on clients in terms of symptoms, functioning, or quality of life. Clients of these services had some improved quality of life; they had fewer reported life problems and improved social functioning. Some clients were less of a burden to their families. In some studies, clients of employees who were or who had been users went for longer until hospital admission and fewer clients needed to be admitted to hospital, or stay in hospital was shorter, although time in hospital was not significantly different in all studies. Services employing people who had been users did not have lower client satisfaction. In one study, clients of employees who were or who had been users were less satisfied with treatment at follow up after one year, but they were not after two years.

Involving users in training gave trainees a more positive attitude toward employees who had been mentally ill and mental illness in general, or they looked at users as individuals. Clients reported being less satisfied with services when interviewed by other users of the service in evaluation research.

Design of study and interpretation

Our review of 298 papers about involving users in delivery of mental health services¹⁷ included only 12 comparative studies. We found five randomised trials, only one of which indicated the randomisation method used (alternate allocation according to an alphabetically ordered list of surnames). Researchers collecting

data were not blinded to treatment group in any of the studies. Four of the trials used intention to treat analysis.^{6 7 9 18} Of the other seven studies, researchers were blinded to treatment group in one study.¹¹ No intention to treat analysis was done in these studies.

Some studies were not set up to investigate users' involvement and the results were from a later analysis of routinely collected data.

Few standardised outcome measures were used unmodified. Some outcome measures were constructed for the particular study. Users were involved in the design of a questionnaire developed for one study. The use of modified rating scales could have led to bias, as has been shown for unpublished scales.¹⁹

Only small numbers of users were involved, with numbers ranging from one user to eight users in a team, making it difficult to apply findings to involving users in general. More users were involved in some studies because some users dropped out, generally for unstated reasons, and were replaced.

Sample sizes of studies were small, so estimates of effect were of low power. Clients were not always willing to see staff who the clients knew had had mental illness.

Discussion

The studies that we identified suggest that users of mental health services can be involved as employees of such services, trainers, or researchers without damaging them. In some studies, benefit was indicated for clients of employees who had been users of services, and, although this was not present across all studies, there were no serious disadvantages. The influence of trainers who had been users on the attitudes of trainees was positive; interviewers who had been users may have brought out negative opinions of services that would not otherwise have been obtained.

We found no comparative studies of users' involvement in planning mental health services, but other evaluations of users' involvement in planning in health services—including mental health services—have recently been reviewed.²⁰

Most of the studies we identified involved few users and have substantial methodological weaknesses. Studies of users as service providers mostly originated in

Table 3 Involving current or former users of a mental health service as interviewers of recipients of the service (clients) in evaluating mental health services

Study	Users involved	Design of study	Measurement of clients' views of service	Significant differences between groups
Clark, 1999, Canada ⁹	Four with severe mental disorder and prior psychiatric hospitalisation	Randomised controlled trial of user interviewers (n=60) and staff interviewers (n=60)	Extremely positive and negative responses and general satisfaction	Clients interviewed by user interviewers gave more extremely negative responses about services
Polowczyk, 1993, USA ¹⁶	People with schizophrenia or affective disorder in remission	Comparative study of user interviewers (n=225) and staff interviewer (n=305)	Satisfaction score	Clients interviewed by user interviewers gave lower service satisfaction scores

What is already known on this topic

Involving health service users in the NHS is recommended in UK government policy

Involving users in mental health services is generally seen as worthwhile, but the effects of involving users have not been thoroughly evaluated, and few attempts to draw evaluations together have been made

What this study adds

The few comparative studies of users' involvement that have been published indicate that involving users as employees, trainers, or researchers has no negative effect on services and may be of benefit

the United States and were confined to a case management model. Government policy in the United Kingdom strongly supports the development of involving users in the delivery and evaluation of mental health services. Little evidence exists on the effectiveness of such programmes, and more formal evaluations are needed.

Contributors: See bmj.com

Funding: Non-conditional grant from Leeds Community and Mental Health Services Trust.

Competing interests: None declared.

- 1 Mental Health Task Force User Group. *Forging our futures: lighting the fire*. London: Department of Health, 1995.
- 2 NHS Health Advisory Service. *Voices in partnership: involving users and carers in commissioning and delivering mental health services*. London: Stationery Office, 1997.
- 3 Department of Health. *National service framework for mental health modern standards and service models*. London: DoH, 1999.
- 4 NHS Centre for Reviews and Dissemination. *Undertaking systematic reviews of research on effectiveness*. CRD, 2001.
- 5 O'Donnell M, Parker G, Proberts M, Matthews R, Fisher D, Johnson B, et al. A study of client-focused case management and consumer advocacy:

- the community and consumer service project. *Aust N Z J Psychiatry* 1999;33:684-93.
- 6 Cook JA, Jonikas JA, Razzano L. A randomized evaluation of consumer versus nonconsumer training of state mental health service providers. *Community Ment Health J* 1995;31:229-38.
 - 7 Solomon P, Draine J. The efficacy of a consumer case management team: 2-year outcomes of a randomized trial. *J Ment Health Adm* 1995;22:135-46.
 - 8 Paulson R, Herinckx H, Demmler J, Clarke G, Cutter D, Birecree E. Comparing practice patterns of consumer and non-consumer mental health service providers. *Community Ment Health J* 1999;35:251-69.
 - 9 Clark CC, Scott EA, Boydell KM, Goering P. Effects of client interviewers on client-reported satisfaction with mental health services. *Psychiatr Serv* 1999;50:961-3.
 - 10 Klein AR, Cnaan RA, Whitecraft J. Significance of peer support with dually diagnosed clients: findings from a pilot study. *Res Soc Work Pract* 1998;8:529-51.
 - 11 Felton CJ, Stastny P, Shern DL, Blanch A, Donahue SA, Knight E, et al. Consumers as peer specialists on intensive case management teams: impact on client outcomes. *Psychiatr Serv* 1995;46:1037-44.
 - 12 Chinman MJ, Rosenheck R, Lam JA, Davidson L. Comparing consumer and nonconsumer provided case management services for homeless persons with serious mental illness. *J Nerv Ment Dis* 2000;188:446-53.
 - 13 Chinman MJ, Weingarten R, Stayner D, Davidson L. Chronicity reconsidered: improving person-environment fit through a consumer-run service. *Community Ment Health J* 2001;37:215-29.
 - 14 Lyons JS, Cook JA, Ruth AR, Karver M, Slagg NB. Service delivery using consumer staff in a mobile crisis assessment program. *Community Ment Health J* 1996;32:33-40.
 - 15 Wood J, Wilson-Barnett J. The influence of user involvement on the learning of mental health nursing students. *NT Research* 1999;4:257-70.
 - 16 Polowczyk D, Brutus M, Orvieto AA, Vidal J, Cipriani D. Comparison of patient and staff surveys of consumer satisfaction. *Hosp Community Psychiatry* 1993;44:589-91.
 - 17 Simpson EL, House AO, Barkham M. *A guide to involving users, ex-users and carers in mental health service planning, delivery or research: a health technology approach*. Leeds: Academic Unit of Psychiatry and Behavioural Sciences, University of Leeds, 2002.
 - 18 Clarke GN, Herinckx HA, Kinney RF, Paulson RI, Cutler DL, Lewis K, et al. Psychiatric hospitalizations, arrests, emergency room visits, and homelessness of clients with serious and persistent mental illness: findings from a randomized trial of two ACT programs vs. usual care. *Ment Health Serv Res* 2000;2:155-64.
 - 19 Marshall M, Lockwood A, Bradley C, Adams C, Joy C, Fenton M. Unpublished rating scales: a major source of bias in randomised controlled trials of treatments for schizophrenia. *Br J Psychiatry* 2000;176:249-52.
 - 20 Crawford MJ, Rutter D, Manley C, Weaver T, Bhui K, Fulop N, et al. Systematic review of involving patients in the planning and development of health care. *BMJ* 2002;325:1263-5.
 - 21 O'Donnell M, Proberts M, Parker G. Development of a consumer advocacy program. *Aust N Z J Psych* 1998;32:873-9.

(Accepted 14 October 2002)



This is an abridged version; the full version is on [bmj.com](http://www.bmj.com)

Assessment of neurocognitive impairment after off-pump and on-pump techniques for coronary artery bypass graft surgery: prospective randomised controlled trial

Vipin Zamvar, David Williams, Judith Hall, Nicola Payne, Clare Cann, Karen Young, S Karthikeyan, John Dunne

Editorial by Taggart

Department of Cardiac Surgery, University Hospital of Wales, Cardiff CF14 4XW
Vipin Zamvar
consultant surgeon
continued over

BMJ 2002;325:1268-71

Abstract

Objective To assess neurocognitive impairment after the off-pump and on-pump techniques for coronary artery bypass graft surgery in patients with triple vessel disease.

Design Randomised controlled trial.

Setting University Hospital of Wales, Cardiff.

Participants 60 patients undergoing coronary artery bypass graft surgery for triple vessel disease prospectively randomised to the off-pump or on-pump technique.

Main outcome measures Change in scores in nine standard neuropsychometric tests administered preoperatively and at 1 and 10 weeks postoperatively.

Results The on-pump group showed a significantly greater deterioration in scores for two and three tests at 1 week and 10 weeks postoperatively, respectively, than the off-pump group. The on-pump group also showed a significantly higher incidence of major deterioration in one of the tests both 1 week and 10 weeks postoperatively. The incidence of neurocognitive impairment at 1 week postoperatively was 27% (8 out of 30) in the off-pump group and 63% (19 out of 30) in the on-pump group ($P=0.004$); and at 10 weeks postoperatively was 10% (3 out of 30) in the off-pump group and 40% (12 out of 30) in the on-pump group ($P=0.017$).

Conclusion Off-pump coronary artery bypass graft surgery results in less neurocognitive impairment than the on-pump technique.