Doctors are not equipped to judge their peers

Instruments developed for physicians to rate their peers in practice are of poor quality, and their validity remains questionable. Reviewing more than 4500 papers, Evans and colleagues (p 1240) identified

only three rating scales that had psychometric data about their development or their validity and reliability. None referred to a theoretical framework, and they all lacked construct and criterion validity. Instruments developed for peer assessment have not been developed in accordance with best practice, say the authors, and should be used with caution.

POEM*

Lowering homocysteine did not prevent recurrent stroke

Question Does lowering the homocysteine level in patients with ischaemic stroke prevent recurrent stroke?

Synopsis Observational studies have indicated that raised homocysteine levels are associated with an increased risk of recurrent stroke, acute coronary events, and mortality. Folic acid supplements have been suggested as a means to lower homocysteine levels and reduce these risks. Fifty six centres in the United States, Canada, and Scotland coordinated to enrol 3680 adults with non-disabling cerebral infarction. The patients received the best medical and surgical care and were randomised in double blind fashion (concealed allocation assignment) to receive an identical daily supplement of either a high dose multivitamin (pyridoxine 25 mg, cobalamin 0.4 mg, and folic acid 2.5 mg) or low dose multivitamin (pyridoxine 200 µg, cobalamin 6 µg, and folic acid 20 µg) preparation. Participants were contacted every three months by telephone or clinic visits for up to two years. A total of 93% of subjects were available for follow up analysis. Although the article does not specify that outcomes were assessed by individuals blinded to treatment group assignment, contact with the corresponding author confirmed that this was true. With intention to treat analysis, the mean reduction in total homocysteine level was significantly greater in the high dose group, but there were no significant differences between the two treatment groups regarding recurrent stroke, acute coronary event, or death. The study had an 80% power to detect a 30% reduction in recurrent ischaemic stroke over two years of follow up.

Bottom line Lowering homocysteine levels with a high dose B vitamins in patients with ischaemic stroke did not lower the risk of recurrent stroke. This study had the ability to detect a 30% difference in stroke rates over two years, if one exists. The likelihood of dying or experiencing a myocardial infarction was not affected by the treatment, although the study was probably too small and too short to find a difference if one truly exists. It seems benign (and possibly helpful) to recommend a multivitamin supplement for all patients at risk of atherosclerotic vascular disease, but there does not seem to be any solid evidence yet supporting routine measurement of homocysteine levels.

Level of evidence 1b (see www.infoPOEMs.com/levels.html). Individual randomised controlled trials (with narrow confidence interval).

Toole JF, Malinow MR, Chambless LE, et al. Lowering homocysteine in patients with ischemic stroke to prevent recurrent stroke, myocardial infarction, and death. The Vitamin Intervention for Stroke Prevention (VISP) randomized controlled trial. *JAMA* 2004;291:565-75.

©infoPOEMs 1992-2003 www.infoPOEMs.com/informationmastery.cfm

* Patient-Oriented Evidence that Matters. See editorial (*BMJ* 2002;325:983)

Editor's choice

Angioplasters and thrombolysers

"Paper will surely take over from screens. It's attractive, highly portable, flexible, smells good, and is easy to notate. Within a year, we hope, the *BMJ* should be available not just electronically but also on paper." History, of course, worked the other way round, and paper preceded electronic communication by centuries—but sometimes the race to be historically first may be close. Thrombolysis arrived as a routine treatment for patients with acute myocardial infarction just before primary angioplasty, but there are now enthusiasts arguing that primary angioplasty should become the first line treatment. Peter Bogarty and James Brophy have imagined a world in which primary angioplasty came first and the enthusiasts are arguing for thrombolysis (p 1257).

There are strong arguments for both treatments, and these are well displayed in a "For and against" by David Smith and Kevin Channer (p 1255). Sometimes we have difficulty urging the combatants to argue their case as strongly as possible. This week—perhaps because both protagonists are cardiologists—there is no such problem. If you were to read either piece alone you would, I suggest, be convinced. Reading them together, you will probably be grateful that it's somebody else's decision on whether to make primary angioplasty available to all.

Smith presents incontrovertible evidence that if normal myocardial blood flow can be restored then mortality can be dramatically reduced—to perhaps less than 1%. Of 100 patients with infarcts about a quarter are not eligible for thrombolysis. Of those treated, a third to a half will have normal blood flow restored, but this would have happened in 10% anyway. Some patients are thus exposed to the hazards of thrombolysis but for no benefit. In contrast, argues Smith, angioplasty can be offered to everybody, needn't be done on those who spontaneously reperfuse, and can achieve normal blood flow in 90-97% of those treated. A meta-analysis of 23 trials shows substantial benefits over thrombolysis in mortality, non-fatal infarction, and haemorrhagic stroke. Surely, you'll think after reading this, the case is made.

But, argues Channer, you must be careful in extrapolating from the results of trials done in selective patients, mostly in the United States, to the real world and most other countries. A large Dutch trial in which patients were transferred from district general hospitals to regional centres found no improvement in all cause mortality. Even in the United States, registry data show less benefit than was expected from trials. The problem is the inevitable delay that occurs with angioplasty.

Perhaps the answer is yet another strategy—possibly home thrombolysis followed by rescue angioplasty. We also need some evidence on costs and benefits and on what it would be like to live in a world full of angioplasters. What will they all do when, as predicted by our heart tsar, deaths from heart disease disappear?

Richard Smith *editor* rsmith@bmj.com

To receive Editor's choice by email each week subscribe via our website: bmj.com/cgi/customalert