but not for irritable bowel syndrome. This antipathy was due to the lack of bodily location, patients failing to conform to the work ethic and traditional "sick role," and disagreement with patients

over causes and management. Referral for mental health interventions was unlikely because doctors were unfamiliar with them or thought them unavailable or unnecessary.

POEM*

Data on treating bronchiolitis is severely limited

Question How effective are the various treatments for bronchiolitis?

Synopsis The authors systematically reviewed Medline and the Cochrane Collaboration's database of controlled clinical trials for randomised controlled trials published in English that assessed the effectiveness of various treatments for bronchiolitis. They used an explicit and reasonable set of search terms and did a limited search for unpublished data. The team assessed the quality of each study, with disagreements adjudicated by consultation and consensus. The authors reported 44 studies of the most commonly used agents: epinephrine, β_2 agonist bronchodilators (albuterol and salbutamol), corticosteroids, and ribavirin. They found a handful of studies evaluating inhaled helium, RSV-immunoglobulin, Chinese herbs, and so forth (see www.ahrq.gov/clinic/evrptfiles.htm#bronch). In general, most studies were quite small, of limited quality, looked at short term improvement, and failed to assess clinically important outcomes. Racemic epinephrine was studied against β_2 agonists in eight randomised controlled trials of 660 infants. Five of these studies assessed hospitalisation; only two reported either fewer admissions or shorter stays. Most of the 13 studies of nebulized β_2 agonists had multiple treatment arms. Seven of the studies assessed hospitalisation; none reported meaningful differences in rate or duration. Four studies evaluated oral corticosteroids and found no consistent effect on hospitalisations or duration of stay. Parenteral corticosteroids had no effect on clinical outcomes. In 10 randomised controlled trials of ribavirin (Copegus, Rebetol), the overall study quality was low. Of the five studies reporting on clinically important outcomes, four failed to show any effect on rate of hospitalisation, length of stay, duration of illness, or use of intensive treatment. The sole study finding a benefit (on use of intensive treatment) used sterile water as the placebo. As sterile water can induce bronchospasm, thereby making ribavirin seem more effective, this study has been criticised.

Bottom line In spite of the large number of studies assessing various treatments for bronchiolitis, in general the studies have been small, of poor quality, and don't assess clinically important end points. The treatments may be effective, however, but just unproved. To really judge their effectiveness, we'd need large, well designed studies that include clinically important outcomes. Until then, bronchiolitis treatment is in the "can do, but not required" category—there are few "musts" or "must nots," so don't obsess about overtreatment or undertreatment.

Level of evidence 1a (www.infoPOEMS.com/levels.html) systematic review of randomised trials displaying worrisome heterogeneity.

King VJ, Viswanathan M, Bordley WC, et al. Pharmacologic treatment of bronchiolitis in infants and children: a systematic review. *Arch Pediatr Adolesc Med* 2004;158:127-37.

©infoPOEMS 1992-2003 www.infoPOEMS.com/informationmastery.cfm

* Patient-Oriented Evidence that Matters. See editorial (*BMJ* 2002;325:983)

Editor's choice

Medicine and man's fall

If medicine were simply a matter of prescribing drugs and wielding scalpels then monkeys—or at least robots—might make adequate doctors. It's the human bit, as in most enterprises, that makes medicine tricky, fascinating, and difficult. Medical journals might be accused of ignoring much of that complexity with their diet of drug trials and systematic reviews. Increasingly, however, we are publishing qualitative research that probes the interactions between doctors and patients.

Three authors from Australia and Canada have asked patients receiving palliative care and their families what they want to be told and found that many were unhappy with the communication process—particularly the disclosure of the initial diagnosis (p 1343). The authors identify six things that are important in communication in such circumstances: playing it straight, making it clear, showing that you (the doctors) care, giving time, pacing the information, and staying the course. Each individual, of course, wants something different and may be far from clear, even to themselves, about what they do want—putting a high premium on the doctor's sensitivity.

The two content areas that are most important are prognosis and hope. The difficulty with prognosis is that many patients want to be told but do not want to know. Not everything needs to be said at once, and patients commonly want lots of information at the beginning but ever less as they come closer to death. Hope, including for some hope of a long life when close to death, is vital, and one of the worst experiences for participants was "to have hope dashed by a rushed or insensitive health carer." To be honest but be able to keep hope alive are two fundamental attributes of a "good doctor."

As I read this paper I couldn't help thinking that it deals with the easier stage of terminal care—when the patients have registered with a palliative care programme. The worst communication failures come, I suspect, in that time between the doctors thinking the patient will die and the patients and families registering for palliative care.

Four authors from London delve into some of the darker aspects of medicine by comparing and contrasting doctors' perceptions of patients with chronic fatigue syndrome and irritable bowel syndrome—both complex, poorly understood conditions where symptoms, outlook, and responses to treatment are similar (p 1354). Yet one doctor says: "I would rather treat a whole surgery full of people with irritable bowel syndrome than people with chronic fatigue." The authors identify five reasons for the difference, but one is nothing more complicated than one illness having a precise bodily location. The most important finding from the study is that with both conditions doctors are reluctant to use "mental health approaches," which may often be effective.

Richard Smith *editor* rsmith@bmj.com

To receive Editor's choice by email each week subscribe via our website: bmj.com/cgi/customalert