

### Summary points

To apply the offside rule correctly in a football game, the referee must be able to keep in his visual field at least five objects at the same time—two players of the attacking team, the last two players of the defending team, and the ball

This is beyond the capacity of the human eye, which may explain why so many offside decisions are controversial

The use of modern technology such as freeze frame television to aid referees' decisions is necessary for the offside rule to be applied correctly

be able to keep in his visual field at least five objects at the same time (two players of the attacking team, the last two players of the defending team, and the ball), and this may not be compatible with the normal eye function—especially as these five objects can be anywhere within the defenders' half of the pitch, an area of at least 3200 m<sup>2</sup>. This may explain at least some of the instances when television replays of a game clearly show that the offside rule was not properly implemented. The key factor in applying this rule correctly is that the player in question must be in the offside position at the exact time when the ball is passed from a team mate, not when the player receives the ball or when the ball is en route between the players.

By reviewing the physiology of the eye movements likely to be involved in assessing an offside position, I have shown that the relative position of four players and the ball cannot be assessed simultaneously by a referee, and unavoidable errors will be made in the attempt. The use of modern technology during games—freeze frame television and frame by frame analysis—is advisable to limit these errors.

I thank Nuria Navarro Zaragoza for her translation of the paper into English and Joaquín Zaragoza Celdrán for producing the figures.

Funding: None.

Competing interests: None declared.

- 1 Burde RM. Control de los movimientos de los ojos. In: Moses R, ed. *Fisiología del ojo de Adler*. 8th ed. Buenos Aires: Panamericana, 1988:122-54.
- 2 Maños Pujol M. Fisiología del sistema sacádico. In: Gavilán C, Gavilán J eds. *Fisiología y fisiopatología del sistema oculomotor*. Madrid: Garsi, 1987:43-53.
- 3 Gavilán J, Gavilán C, Sarriá MJ. Saccadic movements: a computerized study of their velocity and latency. *Acta Otolaryngol* 1983;96:429-36.
- 4 Leigh RJ, Zee DS. The saccadic system. In: Leigh RJ, Zee DS, eds. *The neurology of the eye movements*. Philadelphia: FA Davis, 1983:39-68.
- 5 Shickman GM. Funciones visuales que dependen del tiempo. In: Moses R, ed. *Fisiología del ojo de Adler*. 8th ed. Buenos Aires: Panamericana, 1988:591-633.
- 6 Schmidt D, Abel LA, Dell'Osso LF, Daroff RB. Saccadic velocity characteristics. Intrinsic variability and fatigue. *Aviat Space Environ Med* 1979;50:393.
- 7 Schalén L, Pyykkö I, Juhola M, Magnusson M, Jänti V, Henriksson N. Intraindividual variation in oculomotor performance in man. *Acta Otolaryngol* 1984;406(suppl):212-7.
- 8 Gavilán C, Gavilán J. *Fisiología y fisiopatología del sistema oculomotor*. Madrid: Garsi, 1987:56-66.
- 9 Moses RA. Acomodación. In: *Fisiología del ojo de Adler*. 8th ed. Buenos Aires: Panamericana, 1988:285-302.
- 10 Olivós Arroyo R. *Teoría del fútbol*. Seville: Wanceulen, 1992.
- 11 International Football Association Board. *Laws of the game*. Zurich: Fédération Internationale de Football Association (FIFA), 2003:24.

## An unusual complication of Kocher's manoeuvre

A Pimpalnerkar, A Datta, D Longhino, N Mohtadi

Royal Centre for Defence Medicine, Birmingham B29 6JD

A Pimpalnerkar consultant in orthopaedics and sports

A Datta specialist registrar in orthopaedics

Sports Medicine Centre, Calgary, Canada T2N 1N4

D Longhino orthopaedic resident

N Mohtadi consultant sports surgeon

BMJ 2004;329:1472-3

A 25 year old man presented to the emergency room with a right anterior glenohumeral dislocation after a fall. He was attended to by a 30 year old, left hand dominant orthopaedic resident. Kocher's manoeuvre was attempted to reduce the dislocation.

As the patient's arm was slowly externally rotated the patient resisted forcefully and suddenly internally rotated his arm. The resident immediately heard a "pop" in his own left shoulder, followed by a burning sensation with subsequent weakness.

The resident was an active individual, who did regular weight training and played competitive ice hockey. He had no previous shoulder problems and denied using anabolic steroids.

On examination there was a loss of the normal anterior axillary fold, bruising in the upper arm (figure), and weakness in adduction and internal rotation. Neurovascular examination was normal. Plain radiographs were unremarkable. Magnetic resonance imaging confirmed a near total rupture of the pectoralis major tendon from its humeral attachment. The marrow signal was normal.

The resident had surgical repair of the tendon five days after the injury. The tendon was reattached to the

lateral lip of the bicipital groove using Mitek anchors and number 2 Ethibond. At final follow up at 24 months, the functional range of shoulder movement had returned with good muscle strength and endurance as assessed with isokinetic and functional testing.

### Discussion

Pectoralis major tendon injury, though rare, was first described in 1822 by Patissier.<sup>1</sup> Such injuries are becoming more common owing to an increasing number of recreational and professional athletes. The mechanism is of a violent eccentric contraction and is associated with doing bench presses, wrestling, and water skiing.<sup>2</sup>

The pectoralis major muscle is a powerful adductor, internal rotator and flexor of the shoulder. The sternocostal and clavicular heads insert at the lateral lip of the bicipital groove to form two separate laminae that are oriented perpendicular to each other. When the arm is abducted and externally rotated the sternocostal fibres are maximally stretched.

Numerous reduction techniques for anterior glenohumeral dislocations have been described. They



Loss of normal anterior axillary fold and contour of the pectoralis belly, along with substantial bruising around the axilla

fall into three main categories: traction techniques (Hippocrates,<sup>3</sup> Stimson<sup>4</sup>) leverage techniques (Kocher,<sup>5</sup> Milch<sup>6</sup>), and scapular manipulation.<sup>7</sup>

Kocher's manoeuvre was originally described in 1870.<sup>5</sup> The affected arm is flexed at the elbow and adducted against the side of the body. The forearm is rotated externally and the upper arm lifted in a sagittal plane as far as possible and finally internally rotated. There have been many adjustments to this widely used technique, especially as the additional torque resulting

from adduction and internal rotation have been associated with humeral shaft fractures in elderly people.<sup>8,9</sup>

To perform Kocher's manoeuvre the surgeon must abduct and externally rotate his own arm, this means the fibres of the sternocostal head of his pectoralis major are maximally stretched. If a patient is not well sedated or has not received adequate analgesia he or she is more likely forcefully to resist the reduction. This results in a sudden uncoordinated eccentric contraction of the surgeon's pectoralis major muscle, which in our case resulted in a pectoralis major rupture. If adequate sedation and pain relief are not possible, reduction under general anaesthetic may be necessary to minimise the risk of complications for both the patient and the doctor.

Competing interests: None declared.

- 1 Patisserie P. *Traite des maladies des artisans* Paris, 1882:162-4.
- 2 Bak K, Cameron EA, Henderson JJP. Rupture of pectoralis major: a meta-analysis of 112 cases. *Knee Surg Sports Traumatol Arthrosc*. 2000; 8:113-9.
- 3 Mattick A, Wyatt JP. From Hippocrates to the Eskimo—a history of techniques used to reduce anterior dislocation of the shoulder. *J R Coll Surg Edinb* 2000;45:312-6.
- 4 Stimson LA. An easy method of reducing dislocations of the shoulder and hip. *Med Record* 1900;356.
- 5 Kocher T. Eine neue Reduktionsmethode für Schulterverrenkung. *Berliner Klin Wchenschr* 1870;7:101-5.
- 6 Milch H. Treatment of dislocation of the shoulder. *Surgery* 1938;3:732-40.
- 7 McNamara RM. Reductions of anterior shoulder dislocations by scapular manipulation. *Ann Emerg Med* 1993;22:1140-4.
- 8 Beattie TF, Steedman DJ, McGowan A, Robertson CE. A comparison of Milch and Kocher techniques for the acute anterior dislocation of the shoulder. *Injury* 1986;17:349-52.
- 9 Riebal GD, McCabe JB. Anterior shoulder dislocation: a review of reduction techniques. *Am J Emerg Med* 1991;9:180-8.

## Easy ways to resist change in medicine

Allen F Shaughnessy, David C Slawson

These time honoured techniques will help doctors resist the forces that prevail on them to change their ways of taking care of patients

Numerous forces have been imposed on physicians to make them change their practice behaviours. Under the guise of "quality improvement," managed care organisations, accrediting bodies, and the government are meddling in medicine. Even continuing medical education, previously a form of intellectual entertainment or a forum for much needed sleep, has refocused its efforts towards improving the care of patients (figure).

### Techniques to resist change

Techniques are available, however, that will allow doctors to practise unimpeded by new information that should induce change. To avoid succumbing to both the inner and outer forces prompting change, we offer the following techniques and methods.

#### Don't pay attention

Get so busy with your practice that you do not have the time to read, attend meetings, understand your own practice, or observe the practice of colleagues. Forget about Stephen Covey's seventh habit of "sharpening

the saw."<sup>11</sup> Create your own habit of sawing harder and harder, with a dull saw.

#### Attack the data

When provided with new information that may require a change in practice:

- Firstly, diss the source. No one expects you really to believe information from sources outside your specialty or geographical area.
- Secondly, question the validity of the information. Every study or report contains some loophole in its

#### Levels of belief

- Class 0: Things I believe
- Class 0a: Things I believe despite the available data
- Class 1: Randomised controlled clinical trials that agree with what I believe
- Class 2: Other prospectively collected data
- Class 3: Expert opinion
- Class 4: Randomised controlled clinical trials that don't agree with what I believe
- Class 5: What you believe that I don't

Department of Family Medicine and Community Health, Tufts University School of Medicine, 100 Hospital Road, Malden, MA 02148, USA  
Allen F Shaughnessy  
*faculty*

Department of Family Medicine, Box 800729, University of Virginia Health Sciences Center, Charlottesville, VA 22908, USA  
David C Slawson  
*B Louis Barnett Jr professor of family medicine*

Correspondence to: D C Slawson  
dslawson@virginia.edu

BMJ 2004;329:1473-4