Improving the doctor-patient relationship in chronic illness

Collaborative management of chronic illness is undermined by neglect of emotional and psychological factors in both the patient and the doctor, argue Campbell and McGauley (p 667). They highlight the gap

between the importance of the rising number of chronic diseases and the relatively inadequate attention to the effect of managing chronic illness on the doctor and how this manifests in the therapeutic relationship. Drawing on their work in a high secure psychiatric hospital, they suggest how the gap can be addressed in medical education.

POEM*

Two-drug hypertension regimen increases CVD mortality

Question In the treatment of adults with hypertension, which other drug class added to diuretics most effectively reduces adverse cardiovascular events?

Synopsis Evidence shows that diuretics are equal to or superior to other agents as first line therapy for most patients with hypertension. More than one drug class, however, is frequently required to control hypertension. It is unclear which other drug classes, added to diuretics, optimally reduce adverse cardiovascular events. The investigators evaluated data obtained from women with hypertension enrolled in the women's health initiative observational study, a prospective cohort study of 93 676 women aged 50-79 at baseline, 94% of whom were followed up for a mean of 5.9 years. Antihypertensive medication was determined from original bottles brought to baseline visits and matched to a pharmacy database. End points were ascertained from mailed questionnaires, direct report, telephone follow-up, medical records, and death certificates. The investigators do not specifically state whether outcomes were assessed by individuals blinded to treatment groups. Among women with hypertension but no history of cardiovascular disease (CVD), monotherapy with calcium channel blockers versus diuretics was associated with an increased risk of death from cardiovascular disease (number needed to treat to harm over six years [NNTH/6] = 143; 95% confidence interval 59 to 3898). In similar patients, a two drug regimen of a diuretic plus calcium channel blocker was associated with a statistically significant increase in death from cardiovascular disease compared with both a diuretic plus β blocker and a diuretic plus angiotensin converting enzyme inhibitor (NNTH/6 = 93; 34 to 3898). Both analyses were adjusted for age, race or ethnicity, smoking, high cholesterol requiring medication, body mass index, physical activity, hormone use, and diabetes.

Level of evidence 2b (see www.infoPOEMs.com/levels.html). Individual cohort study or low quality randomised controlled trials < 80% follow up.

Bottom line In women with hypertension and no history of cardiovascular disease, a regimen of a diuretic plus either a β blocker or angiotensin converting enzyme inhibitor reduces the risk of mortality due to cardiovascular disease, compared with a diuretic plus calcium channel blocker. The evidence continues to mount that calcium channel blockers should be the agent of last resort in the treatment of most patients with hypertension.

Wassertheil-Smoller S, Psaty B, Greenland P, et al. Association between cardiovascular outcomes and antihypertensive drug treatment in older women. *JAMA* 2004;292:2849-59.

©infoPOEMs 1992-2003 www.infoPOEMs.com/informationmastery.cfm

* Patient-Oriented Evidence that Matters. See editorial (*BMJ* 2002;325:983)

Editor's choice

Charcot would have approved

As he walked the wards of the Salpetriere Hospital in 19th century Paris, Charcot allegedly gave only two prognoses, solemnly pronouncing at the end of each bed "Il va mourir," or "Il va mourir aujourd'hui." A bleak message definitely, and bluntly delivered, but there is something enviable in its certainty and simplicity. Of course he was short of effective treatments, and accurate prognosis was often the only useful thing medicine could offer, so the art of prognosis was highly valued and was a big part of the medical curriculum.

Today, while death remains stubbornly inevitable, modern medicine can and does postpone it. But has this success made us forget the importance of prognosis? Murray and colleagues think it has (p 611), especially in people with progressive chronic disease. Being busy managing and monitoring a long term illness can prevent (or perhaps protect) us from seeing that the patient is in irreversible decline, and so we rob them of the chance to plan for a good death. To avoid such "prognostic paralysis" and identify patients who could benefit from holistic palliative care, Murray advises asking the simple question, "Would I be surprised if my patient were to die in the next 12 months?"

Thanks to modern medicine and advances in public health we are living longer. But we are also living longer with illness. As this theme issue highlights, chronic diseases now affect six out of ten adults and one in five to six children in the UK (p 657) and by 2020 will account for four fifths of the global burden of disease. Managing chronic disease presents enormous challenges to health systems and individuals. Several of this week's articles tell us that to meet these challenges we need (as well as more money and better systems) new skills and training (pp 633, 637, 644, 651) to help us shift the emphasis away from acute hospital medicine to coordinated, collaborative, patient centred care (p 662). As Campbell and McCauley point out (p 667), unless doctors are trained to understand chronic illness, their unrealistic expectations may foster negative attitudes and the perception that some patients are hard to treat when they fail to respond.

Finding easy and accurate ways to predict response is key to informing patients and planning treatment. So it's good news from Doust and colleagues in their systematic review (p 625) that a simple blood test for B-type natriuretic peptide (BNP) strongly predicts the risk of death from heart failure, both in people with and without symptoms. Clinical assessment of heart failure is notoriously difficult, as is identifying those patients most likely to benefit from treatment. BNP turns out to be better at this than traditional prognostic indicators such as New York Heart Association class, serum creatinine, and possibly even left ventricular ejection fraction. Charcot would have approved.

Fiona Godlee *editor* (fgodlee@bmj.com)

To receive Editor's choice by email each week subscribe via our website: bmj.com/cgi/customalert