What are the ethical considerations? On p 834, Siotia and colleagues present the case of a woman with Down's syndrome who developed postoperative

hypoxia after a dual operation for an umbilical hernia and a total hysterectomy. All *BMJ* readers are invited to use the rapid responses feature on bmj.com to discuss the case.

POEM*

Trial of labour carries small but real increase in risk

Question What are the risks associated with trial of labour after caesarean delivery?

Synopsis This was an observational study in which women with a previous caesarean delivery and a current singleton pregnancy decided whether to pursue a trial of labour (they were not randomised). Among the 45 988 women at the 19 academic medical centres who qualified for the study, 9013 had clear indications for repeat caesarean and were excluded, as were 3276 who presented early in labour without a documented plan for trial of labour. Of the remainder, 17 898 underwent a trial of labour and 15 801 had an elective repeat caesarean delivery. Women choosing trial of labour were younger, more likely to be African-American, less likely to smoke, less likely to be married, and much more likely to have had a successful vaginal delivery (49.8% *v* 15.8%). A multivariate analysis adjusted for potential confounders of the two composite outcomes: adverse maternal events (endometritis, transfusion, uterine rupture, hysterectomy, death, dehiscence, thromboembolic disease, haematoma, cystotomy, bowel injury, and ureteral injury) and adverse neonatal events (intrapartum stillbirth, hypoxic-ischaemic encephalopathy, and neonatal death). Data were gathered prospectively, using standard forms and definitions. About a third of eligible women underwent a trial of labour (the current national rate is 12.7%). A number of important complications were significantly more common in the trial of labour group, most notably uterine rupture (0.7% *v* 0%), endometritis (2.9% *v* 1.8%), and uterine dehiscence (0.7% *v* 0.5%). There was no significant difference in maternal deaths between groups (0.02% *v* 0.04%). The risk of any adverse maternal event was higher in the trial of labour group (5.5% *v* 3.6%; number needed to treat to harm (NNTH) = 53, 95% confidence interval 43 to 68). Uterine rupture was especially common among women with augmented labour (odds ratio 2.4) or induced labour (odds ratio 2.9). Of the 114 uterine ruptures, nine resulted in neonatal death or encephalopathy. The overall risk of stillbirth, hypoxic-ischaemic encephalopathy, or neonatal death was very low, but was significantly higher in the trial of labour group (0.38% *v* 0.13%; NNTH = 398, 269 to 721).

Bottom line The risks of trial of labour after caesarean delivery are small but cannot be ignored. The most important is an increase in the risk of neonatal death, stillbirth, or encephalopathy (number needed to treat to harm = 398), which is higher than that seen in other research (*BMJ* 2004;329:19-25). This article provides some additional guidance for physicians and patients making this important shared decision.

Level of evidence 2b (see www.infoPOEMs.com/levels.html). Individual cohort study or low quality randomised controlled trial < 80% follow-up.

Landon MB, Hauth JC, Leveno KJ, et al. Maternal and perinatal outcomes associated with a trial of labor after prior cesarean delivery. *N Engl J Med* 2004;351:2581-9.

©infoPOEMs 1992-2003 www.infoPOEMs.com/informationmastery.cfm

* Patient-Oriented Evidence that Matters. See editorial (*BMJ* 2002;325:983)

Editor's choice

A good death, or a public one?

Many people might think that part of a good death is privacy—or at least having some say over how wide should be the circle of family and friends who come to talk and touch and give support. But these last two weeks have seen two very public deaths—with room to doubt that the victims would have wanted it that way.

Last weekend the pope died. Even in a non-Catholic country such as Britain, a casual listener to the radio or watcher of TV could not help notice how closely each stage of the pope's dying was marked and reported on. As Sophie Aire says in her media review (p 850), "No patient with Parkinson's disease has ever died such a public death." Almost 30 years ago, when General Franco, Spain's dictator, died, the media was criticised for covering every detail of his dying days. But, as Arie points out, it was an "unprecedented flow of Vatican bulletins that informed journalists that the pope was being fed by a nasogastric tube, had had a urinary tract infection, which was followed by septic shock... and finally, intermittent loss of consciousness." Though some observers didn't find such detail tasteful, Arie's account suggests that "the media friendly way in which the pope died was in character with the way he lived."

Terri Schiavo, on the other hand, died after a bitter argument between her husband and her parents and a politicised public and legal debate among the courts, the media, and US politicians (*BMJ* 2005;330:746). She died after appeal courts refused to overturn an earlier court ruling that she was in a persistent vegetative state and that she should not be kept alive against what were thought to be her wishes. Though the issue in the Schiavo case is not euthanasia as such, the case has inevitably reignited debates about euthanasia—and ferociously so in the US.

In the UK the language is more measured, but last week's report from a committee of the House of Lords marks a softening in attitudes towards euthanasia and assisted suicide in the UK (p 807). Although the committee is split over whether law should be changed, it does want a debate on the subject early in the next Parliament. It also wants euthanasia to be distinguished from assisted suicide, thinking that assisted suicide might be more acceptable to doctors.

Meanwhile, A C Grayling reminds us that humankind's attitudes towards dying are a bit muddled (p 799). He argues that the right to die has its moral basis in the right to a good quality of life. Choosing when and how to die is, he argues, one of life's great questions that should be left to individual choice—like whom to love and whether to have a family. The problems arise when people lose their power to exercise their choice, and didn't voice it beforehand: when they were not, as Montaigne would have it, "booted and spurred, and ready to go."

Jane Smith *deputy editor* (jsmith@bmj.com)

To receive Editor's choice by email each week subscribe via our website: bmj.com/cgi/customalert