Poor adherence to drug treatment increases mortality after a heart attack

Research question Does non-adherence to drug treatments affect survival after a heart attack?

Answer Yes. Patients who stop taking their aspirin, β blockers, and statins are more likely to die within a year than patients who continue to take at least one of them

Why did the authors do the study? They wanted to find out how many patients stopped taking their prescribed drugs after a heart attack, when they stopped, and whether this affected their survival. They also wanted to explore the factors associated with non-adherence to treatment.

What did they do? They followed up a cohort of 1521 US men and women who had been treated for myocardial infarction and then discharged from hospital taking aspirin, a β blocker, and a statin. The authors interviewed all the participants at baseline and then telephoned them at one, six, and 12 months after discharge to find out which drugs they were still taking. The authors also reviewed participants' medical records. Data on deaths came from the social security register of deaths coupled with telephone calls to relatives or the patient's primary care doctor or cardiologist.

The authors derived survival rates at one year for patients who stopped taking all three drugs within a month, and compared them with rates among patients who continued one or more of the treatments. They did further analyses to find out if non-adherence was an independent risk factor for death, and to explore which demographic or treatment characteristics made someone more likely to stop taking treatment.

What did they find? Of the 1521 patients, 184 (12.1%) stopped taking all three drugs within a month of leaving hospital. Another 328 (21.6%) stopped taking one or two, leaving only 1009 (66.3%) taking aspirin, a statin, and a β blocker. Those who stopped all three were significantly more likely to die within a year than those who continued taking one or more of their drugs (11.5% *v* 2.3%, $P < 0.001$), a difference that was independent of age, sex, treatment in hospital, psychosocial factors, and clinical history (adjusted hazard ratio for death 3.81 (95% CI 1.88 to 7.72)). In multivariable analyses, the only independent factors associated with stopping all three drugs were lack of high school education (odds ratio 1.76 (1.2 to 2.6)) and age. Age was a more powerful risk factor for women (odds ratio 1.77 (1.34 to 2.34) for each 10 year increase) than for men (1.23 (1.02 to 1.47)).

What does it mean? A substantial minority of patients stop taking their prescribed long term treatments after a heart attack, and this adversely affects their chances of survival either directly or indirectly. Most people who were going to had stopped their drugs within a month of leaving hospital. Socioeconomic factors seem more important than clinical factors in identifying those most at risk. These findings suggest that patients go through a vulnerable transition period after leaving hospital, and there may be a need to intensify follow-up to see them through it. The authors say a discussion about sticking with medication should be an essential part of the first outpatient consultation after a heart attack.

Ho et al. Impact of medication therapy discontinuation on mortality after myocardial infarction. *Arch Intern Med* 2006;166:1842-7

This summarises a paper that has been selected by bmjupdates. To register for bmjupdates (free email alerts about high quality new papers in your favourite subjects) go to <http://bmjupdates.com/>

Editor's choice

Concerns about revalidation

This week brings an end to the consultation period on Liam Donaldson's report on professional regulation in the United Kingdom. There's been no shortage of comments on the report (this week's *BMJ* carries some more on p 965), with emotions running high on many issues. How should Donaldson respond?

On the face of it his recommendations represent a huge challenge to the profession—an end to professional self regulation with loss of elected representation, removal of the General Medical Council's control of undergraduate education (which effectively excludes it from setting standards at entry to the profession), and the setting up of an independent tribunal to take on final adjudication in serious cases of fitness to practise.

Voices raised against Donaldson's proposals, many of them in the *BMJ*'s rapid responses, decry the planned move to a civil burden of proof in cases of professional misconduct and to appoint local GMC affiliates. They ask who will pay and how much it will cost in terms of time away from patients. They ask what will happen to those who fail and whether large numbers of older doctors will take early retirement rather than go through validation. And underlying all these concerns is the ultimate question: will Donaldson's proposals really deliver better care of patients?

But as Mike Pringle said when the report was first published (*BMJ* 2006;333:161-2) there is much to admire in Donaldson's recommendations. He brings important clarity to the distinction between relicensure and recertification. In so doing he has created a proper role for the royal colleges, which some would say they have lacked in the past, and which Ian Gilmore, president of the Royal College of Physicians, welcomes (p 696).

Donaldson also gives the GMC a much needed kick, in threatening to remove so much of its power. Last week, Jonathan Gornall laid blame for what he saw as Donaldson's heavy handed intervention at the door of the GMC, for its failure to seize the initiative in the wake of Janet Smith's highly critical report (*BMJ* 2006;333:906-11). To his credit the GMC's president, Graham Catto, seems at last to have responded. Catto has outdone Donaldson on proposals to create a less medically dominated council and, as Jim Johnson agrees (p 966), has made a robust defence of the GMC's role in education, with a three pronged proposal to oversee undergraduate, postgraduate, and continuing medical education.

Given Donaldson's mandate for change, even those most opposed to his proposals must accept that revalidation is now inevitable. Better still would be to embrace it as part of, rather than a threat to, medical professionalism. For his part, the chief medical officer needs to bring the UK's doctors with him rather than alienate them further, especially at a time when they already feel their professionalism is under attack from growing political interference. To do this he will need to show, in his revised recommendations, that he has heard their concerns.

Fiona Godlee *editor* (fgodlee@bmj.com)