



ANALYSIS

Fetal exposure to dietary carcinogens and risk of childhood cancer: what the NewGeneris project tells us

Jos Kleinjans and colleagues summarise the evidence from a European study using biomarkers to assess how maternal diet affects the child

Jos Kleinjans *professor of environmental health science*¹, Maria Botsivali *research associate*², Manolis Kogevinas *professor*³, Domenico Franco Merlo *senior epidemiologist*⁴, on behalf of the NewGeneris consortium

¹Department of Toxicogenomics, Maastricht University, Maastricht, Netherlands; ²Institute of Biology, Medicinal Chemistry and Biotechnology, National Hellenic Research Foundation, Athens, Greece; ³Center for Research in Environmental Epidemiology (CREAL), Barcelona, Spain; ⁴IRCCS AOU San Martino-IST- Istituto Nazionale per la Ricerca sul Cancro, Genova, Italy

Cancer in childhood is rare. Globally, there are around 175 000 new cancer cases a year among children aged 0-14 years.¹ However, in Europe, since the 1950s the incidence of cancer in this age group has increased by about 1% a year, with leukaemia, brain tumours, and lymphomas accounting for most cases.² The increases in incidence of lymphoid leukaemia, in particular, are more apparent in European than in Asian or African children.³ The development of childhood cancer thus seems to be affected by both genetic and environmental factors. Given that the highest incidences of childhood leukaemia are reported in children younger than 6-7 years, that the latency period of leukaemia in children is relatively short,⁴ and that adverse genetic events in utero have been shown to give rise to leukaemia in childhood,⁵ we hypothesised that fetal exposure to environmental carcinogens may be an underlying cause. Diet is an important source of carcinogenic compounds because of the accumulation of environmental carcinogens within the food chain (dioxins, polychlorinated biphenyls (PCBs), polycyclic aromatic hydrocarbons (PAHs)), as well as of formation of carcinogens such as PAHs, heterocyclic amines, and acrylamide during baking, frying, and grilling of food.⁶ The New Generis (Newborns and Genotoxic Exposure Risks) project therefore set out to investigate whether intake of dietary carcinogens by the pregnant mother leads to exposures of the fetus and initiates adverse biological responses that can induce cancer in later childhood.

Investigation of this hypothesis in an epidemiological study would require a huge sample size because of the relatively low incidence of childhood cancer (about 140 cases per million children).^{2,3} Case-control studies have been informative but also have limitations.⁷ Instead, we designed a biomarker based study.⁸ We used umbilical cord blood samples from 1151 newborns

from mother-child birth cohorts and biobanks in Norway, Denmark, UK, Spain, and Crete to investigate biochemical and cytogenetic markers indicative of prenatal exposure to chemical carcinogens and longer term cancer risk. We focused on fetal exposure to the following dietary carcinogens: acrylamide, PAHs, nitrosamines, fat oxidation metabolites (malondialdehyde), and xenoestrogens (dioxins and dioxin-like PCBs). Here, we summarise the major findings.

Do dietary carcinogens pass the placenta?

For fetal exposure to occur, dietary carcinogens taken in by the mother have to pass through the placenta. We investigated this using perfusion methods on placentas donated by 93 healthy mothers without any reported complications of pregnancy.⁹ Although the transfer rates of the different compounds varied, all tested carcinogens readily crossed the placental barrier. Nitrosamines—for example, nitrosodimethylamine—passed through most rapidly, followed by acrylamide and PAHs such as benzo(a)pyrene, with PCB 52 and dioxin being the slowest.⁹ However, since placental morphology profoundly changes throughout gestation, data on carcinogen transfer rates from term placenta may not be representative of transfer occurring earlier in gestation.¹⁰ We concluded that the placenta does not protect the fetus against exposure to dietary carcinogens.

Is the fetus actually exposed to dietary carcinogens?

We quantified the presence of selected dietary carcinogens in umbilical cord blood samples taken from 1151 newborns. For nitrosamines, acrylamide, oxidative fat metabolites, and PAHs

we assessed exposure by measuring levels of reactive metabolites bound to macromolecules such as DNA and haemoglobin. Plasma levels of dioxins and dioxin-like PCBs were measured with a validated bioassay (DR CALUX). The table^{||} shows the levels of these biomarkers of exposure in cord blood. In most infants, fetal exposure was confirmed. We noticed large variations in all biomarkers, with a skewed distribution of exposure across newborns (for instance, acrylamide-haemoglobin adducts ranged from 4.4 to 124.8 pmol/g Hb).¹¹

Does fetal exposure induce potentially carcinogenic events?

To assess the cancer risks potentially associated with fetal exposures to selected dietary carcinogens, we measured the numbers of micronuclei in lymphocytes from 467 umbilical cord blood samples. Micronuclei are a cytogenetic biomarker for chromosomal damage and have been prospectively associated with cancer risks in adults occupationally exposed to carcinogenic compounds.¹⁵ We found significant associations between particular features of micronuclei and exposure to oxidative fat metabolites and dioxins and PCBs, predominantly in the highest quarters of exposure.¹¹

To investigate further the molecular mechanisms underlying carcinogenic events in utero we determined global gene expression levels in 120 Norwegian neonates in relation to fetal exposure to acrylamide and dioxins. While exposure did not seem to differ between the sexes, we discovered important sex-specific differences in genomic responses, in particular associated with effects on cell cycle and the immune system.¹⁶ Specifically in boys, dioxin exposure was associated with activation of the proinflammatory transcription regulator tumour necrosis factor α , which then induces the nuclear factor- κ B; acrylamide exposure was associated with activation of the Wnt pathway, which is associated with uncontrolled cell growth. This may explain the higher incidence of leukaemia² and cancer overall among boys.³

Subsequently, from these gene expression profiles, we selected 36 genes and developed a high throughput screen based on polymerase chain reaction technology. This allowed us to evaluate the associations of these genes with established fetal exposures in all 1151 newborns for whom we had umbilical cord blood. Compared with the lowest exposure groups, expression of five genes was significantly higher in those with highest exposure to polycyclic aromatic hydrocarbons and six genes for those with highest exposure to fat metabolites. Expression levels were significantly lower for seven genes in those with highest exposure to dioxins compared with the lowest category.¹¹ The affected genes have roles in cell cycle regulation and apoptosis, which are generic processes involved in carcinogenesis.

Overall, fetal exposure to selected dietary carcinogens seems to induce molecular events that indicate increased cancer risks, thus supporting the hypothesis that childhood cancer, in particular leukaemia among boys, is causally related to the dietary intake of carcinogenic substances by their mothers during pregnancy.

Are particular children more susceptible to cancer risks?

Childhood cancer remains uncommon despite widespread exposure to dietary carcinogens in utero. This suggests that particular children may be more susceptible to the cancerous

effects of toxic chemicals.¹⁷ Such increased susceptibility may be genetically predisposed—some people may lack the gene activity for, say, detoxifying carcinogens or repairing induced DNA damage, or express a higher activity in metabolic transformation of carcinogens into DNA damaging daughter compounds. Relevant genes may be polymorphic in the general population, being non-homogeneously distributed at frequencies of 5–40%. Carriers of such single nucleotide polymorphisms (SNPs) may be prone to increased cancer risks after exposure to carcinogenic agents.¹⁸

To investigate whether genetic predisposition increased susceptibility to fetal exposure to carcinogens, we analysed cord blood DNA from 435 subjects using a genome-wide association method.¹⁹ We found a strong signal that the folate hydrolase 1 (*FOLH1*) gene interacted with the association between fetal exposure to dioxin related androgens and frequencies of micronuclei. The *FOLH1* polymorphism may cause low serum levels of folate thus leading to chromosome instability and formation of micronuclei. In addition, polymorphisms in the epoxyhydrolase 1/2 (*EPHX1/2*) and cytochrome p450 2E1 (*CYP2E1*) genes coding for metabolic enzymes, which may bioactivate precarcinogenic chemicals, seem to be associated with the micronuclei frequencies.¹¹

In conclusion, although the evidence is still limited, our data suggest some genetic predisposition to risk of childhood cancer associated with fetal exposure to dietary carcinogens.

Are there other effects on health?

We also investigated the effect of fetal exposure to dietary carcinogens on birth weight, head circumference, and gestational age. Fetal exposure to acrylamide seemed associated with a reduction in birth weight and head circumference ($n=1101$).²⁰ Biomarkers of exposure to PAHs ($n=612$) were associated with lower birth weight only in newborns from Norway, Denmark, and England.²¹ In addition, birth weight was lower in infants in the medium and high thirds of dioxin and dioxin-like PCB levels in cord blood ($n=967$) compared with those in the lowest third.¹² Furthermore, gestational age was shorter by about half a week in the highest compared with the lowest exposure levels; this association seemed to be stronger for boys than for girls. Reduced birth weight is regarded as a risk factor for cardiovascular disease, type 2 diabetes, and osteoporosis in later life,²² while reduced head circumference at birth may be related to delayed neurodevelopment.²³

Furthermore, we showed that maternal intake of dioxin and PCBs estimated from food frequency questionnaires is associated with immunotoxic events during early childhood. In 111 newborns from the Norwegian cohort, dioxin exposure was associated with gene expression responses in cord blood cells and with reduced measles vaccination responses at age 3 years.²⁴ Overall, the functions of affected genes—for instance, those belonging to the HLA system related to antibody production²⁵—pointed towards immunosuppression, girls showing more immune related genes deregulated in association with dioxin and PCB exposure than boys.

Based on the above results, fetal exposure to dietary carcinogens may have other adverse effects on health in addition to cancer risk.

Can pregnant woman reduce their risk of exposure?

The NewGeneris project suggests that pregnant women's consumption of carcinogens present in the European diet results

in measurable fetal exposures. Furthermore, such exposure seems to result in molecular and functional changes that are thought to be biomarkers for the risk of developing cancer and other diseases during childhood and later life. We observed higher susceptibilities among boys, as well as among carriers of particular gene polymorphisms, suggesting that genetic predisposition may contribute to health risks. Such risks might be reduced by decreasing the intake of dietary carcinogens. Although pregnant women can limit consumption by avoiding foods such as processed meats and fried and grilled food, eliminating carcinogens altogether is difficult because of their ubiquity in the diet.

Food manufacturers could have a role in reducing the presence of carcinogenic agents. For instance, manufacturers of French fries and crisps could switch to low sugar potato varieties to reduce the presence of acrylamide, which is formed during deep frying by the reaction of asparagine and reducing sugars (both naturally occurring in potatoes). Reduction of use of nitrite for curing meat products would also lead to less formation of nitrosamines in the stomach. For carcinogens that are present in multiple types of food, such as dioxin-like PCBs which are generically present in meat and fish as well as in milk and milk products,¹³ food manufacturers could be required to assess the levels of these compounds in raw products and take preventive measures. We propose that national food safety agencies consider exposure risks in more detail and implement policy measures where appropriate.

In general, our findings support the recommendation to pregnant mothers to eat a well balanced diet. Such a diet also has benefits for the child by reducing the risks of preterm delivery,²⁶ low birth weight,²⁷ childhood leukaemia,⁷ and adult cardiovascular disease.²⁸ It should also not be forgotten that smoking and alcohol intake during pregnancy pose the largest health risks to the unborn child.

NewGeneris is an integrated project conducted within the European Union's 6th Framework Programme, priority area Food Quality and Safety. We thank Soterios Kyrtopoulos from the National Hellenic Research Foundation for critically reading the manuscript.

Contributors and sources: JK coordinated the NewGeneris project, supervised the analysis of biomarkers of genotoxic exposure and effects, and wrote this summary. MB calculated maternal dietary intake of relevant carcinogens and identified the main food items contributing to these intakes. MK supervised the analysis of additional health outcomes and contributed to the calculations of dietary intake of carcinogens. DFM was responsible for the data analysis plan and contributed to the analysis of the GWAS data.

Competing interests: We have read and understood BMJ policy on declaration of interests and have no relevant interests to declare.

Provenance and peer review: Not commissioned; externally peer reviewed.

- 1 American Cancer Society. Global cancer facts and figures. 2nd ed. American Cancer Society, 2011.
- 2 Kaatsch P. Epidemiology of childhood cancer. *Cancer Treat Rev* 2010;36:277-85.
- 3 Steliarova-Foucher E, Stiller C, Kaatsch P, et al. Geographical patterns and time trends of cancer incidence and survival among children and adolescents in Europe since the 1970s (the ACCIS project): an epidemiological study. *Lancet* 2004;364:2097-105.
- 4 Greaves MF. Aetiology of acute leukaemia. *Lancet* 1997;349:344-9.
- 5 Wiemels JL, Xiao Z, Buffler PA, et al. In utero origin of t(8;21) AML1-ETO translocations in childhood acute myeloid leukemia. *Blood* 2002;99:3801-5.
- 6 Sugimura T. Nutrition and dietary carcinogens. *Carcinogenesis* 2000;21:387-95.
- 7 Kwan ML, Jensen CD, Block G, Hudes ML, Chu LW, Buffler PA. Maternal diet and risk of childhood acute lymphoblastic leukemia. *Public Health Rep* 2009;124:503-14.
- 8 Merlo DF, Wild CP, Kogevinas M, et al. NewGeneris: a European study on maternal diet during pregnancy and child health. *Cancer Epidemiol Biomarkers Prev* 2009;18:5-10.
- 9 Mose T, Mathiesen L, Karttunen V, et al. Meta-analysis of data from human ex vivo placental perfusion studies on genotoxic and immunotoxic agents within the integrated European project NewGeneris. *Placenta* 2012;33:433-9.
- 10 Nanovskaya TN, Nekhayeva IA, Hankins GD, Ahmed MS. Transfer of methadone across the dually perfused preterm human placental lobule. *Am J Obstet Gynecol* 2008;198:126.e1-4.
- 11 Merlo DF, Agramunt S, Anna L, et al. Micronuclei in cord blood lymphocytes and associations with biomarkers of exposure to carcinogens and hormonally active factors, gene polymorphisms, and gene expression: the NewGeneris cohort. *Environ Health Perspect* 2014;122:193-200.
- 12 Vafeiadi M, Agramunt S, Pedersen M, et al. In utero exposure to compounds with dioxin-like activity and birth outcomes. *Epidemiol* 2014;25:215-24.
- 13 Papadopoulou E, Kogevinas M, Botsivali M, et al. Maternal diet, prenatal exposure to dioxin-like compounds and birth outcomes in a European prospective mother-child study (NewGeneris). *Sci Total Environ* 2014;484:121-8.
- 14 Van de Look K, Botsivali M, Zangogianni M, et al. The effect of dietary estimates calculated using food frequency questionnaires on micronuclei formation in European pregnant women: a NewGeneris study. *Mutagenesis* 2014;29:393-400.
- 15 Bonassi S, Znaor A, Ceppi M, et al. An increased micronucleus frequency in peripheral blood lymphocytes predicts the risk of cancer in humans. *Carcinogenesis* 2007;28:625-31.
- 16 Hochstenbach K, van Leeuwen DM, Gmuender H, et al. Global gene expression analysis in cord blood reveals gender-specific differences in response to carcinogenic exposure in utero. *Cancer Epidemiol Biomarkers Prev* 2012;21:1756-67.
- 17 Wild CP, Kleinjans J. Children and increased susceptibility to environmental carcinogens: evidence or empathy? *Cancer Epidemiol Biomarkers Prev* 2003;12:1389-94.
- 18 Williams JA. Single nucleotide polymorphisms, metabolic activation and environmental carcinogenesis: why molecular epidemiologists should think about enzyme expression. *Carcinogenesis* 2001;22:209-14.
- 19 Norrgard, K. Genetic variation and disease: GWAS. *Nat Educ* 2008;1:87-9.
- 20 Pedersen M, von Stedingk H, Botsivali M, et al. Birth weight, head circumference, and prenatal exposure to acrylamide from maternal diet: the European prospective mother-child study (NewGeneris). *Environ Health Perspect* 2012;120:1739-45.
- 21 Pedersen M, Schoket B, Godschalk RW, et al. Bulky DNA adducts in cord blood, maternal fruit-and-vegetable consumption, and birth weight in a European mother-child study (NewGeneris). *Environ Health Perspect* 2013;121:1200-6.
- 22 Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med* 2008;359:61-73.
- 23 Gale CR, O'Callaghan FJ, Bredow M, Martyn CN. The influence of head growth in fetal life, infancy, and childhood on intelligence at the ages of 4 and 8 years. *Pediatrics* 2006;118:1486-92.
- 24 Hochstenbach K, van Leeuwen DM, Gmuender H, et al. Toxicogenomic profiles in relation to maternal immunotoxic exposure and immune functionality in newborns. *Toxicol Sci* 2012;129:315-24.
- 25 Ovsyannikova IG, Jacobson RM, Vierkant RA, Shane Pankratz V, Jacobsen SJ, Poland GA. Associations between human leukocyte antigen (HLA) alleles and very high levels of measles antibody following vaccination. *Vaccine* 2004;22:1914-20.
- 26 Englund-Ögge L, Brantsæter AL, Sengpiel V, et al. Maternal dietary patterns and preterm delivery: results from large prospective cohort study. *BMJ* 2014;348:g1446.
- 27 Kind KL, Moore VM, Davies MJ. Diet around conception and during pregnancy-effects on fetal and neonatal outcomes. *Reprod Biomed Online* 2006;12:532-41.
- 28 Godfrey KM, Barker DJ. Fetal programming and adult health. *Public Health Nutr* 2001;4:611-24.

Accepted: 21 Jul 2015

Cite this as: *BMJ* 2015;351:h4501

© BMJ Publishing Group Ltd 2015

Key messages

- An estimated 175 000 new cancer cases occur globally among children aged 0-14
- The fetus is exposed to dietary carcinogens consumed by the mother during pregnancy
- Exposure of the fetus causes perturbations of the molecular pathways linked with functional markers of carcinogenicity
- Genetic polymorphisms and male sex may predispose to susceptibility to carcinogen exposure before birth and increase risks of developing leukaemia, in particular
- Fetal exposure to dietary carcinogens may also affect birth weight and cause immune suppression

Table

Table 1 | Median (range) exposure to dietary (pre)carcinogens in newborns and maternal dietary intake during pregnancy

Carcinogen (biomarker)	Cord blood biomarker level	Maternal daily dietary intake*	Foods with high levels
Acrylamide (acrylamide-haemoglobin adducts)	14.4 (4.4-124.8) pmol/g Hb	22 (1-135) µg	Deep fried potato products (crisps, french fries)
			Crisp breads, biscuits, crackers
			Coffee
Polycyclic aromatic hydrocarbons (bulky DNA adducts)	8.4 (0.6-116.6) 10-8 nucleotides	180 (42-717) ng	Grilled/barbecued meats
			Smoked and processed fish
			Smoked and processed meat
Oxidative fat metabolites (M1dG-DNA adducts)	34.2 (0.5-324.7) 10-9 nucleotides	Omega 6 fatty acids 11.3 (1.2-77.0) g	Vegetable oils (soya, rape, sunflower)
			Fatty meat
			Eggs
Nitrosamines (O ⁶ -MG-DNA adducts)	0.40 (0.08-3.03) 10-8 nucleotides	N-nitrosodimethylamine 82 (2-547) ng	Processed meat
			Preserved fish
			Smoked meat
			Smoked fish
Dioxin/PCBs (plasma DR CALUX)	0.13 (0.01-104) pg/ml†	77.2 (5.1-945.5) pg†	Fatty meat
			Fatty fish
			Full fat milk and dairy products

*Derived from the NewGeneris database on dietary consumption patterns taken from food frequency questionnaires and data on chemical levels in food.^{12 13 14}
†2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) toxic equivalency quotients.