

Corticosteroids for sore throat: a clinical practice guideline

Bert Aertgeerts,^{1 2} Thomas Agoritsas,^{3 4} Reed A C Siemieniuk,^{3 5} Jako Burgers,^{6 7} Geertruida E Bekkering,^{1 2} Arnaud Merglen,⁸ Mieke van Driel,⁹ Mieke Vermandere,¹ Dominique Bullens,^{10 11} Patrick Mbah Okwen,¹² Ricardo Niño,¹³ Ann van den Bruel,^{14 15} Lyubov Lytvyn,¹⁶ Carla Berg-Nelson,^{17 18} Shunjie Chua,¹⁹ Jack Leahy,²⁰ Jennifer Raven,²¹ Michael Weinberg,²² Behnam Sadeghirad,^{3 23} Per O Vandvik,^{15 24} Romina Brignardello-Petersen^{2 25}

Full author details can be found at the end of the article

Correspondence to: B Aertgeerts bert.aertgeerts@kuleuven.be

Cite this as: *BMJ* 2017;358:j4090
doi: 10.1136/bmj.j4090

This BMJ Rapid Recommendation article is one of a series that provides clinicians with trustworthy recommendations for potentially practice changing evidence. BMJ Rapid Recommendations represent a collaborative effort between the MAGIC group (www.magicproject.org) and The BMJ. A summary is offered here and the full version including decision aids is on the MAGICapp (www.magicapp.org), for all devices in multilayered formats. Those reading and using these recommendations should consider individual patient circumstances, and their values and preferences and may want to use consultation decision aids in MAGICapp to facilitate shared decision making with patients. We encourage adaptation and contextualisation of our recommendations to local contexts. Those considering use or adaptation of content may go to MAGICapp to link or extract its content or contact The BMJ for permission to reuse content in this article.

What is the role of a single dose of oral corticosteroids for those with acute sore throat? Using the GRADE framework according to the BMJ Rapid Recommendation process, an expert panel make a weak recommendation in favour of corticosteroid use. The panel produced these recommendations based on a linked systematic review triggered by a large randomised trial published in April 2017. This trial reported that corticosteroids increased the proportion of patients with complete resolution of pain at 48 hours. Box 1 shows all of the articles and evidence linked in this Rapid Recommendation package. The infographic provides the recommendation together with an overview of the absolute benefits and harms of corticosteroids in the standard GRADE format. Table 2 below shows any evidence that has emerged since the publication of this article. Clinicians and their patients can find consultation decision aids to facilitate shared decision making in MAGICapp (www.magicapp.org/goto/guideline/JjXYAL/section/j79pvn).

Acute sore throat is defined as pain in the throat for less than 14 days. Acute sore throat could be caused by pharyngitis, nasopharyngitis, tonsillitis, peritonsillar abscess, or retropharyngeal abscess. Some patients with sore throat also experience headache, fever, muscle stiffness, cough, and general malaise.

Acute sore throat is common, but only a minority of patients will visit their general practitioner.¹ A survey reported that the main reasons are to establish the cause of the symptoms, obtain pain relief, and to gain information on the course of the disease.² Data from Dutch and Flemish primary care databases show that, for every 1000 consecutive patients consulting a general practitioner, 50 present with an acute sore throat.^{3 4} In the US, more than 92 million visits by adults to primary care practices and emergency departments between 1997 and 2010 were recorded.⁵ Sore throat presenting as acute tonsillitis is also the commonest cause for emergency admission to otorhinolaryngology services in the US.⁶

Acute sore throat is a self limiting disease and typically resolves after 7-10 days in adults and 2-7 days in children.⁷ Most infections are of viral origin; only a few are caused by a bacterial infection, of which group A

WHAT YOU NEED TO KNOW:


- Sore throat is one of the most common reasons for primary care appointments, and international guidance varies about whether to use corticosteroids to treat it, but a trial published in April 2017 suggested that corticosteroids might be effective
- We make a weak recommendation to use a single dose of oral corticosteroids, in those presenting with acute sore throat, after performing a systematic review of the new evidence in this rapid recommendation publication package
- The recommendation is weak and shared decision making is needed because corticosteroids did not help all patient reported outcomes and patients' preferences varied substantially
- Steroids somewhat reduced the severity and duration of pain by one day, but time off school or work was unchanged. Harm seems unlikely with one steroid dose.
- The treatment is inexpensive and likely to be offered in the context of a consultation that would have taken place anyway

β -haemolytic streptococcus, *Haemophilus influenzae*, and *Moraxella catarrhalis* are the most common pathogens. Evidence suggests that the time to resolution is not associated with the type of pathogen.⁷ About 2% of patients initially presenting with sore throat will have a mononucleosis infection caused by an Epstein-Barr virus, which could prolong the duration of symptoms.⁸

Some patients experience unacceptable morbidity and inconvenience, and miss school or work due to recurrent sore throat.⁹ Pain is a common reason for work or school absence. Complications of sore throat are rare: about 0.2% of patients with tonsillitis will develop a peritonsillar abscess.¹⁰

The diagnosis of an acute sore throat is based on signs and symptoms. The Centor clinical prediction rules can be used to help predict whether the sore throat is caused by a bacterial pathogen, and thus guide the decision whether to prescribe an antibiotic.^{11 12}

Population



People with sore throat

This recommendation applies to almost all patients with sore throat:

- ✓ Children 5 years and older and all adults
- ✓ Severe and not severe sore throat
- ✓ Emergency and primary care settings
- ✓ Patients with a viral or bacterial sore throat
- ✓ Patients who receive immediate or deferred antibiotics

However the recommendation is **not** applicable to patients with:

- ✗ Infectious mononucleosis
- ✗ Immunocompromising conditions
- ✗ Sore throat following surgery or intubation
- ✗ Children under 5 years old

Comparison

Short course of steroids

1–2 doses of oral Dexamethasone (or equivalent dose of alternative corticosteroid) + standard care

Adults:
10 mg

Children:
0.6 mg
per kg

+

Standard care
+ Analgesics
+/- Antibiotics

Favours steroids

or

No steroids

Standard clinical care, which typically includes analgesics, and may include antibiotics

Standard care
+ Analgesics
+/- Antibiotics

Favours no steroids

Strong ← Weak → Weak → Strong

We suggest short course of steroids. Discuss with patients in shared decision making.

Preferences and values

The panel believes that there is a great variability on how much reduction in pain severity or time to complete pain resolution each patient would consider important. Shared decision making may help establish what matters most to each patient.

Serious adverse events

One-dose administration of steroids is not likely to cause serious adverse events. Very low quality evidence exists for extremely rare but serious adverse effects following higher doses or longer courses of steroids (up to 30 days).

Multiple doses

Risks may outweigh benefits when cumulative doses of steroids are given for multiple episodes of sore throat. To mitigate this issue, clinicians could administer the medication in office if possible, or prescribe only one dose per visit.

Disclaimer: This infographic is not a validated clinical decision aid. This information is provided without any representations, conditions or warranties that it is accurate or up to date. BMJ and its licensors assume no responsibility for any aspect of treatment administered with the aid of this information. Any reliance placed on this information is strictly at the user's own risk. For the full disclaimer wording see BMJ's terms and conditions: <http://www.bmj.com/company/legal-information/>

BMJ: first published as 10.1136/bmj.j4090 on 20 September 2017. Downloaded from <http://www.bmj.com/> on 21 May 2024 by guest. Protected by copyright.

Box 1 | Linked articles in this *BMJ* Rapid Recommendations cluster

- Aertgeerts B, Agoritsas T, Siemieniuk RAC, et al. Corticosteroids for sore throat: a clinical practice guideline. *BMJ* 2017;358:j4090. doi:10.1136/bmj.j4090
– Summary of the results from the Rapid Recommendation process
- Sadeghirad B, Siemieniuk RA, Brignardello-Petersen R, et al. Corticosteroids for treatment of sore throat: a systematic review and meta-analysis of randomised trials. *BMJ* 2017;358:j3887. doi:10.1136/bmj.j3887
– Review of all available randomised trials that assessed corticosteroids as adjunct treatment versus standard care for sore throat.
- MAGICapp (www.magicapp.org/goto/guideline/jjXYAL/section/j79pvn)
– Expanded version of the results with multilayered recommendations, evidence summaries, and decision aids for use on all devices

Most guidelines recommend paracetamol or ibuprofen as the first choice treatment.¹³ The use of corticosteroids is mentioned in few, and is generally discouraged (table 1). Antibiotics are probably not helpful for pain relief in an episode of acute sore throat caused by viruses, but may help those with a bacterial infection.^{14 15} Recommended management of sore throat varies widely, and table 1 summarises current guidelines.

The evidence

The linked systematic review reports the effects of corticosteroids when added to standard care in patients with acute sore throat.¹⁶

Figure 1 gives an overview of the number and types of patients included, the study funding, and patient involvement, as well as a summary of the benefits and harms of corticosteroids for treating acute sore throat.

The panel identified eight patient-important outcomes needed to inform the recommendation: complete resolution of pain, time to onset of pain relief, pain severity, need for antibiotics, days missed from school or work, recurrence of symptoms, duration of bad or non-tolerable symptoms, and adverse effects. The included studies reported on all patient-important outcomes, except for duration of bad or non-tolerable symptoms. Regarding pain, the panel appraised the likelihood of complete resolution of pain at 24 hours and 48 hours, as well as the mean time to complete resolution of pain and the mean time to onset of pain relief.

Although most of the studies (80%) were conducted in emergency departments, they accounted for 54% of all patients enrolled across studies. The remaining 46% were enrolled in the studies conducted in primary care

Table 1 | Current guidance for treatment of patients with sore throat

	Ibuprofen	Paracetamol	Antibiotics	Corticosteroids	
				For adults	For children
EBM guidelines ¹¹	Supportive	Supportive	Conditionally	Supportive	Not applicable
SIGN ⁶	Supportive	Supportive	Conditionally	Not supportive	No comment
NHG ¹²	Supportive	Supportive	Conditionally	Not recommended	No comment
BC guidelines ¹³	No comment	No comment	Against	No comment	No comment
UpToDate ¹⁴	Against	No comment	No comment	Supportive	No comment

HOW THE RECOMMENDATION WAS CREATED

A large randomised controlled trial published in April 2017²¹ found that corticosteroids increased the proportion of patients with complete resolution of symptoms at 48 hours. However, corticosteroids did not seem to decrease the duration of moderately bad symptoms, pain severity, healthcare attendance, days missed from school or work, or the consumption of delayed antibiotics. This study adds to the body of evidence that suggests that, although corticosteroids probably have benefits in patients with sore throat, these benefits may be modest.²²⁻²⁵ The Rapid Recommendations team felt that the study, when considered in context of the full body of evidence, might change practice.²⁶

Our international panel—including general practitioners, general internists, paediatricians, an otorhinolaryngologist, epidemiologists, methodologists, statisticians, and people with lived experience of sore throat—decided what was the scope of the recommendation and the outcomes that are most important to patients. After a parallel team conducted a systematic review on the benefits and harms of corticosteroids,¹⁶ and a systematic search for evidence about patients' values and preferences (appendix 1 on bmj.com), the panel met to discuss the evidence and formulate a recommendation. No person had financial conflicts of interest; intellectual and professional conflicts were minimised and managed (appendix 2 on bmj.com).

The panel followed the *BMJ* Rapid Recommendations procedures for creating a trustworthy recommendation,^{26 27} including using the GRADE approach to critically appraise the evidence and create recommendations (appendix 3 on bmj.com).²⁸ The panel considered the balance of benefits, harms, and burdens of the drug, the quality of the evidence for each outcome, typical and expected variations in patient values and preferences, and acceptability.²⁹ Recommendations can be strong or weak, for or against a course of action.

settings, and the panel was therefore confident that the evidence was applicable to them as well. Most of the studies focused in adults only (60%). The studies that focused only on children (three studies, 2% of all the patients enrolled in the studies) did not include children younger than 5 years old, and thus the recommendation does not apply to younger ages.

Since the randomised controlled trials focused on patients who did not have recurrent episodes of sore throat, the panel was less confident of the applicability of the evidence to such patients, and the recommendation therefore does not apply to them. Similarly, the panel did not consider patients with sore throat after surgery or intubation, nor immunocompromised patients.

Understanding the recommendation

The recommendation for using corticosteroids made by the panel was weak because of the modest reduction of symptoms and the large variability in patient preferences.

The panel is confident that the recommendation applies to almost all patients with acute sore throat: children 5 years and older and adults, severe and not severe sore throat, patients who receive immediate antibiotics and those who receive deferred antibiotics, patients with a viral or bacterial sore throat, and patients who seek




Fig 1 | Characteristics of patients and trials included in systematic review of effects of corticosteroids on acute sore throat

care in the emergency department as well as those who attend primary care. The systematic review contained adequate representation from such groups and settings, and results were consistent (that is, absence of credible subgroup effects), for example, between trials of children and adults, and those seen in emergency departments and in primary care offices.¹⁶

Absolute benefits and harms

Although the evidence indicates that the treatment works on average, it did not reduce the severity of pain dramatically and failed to improve several other patient-important outcomes.

The infographic explains the recommendation and provides an overview (GRADE summary of findings) of the absolute benefits and harms of corticosteroids. Estimates of baseline risk for effects come from the control arms of the trials.¹⁶ The infographic also leads to point-of-care formats in the MAGICapp, including consultation decision aids designed to support shared decision making with patients.¹⁷

Considering the evidence and its certainty, the panel was confident that:

- Corticosteroids increase the chance of complete resolution of pain at 24 and 48 hours, reduce the severity of pain, and shorten the time to onset of pain relief (GRADE high to moderate quality evidence)
- Corticosteroids are unlikely to reduce recurrence or relapse of symptoms or days missed from school or work (GRADE moderate quality evidence)

- A single dose of corticosteroids is unlikely to cause serious adverse events
 - The randomised trials did not report any major event attributable to single dose corticosteroids (GRADE moderate quality evidence)
 - The panel also considered evidence from observational studies that used higher doses of steroids. A large retrospective US cohort study of private insurance claims assessed adverse events in 327 452 adults who received an outpatient prescription of corticosteroids.¹⁸ There was a small absolute increase in the rate of sepsis, venous thromboembolism, and fracture in the first 30 days (GRADE low quality evidence, due to suboptimal verification of diagnosis in large databases and confounding by indication¹⁹). The panel agreed that such events seemed unlikely with single dose steroids
 - Similarly, among paediatric populations, indirect evidence from a meta-analysis of 44 randomised trials did not report any major adverse events in patients with conditions requiring a short course of corticosteroids (such as asthma, bronchiolitis, croup, wheeze, and pharyngitis or tonsillitis)²⁰
- There are no differences in the relative effects of corticosteroids (when compared with usual care) between primary care settings and emergency departments
- It is unlikely that new information will change interpretation for outcomes that are high to moderate quality of evidence.

PRACTICAL ISSUES		
	Steroids	No steroids
MEDICATION ROUTINE	<ul style="list-style-type: none"> One (or two) doses of steroids, taken as pill(s) or intramuscular injection(s) May require concomitant antibiotics, and/or over the counter pain relievers 	
TESTS & VISITS	<ul style="list-style-type: none"> May need additional visits if symptoms do not resolve or worsen 	
ADVERSE EFFECTS	<ul style="list-style-type: none"> Serious adverse events are unlikely with one-dose steroids. There may be risks with repeated doses across multiple episodes of sore throat, or through self-medication May require concomitant antibiotics, and or over the counter pain relievers 	
EMOTIONAL WELL-BEING	<ul style="list-style-type: none"> May cause transient sleep disturbance and excitability, although infrequently with one-dose steroids 	
PREGNANCY & NURSING	<ul style="list-style-type: none"> Dexamethasone crosses the placenta, and is generally avoided during pregnancy. There is, however, probably no risk of malformation 	
COSTS & ACCESS	<ul style="list-style-type: none"> Inexpensive, available by prescription 	
FOOD & DRINK	<ul style="list-style-type: none"> May increase appetite, particularly in children 	

Fig2 Practical issues about use of corticosteroids to treat acute sore throat

- The panel was less confident about whether:
- Corticosteroids reduced antibiotic use, due to a lack of improvement or worsening of symptoms in patients not prescribed antibiotics immediately when consulting the physician (GRADE low quality evidence)
 - Corticosteroids reduced the average time to complete resolution of pain (GRADE low quality evidence).

Values and preferences

The weak recommendation for corticosteroids reflects a high value on a modest reduction of symptom severity and the time that it takes to achieve such improvement, and a substantial and important increase in the chance of complete resolution of pain at 48 hours.

The panel, including the patient representatives, felt that the values and preferences are likely to vary greatly across patients, which justifies a weak recommendation. For example, achieving complete pain resolution 12 hours earlier may be of little importance for patients who feel less busy in their daily life, have higher tolerance to pain, or whose symptoms are not so severe; whereas it may be very important to patients whose ability to go

EDUCATION IN PRACTICE

- How do you currently approach giving advice for those with acute sore throat? Do you consider offering corticosteroids?
- The recommendation for corticosteroid use is weak, and patient’s preferences are likely to vary. What information could you share with your patient to help reach a decision together?
- Have you learnt one thing from this article that might alter how you consult with patients with sore throat? How might you share this information with colleagues to learn together?
- To what extent do you practice shared decision making for such preference-sensitive decisions?

HOW PATIENTS WERE INVOLVED IN THE CREATION OF THIS ARTICLE

Five people with lived experience of sore throat were full panel members. These panel members identified important outcomes, and led the discussion on values and preferences. These patient representatives agreed that while small reductions in pain severity and time to complete pain resolution (for example 12 compared to 24 hours) were important to them, these values may not be shared by all patients; they expected moderate to great variability in how much importance other patients would place in small reductions in pain. These panel members participated in the teleconferences and email discussions and met all authorship criteria.

to school or to perform at work are compromised, care givers wishing to reduce their children’s pain, or patients experiencing their pain as severe.

The panel believes that there is great variability in how much reduction in pain severity or time to complete pain resolution each patient would consider important. However, the greater the reduction in hours to achieve complete resolution of pain, the more likely it is that typical patients would place high value on those outcomes. Patients who place a high value in reducing the symptoms by any amount (such as patients with lower tolerance to pain or with severe symptoms) are more likely to accept receiving corticosteroids.

The weak recommendation for corticosteroids also reflects the concerns that the panel had with acceptability. Specifically, how acceptable is it to treat a condition that is usually not severe and is self limiting with a drug that many patients, practitioners, and other stakeholders know is almost always used for more severe diseases.

The systematic search for empirical data on patients’ values and preferences related to sore throat identified 4149 references that were screened at the title and abstract level. From these, we screened 99 full text articles, from which only two provided relevant information on patients’ values and preferences (see appendix 1 on bmj.com). Neither of the studies provided additional data that had not been raised by the panel members: the panel had identified appropriate patient-important outcomes and considered the variability in patient values and preferences regarding sore throat management.

Table 2 | New evidence which has emerged after initial publication

Date	New evidence	Citation	Findings	Implications for recommendation(s)
There are currently no updates to the article				

Practical issues and other considerations

Figure 2 outlines the key practical issues for patients and clinicians discussing adjunct steroids for sore throat, which are also accessible along with the evidence as decision aids to support shared decision-making in MAGI-Capp. Steroids are typically given as 10 mg dexamethasone (or adapted to weight for children: 0.6 mg/kg, up to a maximum dose of 10 mg), typically taken as pill or intramuscular injection.

The risks may outweigh the benefits when larger cumulative doses of corticosteroids are given to patients who experience multiple episodes of sore throat, either through multiple visits or for patients who self medicate if prescribed more than one pill for their previous episode. To mitigate this issue, clinicians should administer the medication in office if possible or prescribe only one dose per visit.

Costs and resources

The panel focused on the patient perspective rather than that of society when formulating the recommendation. Given the low cost of corticosteroids for treating sore throat, implementation of this recommendation is unlikely to have an important impact on the costs for health funders. The treatment is inexpensive and likely to be offered in the context of a consultation that would have taken place anyway. Nevertheless, it remains uncertain whether it may increase the proportion of patients visiting a doctor to get a prescription of corticosteroids.

Uncertainties for future research

Key research questions to inform decision makers and future guidelines include:

- Are there any severe adverse effects of using one-dose of steroids for treating sore throat?
- What are the effects of corticosteroids, in addition to standard care, in patients with recurrent episodes of acute sore throat?

Updates to this article

Table 2 shows evidence which has emerged since the publication of this article. As new evidence is published, a group will assess the new evidence and make a judgment on to what extent it is expected to alter the recommendation.

Competing interests: All authors have completed the *BMJ Rapid Recommendations* interests disclosure form and a detailed, contextualised description of all disclosures is reported in appendix 2 on bmj.com. As with all *BMJ Rapid Recommendations*, the executive team and *The BMJ* judged that no panel member had any financial conflict of interest. Professional and academic interests are minimised as much as possible, while maintaining necessary expertise on the panel to make fully informed decisions.

Funding: This guideline was not funded.

Transparency: B Aertgeerts affirms that the manuscript is an honest, accurate, and transparent account of the recommendation being reported; that no important aspects of the recommendation have been omitted; and that any discrepancies from the recommendation as planned (and, if relevant, registered) have been explained.

- Evans CE, McFarlane AH, Norman GR, Neale KA, Streiner DL. Sore throats in adults: who sees a doctor? *Can Fam Physician* 1982;28:453-8. PMID:21286075.
- van Driel ML, De Sutter A, Deveugele M, et al. Are sore throat patients who hope for antibiotics actually asking for pain relief? *Ann Fam Med* 2006;4:494-9. doi:10.1370/afm.609 PMID:17148626.
- NIVEL Primary care database: Netherlands Institute for Health Services Research; 2012. www.nivel.nl/en/dossier/nivel-primary-care-database.
- Truyers C, Goderis G, Dewitte H, Akker Mv, Buntinx F. The Intego database: background, methods and basic results of a Flemish general practice-based continuous morbidity registration project. *BMC Med Inform Decis Mak* 2014;14:48. doi:10.1186/1472-6947-14-48 PMID:24906941.
- Barnett ML, Linder JA. Antibiotic prescribing to adults with sore throat in the United States, 1997-2010. *JAMA Intern Med* 2014;174:138-40. doi:10.1001/jamainternmed.2013.11673. PMID:24091806.
- Bird JH, Biggs TC, King EV. Controversies in the management of acute tonsillitis: an evidence-based review. *Clin Otolaryngol* 2014;39:368-74. doi:10.1111/coa.12299. PMID:25418818.
- Thompson M, Vodicka TA, Blair PS, Buckley DI, Heneghan C, Hay AD. TARGET Programme Team. Duration of symptoms of respiratory tract infections in children: systematic review. *BMJ* 2013;347:f7027. doi:10.1136/bmj.f7027. PMID:24335668.
- Rea TD, Russo JE, Katon W, Ashley RL, Buchwald DS. Prospective study of the natural history of infectious mononucleosis caused by Epstein-Barr virus. *J Am Board Fam Pract* 2001;14:234-42. PMID:11458965.
- Scottish Intercollegiate Guidelines Network. Management of sore throat and indications for tonsillectomy (SIGN 117). 2010. www.sign.ac.uk/sign-117-management-of-sore-throat-and-indications-for-tonsillectomy.html.
- Little P, Stuart B, Hobbs FD, et al. DESCARTE investigators. Predictors of suppurative complications for acute sore throat in primary care: prospective clinical cohort study. *BMJ* 2013;347:f6867. doi:10.1136/bmj.f6867 PMID:24277339.
- Atula T. Corticosteroids for sore throat. *Evidence-Based Medicine Guidelines*. 2014. www.ebm-guidelines.com/ebmg/ltk.naytaartikkeli?p_artikkeli=evd07095#refs.
- Ebell MH, Smith MA, Barry HC, Ives K, Carey M. The rational clinical examination. Does this patient have strep throat? *JAMA* 2000;284:2912-8. doi:10.1001/jama.284.22.2912 PMID:11147989.
- Chiappini E, Regoli M, Bonsignori F, et al. Analysis of different recommendations from international guidelines for the management of acute pharyngitis in adults and children. *Clin Ther* 2011;33:48-58. doi:10.1016/j.clinthera.2011.02.001. PMID:21397773.
- Del Mar CB, Glasziou PP, Spinks AB. Antibiotics for sore throat. *Cochrane Database Syst Rev* 2006;(4):CD000023. doi:10.1002/14651858.CD000023.pub3. PMID:17054126.
- van Driel ML, De Sutter A, Habraken H, Thorning S, Christiaens T. Different antibiotic treatments for group A streptococcal pharyngitis. *Cochrane Database Syst Rev* 2016;9:CD004406. doi:10.1002/14651858.CD004406.pub4. PMID:27614728.
- Sadhegirad B, Siemieniuk R, Brignardello-Petersen R, et al. Corticosteroids for treatment of sore throat: a systematic review and meta-analysis of randomized trials. *BMJ* 2017;358:j3887. doi:10.1136/bmj.j3887.
- Agoritsas T, Heen AF, Brandt L, et al. Decision aids that really promote shared decision making: the pace quickens. *BMJ* 2015;350:g7624. doi:10.1136/bmj.g7624. PMID:25670178.
- Waljee AK, Rogers MA, Lin P, et al. Short term use of oral corticosteroids and related harms among adults in the United States: population based cohort study. *BMJ* 2017;357:j1415. doi:10.1136/bmj.j1415. PMID:28404617.
- Agoritsas T, Merglen A, Shah ND, O'Donnell M, Guyatt GH. Adjusted analyses in studies addressing therapy and harm: users' guides to the medical literature. *JAMA* 2017;317:748-59. doi:10.1001/jama.2016.20029. PMID:28241362.
- Fernandes RM, Oleszczuk M, Woods CR, Rowe BH, Cates CJ, Hartling L. The Cochrane Library and safety of systemic corticosteroids for acute respiratory conditions in children: an overview of reviews. *Evid Based Child Health* 2014;9:733-47. doi:10.1002/ebch.1980 PMID:25236311.
- Hayward GN, Hay AD, Moore MV, et al. Effect of oral dexamethasone without immediate antibiotics vs placebo on acute sore throat in adults: a randomized clinical trial. *JAMA* 2017;317:1535-43. doi:10.1001/jama.2017.3417 PMID:28418482.
- Hayward G, Thompson M, Heneghan C, Perera R, Del Mar C, Glasziou P. Corticosteroids for pain relief in sore throat: systematic review and meta-analysis. *BMJ* 2009;339:b2976. doi:10.1136/bmj.b2976. PMID:19661138.
- Hayward G, Thompson MJ, Perera R, Glasziou PP, Del Mar CB, Heneghan CJ. Corticosteroids as standalone or add-on treatment for sore throat. *Cochrane Database Syst Rev* 2012;10:CD008268. doi:10.1002/14651858.CD008268.pub2. PMID:23076943.
- Korb K, Scherer M, Chenot JF. Steroids as adjuvant therapy for acute pharyngitis in ambulatory patients: a systematic review. *Ann Fam Med* 2010;8:58-63. doi:10.1370/afm.1038. PMID:20065280.
- Wing A, Villa-Roel C, Yeh B, Eskin B, Buckingham J, Rowe BH. Effectiveness of corticosteroid treatment in acute pharyngitis: a systematic review of the literature. *Acad Emerg Med* 2010;17:476-83. doi:10.1111/j.1553-2712.2010.00723.x PMID:20536799.

- 26 Siemieniuk RA, Agoritsas T, Macdonald H, Guyatt GH, Brandt L, Vandvik PO. Introduction to BMJ Rapid Recommendations. *BMJ* 2016;354:i5191. doi:10.1136/bmj.i5191. PMID:27680768.
- 27 Vandvik PO, Otto CM, Siemieniuk RA, et al. Transcatheter or surgical aortic valve replacement for patients with severe, symptomatic, aortic stenosis at low to intermediate surgical risk: a clinical practice guideline. *BMJ* 2016;354:i5085. doi:10.1136/bmj.i5085. PMID:27680583.
- 28 Guyatt GH, Oxman AD, Vist GE, et al. GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924-6. doi:10.1136/bmj.39489.470347.A. PMID:18436948.
- 29 Andrews JC, Schünemann HJ, Oxman AD, et al. GRADE guidelines: 15. Going from evidence to recommendation—determinants of a recommendation's direction and strength. *J Clin Epidemiol* 2013;66:726-35. doi:10.1016/j.jclinepi.2013.02.003. PMID:23570745.

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to <http://group.bmj.com/group/rights-licensing/permissions>

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

¹Academic Centre for General Practice, Department of Public Health and Primary Care, KU Leuven, Belgium

²CEBAM, Belgian Centre for Evidence-Based Medicine, Cochrane Belgium, Leuven, Belgium

³Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada L8S 4L8

⁴Division General Internal Medicine & Division of Clinical Epidemiology, University Hospitals of Geneva, CH-1211, Geneva, Switzerland

⁵Department of Medicine, University of Toronto, Toronto, Ontario, Canada

⁶Dutch College of General Practitioners, Utrecht, The Netherlands

⁷School CAPHRI, Department Family Medicine, Maastricht, The Netherlands

⁸Division of General Pediatrics, University Hospitals of Geneva & Faculty of Medicine, University of Geneva, Geneva, Switzerland

⁹Primary Care Clinical Unit, Faculty of Medicine, University of Queensland, Brisbane, Australia

¹⁰Pediatric Immunology, Department of microbiology and immunology, KU Leuven, Belgium

¹¹Pediatric allergy, Clinical division of pediatrics UZ Leuven, Leuven, Belgium

¹²Bali District Hospital, Bali and Centre for Development of Best practices in Health Yaounde, Cameroon

¹³Otorhinolaryngology—Head and Neck Surgery, Clinica del Country, Bogota, Colombia

¹⁴NIHR Oxford Diagnostic Evidence Cooperative, Oxford, UK

¹⁵Department of Primary Care Health Sciences, University of Oxford, Oxford, UK

¹⁶Oslo University Hospital, Forskningsveien 2b, Blindern 0317 Oslo, Norway

¹⁷The Society for Participatory Medicine, Newburyport, MA 01950-1183, USA

¹⁸Arizona Senior Academy, Tucson, AZ 85747, USA

¹⁹MOH Holdings, 1 Maritime Square, Singapore, Singapore 099253

²⁰Cochrane UK, London, UK

²¹Cochrane Consumers Group, Halifax, Canada

²²Washington DC, USA

²³HIV/STI Surveillance Research Center, and WHO Collaborating Center for HIV Surveillance, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

²⁴Department of Medicine, Innlandet Hospital Trust - division Gjøvik, Norway

²⁵Faculty of Dentistry, Universidad de Chile, Santiago, Chile