OPEN ACCESS

Check for updates

Genetic predisposition to increased serum calcium, bone mineral density, and fracture risk in individuals with normal calcium levels: mendelian randomisation study

Agustin Cerani,^{1,2} Sirui Zhou,^{1,2} Vincenzo Forgetta,¹ John A Morris,^{1,3} Katerina Trajanoska,^{4,5} Fernando Rivadeneira,^{4,5} Susanna C Larsson,⁶ Karl Michaëlsson,⁷ J Brent Richards^{1,2,3}

For numbered affiliations see end of the article.

Correspondence to: J B Richards
brent.richards@mcgill.ca

(ORCID 0000-0002-3746-9086)

Cite this as: *BMJ* 2019;366:l4410
<http://dx.doi.org/10.1136/bmj.l4410>

Accepted: 11 June 2019

ABSTRACT

OBJECTIVE

To determine if genetically increased serum calcium levels are associated with improved bone mineral density and a reduction in osteoporotic fractures.

DESIGN

Mendelian randomisation study.

SETTING

Cohorts used included: the UK Biobank cohort, providing genotypic and estimated bone mineral density data; 25 cohorts from UK, USA, Europe, and China, providing genotypic and fracture data; and 17 cohorts from Europe, providing genotypic and serum calcium data (summary level statistics).

PARTICIPANTS

A genome-wide association meta-analysis of serum calcium levels in up to 61 079 individuals was used to identify genetic determinants of serum calcium levels. The UK Biobank study was used to assess the association of genetic predisposition to increased serum calcium with estimated bone mineral density derived from heel ultrasound in 426 824 individuals who had, on average, calcium levels in the normal range. A fracture genome-wide association meta-analysis comprising 24 cohorts and the UK Biobank including a total of 76 549 cases and 470 164 controls, who, on average, also had calcium levels in the normal range was then performed.

RESULTS

A standard deviation increase in genetically derived serum calcium (0.13 mmol/L or 0.51 mg/dL) was not associated with increased estimated bone mineral density (0.003 g/cm², 95% confidence interval -0.059 to 0.066; P=0.92) or a reduced risk of fractures (odds ratio 1.01, 95% confidence interval

0.89 to 1.15; P=0.85) in inverse-variance weighted mendelian randomisation analyses. Sensitivity analyses did not provide evidence of pleiotropic effects.

CONCLUSIONS

Genetic predisposition to increased serum calcium levels in individuals with normal calcium levels is not associated with an increase in estimated bone mineral density and does not provide clinically relevant protection against fracture. Whether such predisposition mimics the effect of short term calcium supplementation is not known. Given that the same genetically derived increase in serum calcium is associated with an increased risk of coronary artery disease, widespread calcium supplementation in the general population could provide more risk than benefit.

Introduction

Fragility fractures are a large problem worldwide in both women and men, with an impact on quality of life and mortality.^{1 2} Calcium supplementation is promoted and emphasised by prevention and treatment guidelines to reduce the risk of osteoporosis and fractures,³⁻⁵ and is now common among the general adult population in high income countries.⁶⁻⁸ For example, in the NHANES study, 53% of Americans used dietary calcium supplements and 43% reported daily use.⁹

Evidence, however, from multiple studies indicates that increased serum calcium, a short term consequence of calcium supplementation, is associated with an increased risk of cardiovascular disease and mortality.^{10 11} A meta-analysis of serum calcium on incident risk of cardiovascular disease indicated that serum calcium was associated with an increased risk of cardiovascular disease.¹² Higher circulating calcium levels have also been found to be associated with an increased risk of stroke in observational studies.^{13 14}

However, observational associations of serum calcium with cardiovascular disease can be susceptible to confounding, even after controlling for known risk factors of the disease. Consequently, randomised controlled trials of calcium supplement use were undertaken because the process of randomisation breaks the association with confounding variables. Although these randomised controlled trials did not prespecify cardiovascular disease as a primary outcome, several meta-analyses have provided conflicting evidence, and most of these studies relied on short term calcium supplementation.^{10 15-19}

WHAT IS ALREADY KNOWN ON THIS TOPIC

Calcium supplementation in the general population is common and often intended to reduce the risk of fracture

Calcium supplementation has been associated with an increased risk of coronary artery disease and its protective effects on bone health remain unclear

WHAT THIS STUDY ADDS

Genetic predisposition to increased serum calcium levels in individuals with normal calcium levels is not associated with an increase in estimated bone mineral density and does not provide clinically relevant protection against fracture

Given that the same genetically derived increase in serum calcium is associated with an increased risk of coronary artery disease, widespread calcium supplementation in the general population might provide more risk than benefit

Another way to overcome confounding is by mendelian randomisation. Mendelian randomisation is an established genetic epidemiology method that uses natural genetic variation to strengthen causal inference by mimicking a lifelong randomised controlled trial.²⁰ Specifically, genetic variants are identified that are reproducibly associated with the risk factor and are then tested for their combined effect on the disease outcome. Since genetic variants are randomly assigned at conception, this method greatly decreases confounding. Further, since conception always precedes disease onset, such studies are not prone to reverse causation. Mendelian randomisation studies are less prone to regression dilution bias than observational studies because genotypes are measured with a high degree of precision. Lastly, mendelian randomisation provides an estimate of lifelong exposure. Nonetheless, mendelian randomisation studies are limited by potential bias owing to horizontal pleiotropy, where the genetic variant influences the outcome, independently of the exposure, among other limitations.²¹

Our recent mendelian randomisation study found that lifelong genetically predicted increased serum calcium levels were associated with a higher risk of coronary artery disease and myocardial infarction,²² such that a one standard deviation increase in serum calcium (0.13 mmol/L or 0.51 mg/dL) was associated with an increased odds of coronary artery disease (odds ratio 1.25, 95% confidence interval 1.08 to 1.45, $P=0.003$), comparable with previous randomised controlled trial meta-analysis estimations.¹² Thus, given the risks of increased serum calcium and the high prevalence of calcium supplementation, it is important to understand the potential beneficial effects of calcium on skeletal health and fracture so that patients and their doctors can balance potential risks against potential benefits.

Clearly calcium is required for normal skeletal development and maintenance since net calcium excretion must be replaced.⁵ Indeed, severe hypocalcemia due to deficient calcium or vitamin D intake, or both, leads to diminished bone density and increased risk of fracture, which is improved with increased calcium intake.²³⁻³² Yet, what remains unclear is whether additional calcium supplementation to an ordinary diet can lead to clinically relevant improvements in heel ultrasound bone mineral density and prevent fractures in the general adult population, who in general, have normal serum calcium and parathyroid hormone levels. Randomised controlled trial data supporting calcium supplementation to prevent fractures is inconsistent and even large trials, such as the Women's Health Initiative, have not shown any reduction in the risk of fracture with calcium plus vitamin D in community dwelling older women and men.³³⁻³⁸ Calcium supplementation alone might even increase the risk of hip fracture, the most devastating type of fragility fracture.³⁸⁻³⁹ Further, a recent randomised controlled trial using bisphosphonates to prevent bone fractures found profound beneficial

effects without calcium supplementation,⁴⁰ a result also supported by another randomised controlled trial with bone mineral density as an outcome.⁴¹

Given the potential risks of calcium supplements and their widespread use, it is important to understand if increasing calcium results in a reduced risk of osteoporosis and fracture. We therefore assessed whether genetically predicted lifelong higher serum calcium levels were associated with bone mineral density and the risk of fracture by using mendelian randomisation. To do so, we identified the genetic determinants of serum calcium levels in 61079 individuals and tested their effect on estimated bone mineral density ($n=426\,824$) and the risk of fracture (76 549 cases and 470 164 controls).

Methods

Study design and data sources

Selection of instrumental variables

Figure 1 shows that the causal interpretation of mendelian randomisation estimates relies on three assumptions. Firstly, the genetic variants, termed single nucleotide polymorphisms, must be associated with the risk factor of interest. Secondly, the genetic variant must not be associated with confounders (common causes of the risk factor and outcome association), which are not in the causal pathway between the risk factor and the outcome. Thirdly, the genetic variant is independent of the outcome conditional on the risk factor and confounders (that is, absence of horizontal pleiotropy). This means that the genetic variant's effect on the outcome should only be mediated by the risk factor and, thus, not have a direct effect on the outcome independent of the risk factor. We undertook a two sample mendelian randomisation approach to test the effect of increased serum calcium on bone mineral density and fracture.⁴² Two sample mendelian randomisation identifies genetic variants to be associated with the exposure in one dataset and then tests the association of these variants with the outcome in a separate dataset. The advantage of this approach is that it allows for larger sample sizes, providing more precise estimates of effect of the exposure on the outcome. For an overview of the concepts and methods deployed in mendelian randomisation studies, we refer interested readers to a recent review by Holmes and colleagues.⁴³

Associations between single nucleotide polymorphisms and serum calcium concentration

We obtained seven single nucleotide polymorphisms associated with total serum calcium concentrations at a genome-wide significant level of $P<5\times 10^{-8}$ from the largest serum calcium genome-wide association study meta-analysis to date.⁴⁴ Genome-wide significant associations between the single nucleotide polymorphism and serum calcium comply with the first assumption of mendelian randomisation (that is, association between instrument and exposure). The study consisted of a discovery cohort of 39 400 individuals of European descent from 17 population

Fig 1 | Mendelian randomisation assumptions

based studies. Serum calcium concentrations were quantified by colorimetric assay in most cohorts.

Discovery analysis identified 14 genome-wide significant single nucleotide polymorphisms associated to serum calcium concentration, of which seven were replicated in a subsequent replication cohort of 21 676 individuals of European descent. The replicated single nucleotide polymorphisms were rs1801725 in *CASR*, rs1550532 in *DGKD*, rs780094 in *GCKR*, rs7336933 near *VWA8* and *DGKH*, rs10491003 near *GATA3*, rs7481584 in *CARS*, and rs1570669 near *CYP24A1* (table 1). In addition to the seven replicated single nucleotide polymorphisms, we included a genetic variant, rs17711722: *VKORC1L1* locus, which failed replication as defined by O'Seaghdha and colleagues ($P > 0.05$ for replication),⁴⁴ although the single nucleotide polymorphism remained genome-wide significant after meta-analysis of their discovery and replication results. In addition, rs17711722: *VKORC1L1* locus displayed functional plausibility given that it encodes an enzyme involved in vitamin K physiology, which is associated with calcium homeostasis.^{45 46}

Bone mineral density and fracture genome-wide association study

Bone mineral density is a clinically relevant measure used to diagnose osteoporosis and to risk-stratify for fracture. Estimated bone mineral density is derived from two heel ultrasound measures, velocity

of sound, and broadband ultrasound attenuation, which is highly heritable (50% to 80%),⁴⁷ and is a strong predictor of risk of fracture.^{48 49} To estimate the effect of serum calcium level on estimated bone mineral density and fracture we obtained summary statistics for the associations between the eight calcium modifying single nucleotide polymorphisms and estimated bone mineral density and fractures risk from our recent genome-wide association study on estimated bone mineral density consisting of 426 824 white British individuals from the UK Biobank.⁵⁰ In this well powered genome-wide association study, 518 genome-wide significant loci accounted for 20% of the total variance in estimated bone mineral density, using 13.7 million single nucleotide polymorphisms imputed to the Haplotype Reference Consortium panel.⁵¹

We performed an updated fracture fixed effect meta-analysis comprising a total of 24 cohorts from two recently published fracture genome-wide association studies, which included 23 cohorts from Genetic Factors for Osteoporosis consortium (GEFOS), EPIC-Norfolk study, and UK Biobank's full release.^{50 52} The meta-analysis involved a total of 76 549 cases and 470 164 controls (table 2). Fracture cases in GEFOS and EPIC-Norfolk cohorts included adults who had any fractures confirmed by medical, radiological, or questionnaires with the exclusion of skull, fingers, and toes as well as high trauma fractures, when available. The details of case ascertainment of each cohort were described previously.⁵² For UK Biobank, we included fracture cases reported from either hospital based fracture diagnosis according to ICD-10 (international classification of diseases, 10th revision) codes, or questionnaire based self reported fracture. Fractures located at skull, face, hands and feet, pathological, and atypical fractures; periprosthetic fractures were excluded. Detailed description of fracture ascertainment in UK Biobank was described previously.⁵⁰ Controls from all cohorts were defined as patients without a history of fracture. Approximately 69.5% of all fracture cases and 78.1% of all samples in the 24 cohorts were from UK Biobank, whereas GEFOS

Table 1 | Summary statistics for calcium estimated bone mineral density (eBMD) and fracture for single nucleotide polymorphisms (SNPs) influencing serum calcium

Nearby gene	Chr	Associated SNP	Calcium increasing allele	Calcium serum GWAS			eBMD GWAS *			Fracture GWAS		
				Allele freq	Effect (mmol/L)	P value	Allele freq	Effect (g/cm ²)	P value	Allele freq	Odds ratio (95% CI)	P value
<i>DGKD</i>	2	rs1550532	C	0.32	0.0045	8×10^{-11}	0.32	0.0022	0.50	0.32	1.004 (0.992 to 1.016)	0.51
<i>CASR</i>	3	rs1801725	T	0.15	0.0178	9×10^{-86}	0.13	0.0046	0.077	0.13	0.993 (0.976 to 1.011)	0.45
<i>GATA3</i>	10	rs10491003	T	0.09	0.0068	5×10^{-9}	0.09	-0.0002	0.77	0.09	1.013 (0.993 to 1.034)	0.21
<i>CARS</i>	11	rs7481584	G	0.72	0.0045	1×10^{-10}	0.72	-0.0067	0.001	0.71	1.003 (0.990 to 1.016)	0.62
<i>DGKH, VWA8</i>	13	rs7336933	G	0.85	0.0055	9×10^{-10}	0.85	0.00098	0.83	0.85	1.018 (1.000 to 1.033)	0.05
<i>CYP24A1</i>	20	rs1570669	G	0.34	0.0045	9×10^{-12}	0.34	-0.0045	0.016	0.34	0.993 (0.981 to 1.005)	0.25
<i>VKORC1L1</i>	7	rs17711722	T	0.47	0.00375	8×10^{-9}	0.44	0.0043	0.0180	0.45	0.998 (0.986 to 1.010)	0.68

*Mean (SD) eBMD is 0.54 g/cm² (0.12 g/cm²)

GWAS=genome-wide association study

Table 2 | Cohort descriptions

Source	Study design	Country of origin	Ethnicity	Mean (SD) age	Study type	Assessment method	Fracture	Non-fracture	Total
Estimated bone mineral density cohort									
UK Biobank	Cohort	UK	Mixed (white British subset used for analysis)	56.6 (8.1)	Cohort	Heel quantitative ultrasound (heel BMD)	NA	NA	426 824
Fracture cohort									
Age, Gene/Environment Susceptibility Reykjavik Study	Cohort	Iceland	Northern European	76.4 (5.5)	Population based	Medical and radiographic records	1458	1727	3185
Anglo-Australasian Osteoporosis Genetics Consortium	Population cohort, and case control for fracture cases	Australia	North western European	69.6 (8.6)	Population based, clinical based	Questionnaire, radiography	685	1113	1798
B-vitamins for the PRevention Of Osteoporotic Fractures	Intervention study	Netherlands	North western European	74.3 (6.5)	General population	History of fractures before baseline: questionnaire. Incident fractures: self report, validated at GP or hospital	715	1483	2198
Cardiovascular Health Study	Cohort	US	European American	73.2 (5.9)	Population based	Self report of incident fracture of the hip, leg, arm, or vertebra	519	2742	3261
DeCODE Genetics Study	Cross sectional	Iceland	North western European	60.7 (13.9)	Population based, clinical based	Medical records, radiographic documentation, questionnaire	1836	14 560	16 396
Estonian Genome Center University of Tartu-I	Cohort	Estonia	Northern European	55.4 (20.2)	Population based	Medical records, questionnaire	217	4296	4513
Estonian Genome Center University of Tartu-II	Cohort	Estonia	Northern European	40.3 (16.1)	Population based	Medical records, questionnaire	71	1717	1788
Erasmus Rucphen Family	Cohort	Netherlands	North western European	48.8 (14.6)	Family based isolate	Interview	260	1342	1602
Framingham Heart Study	Cohort	US	European American	64.7 (11.2)	Population and family based	Medical records, questionnaire	1520	2782	4302
Göteborg Osteoporosis and Obesity Determinants Study	Cohort	Sweden	Northern European	18.9 (0.6)	Population based	Radiographic document	273	597	870
Health Aging and Body Composition	Cohort	US	European American	73.8 (2.9)	Population based	Radiographic	308	1353	1661
Hong Kong Southern Chinese	Case control	China	Southern Chinese	48.9 (15.6)	Population based, clinic based	Medical records, radiographic, and questionnaire	79	627	706
MrOS	Cohort	US	European American	73.9 (5.9)	Clinic based	Questionnaire, radiographic documentation	918	3555	4473
The PROspective Study of Pravastatin in the Elderly at Risk	Cohort, randomised controlled trial	Netherlands, UK, Ireland	North western European	75.4 (3.4)	Clinic based	Medical records	426	4816	5242
Rotterdam Study I	Cohort	Netherlands	North western European	69.4 (9.0)	Population based	Medical records, questionnaire	2163	3574	5737
Rotterdam Study II	Cohort	Netherlands	North western European	63.8 (7.1)	Population based	Medical records, questionnaire	932	1220	2152
Rotterdam Study III	Cohort	Netherlands	North western European	56.1 (5.4)	Population based	Medical records, questionnaire	505	2421	2926
Study of Osteoporotic Fractures	Cohort	US	European American	71.5 (5.2)	Clinic based	Questionnaire, radiographic documentation	1611	1698	3309
TwinsUK	Cohort	UK	North western European	49.9 (13.6)	Population based, family based	Medical records, radiographic, and questionnaire	839	4111	4950
Women's Genome Health Study	Cohort	US	European American	54.1 (7.1)	Population based	Questionnaire	1832	20 498	22 330
Women's Health Initiative Clinical Trial	Quasi case/control	US	European American	69.0 (6.4)	Population based	Medical records	1058	647	1705
Women's Health Initiative Observational Study	Quasi case/control	US	European American	69.0 (6.5)	Population based	Medical records	1603	989	2592
CV risk in Young Finns Study	Cohort	Finland	Northern European	38.0 (5.0)	Population based	Medical records	611	975	1586
European Prospective Investigation into Cancer, Norfolk study	Cohort	UK	North western European	59.1 (9.3)	Population based	Medical records	2926	17 710	20 636
UK Biobank	Cohort	UK	Mixed (white British subset used for analysis)	56.8 (8.0)	Cohort	Questionnaire, based on answering yes to the question "Have you fractured/broken any bones in the last 5 years?" at either baseline or first follow up	53 184	373 611	426 795

cohorts represented 26.7% of cases and 18.2% of all samples and EPIC-Norfolk provided 2.8% of all cases and 3.8% of all samples. Table 2 shows the mean age of the GEFOS and EPIC-Norfolk cohorts.

Linear regression involving UK Biobank's fracture cohort was performed using BOLT-LMM,⁵³ an algorithm that allows for large scale mixed model association testing. Our genome-wide association study analysis included adjustment for sex, age, assessment centre, genotyping array, and first 20 principle components. Before meta-analysis with GEFOS and EPIC-Norfolk, and owing to the use of BOLT-LMM,^{53 54} the effect size estimate and standard error of each genetic variant (single nucleotide polymorphism) from UK Biobank's genome-wide association study analysis was transformed to odds ratios by using the following approximation: $\log \text{odds ratio} = \beta / (\mu * (1 - \mu))$, where μ is the case fraction in UK Biobank's fracture cohort.⁵²

The UK Biobank full release fracture genome-wide association study was performed on its white British subpopulation, which included 53 184 fracture cases and 373 611 controls. The mean age of the UK Biobank fracture cohort is 56.8 ± 8.0 (table 2). The study cohorts involved in the serum calcium genome-wide association study were mostly of European descent.⁴⁴ The estimated bone mineral density genome-wide association study cohort consisted of all white British patients from the UK.⁵⁵ Fracture cohorts were predominantly of European descent: Europe (91.6%), North America (8.0%), Australia (0.3%), and East Asia (0.1%).⁵⁰

We used METAL to perform fixed effects meta-analysis of these results. Individual genome-wide association studies were corrected by genomic control,⁵⁶ and a total of 8 818 767 autosomal single nucleotide polymorphisms were included in the meta-analysis. The average genomic inflation λ of 24 cohorts was 1.025 and was adjusted accordingly when performing the meta-analysis. Finally, summary statistics from the eight calcium associated single nucleotide polymorphisms (table 1) were extracted from the fracture meta-analysis described above.

Single nucleotide polymorphism validation and pleiotropy assessment

We next undertook sensitivity analysis to understand if any of the single nucleotide polymorphisms might violate assumptions of mendelian randomisation.

Linkage disequilibrium

Single nucleotide polymorphisms used for mendelian randomization analysis are assumed to be independent of each other.²¹ Given that the eight identified calcium associated single nucleotide polymorphisms are located on different chromosomes, they would segregate independently of each other and, hence, are not in linkage disequilibrium.

Pleiotropy

To avoid a biased estimation of the effect of serum calcium (risk factor) on either the risk of estimated bone mineral

density or fracture (outcomes), the genetic variants (instruments) used in the mendelian randomisation analysis should only affect the outcome only through serum calcium. Thus, we evaluated potential associations of our selected calcium associated single nucleotide polymorphisms with known determinants of bone mineral density and fracture by searching the selected single nucleotide polymorphisms in Phenoscanner, a database of genome-wide association study results.⁵⁷ We further assessed whether each of the eight single nucleotide polymorphisms were expression quantitative trait loci for genes that could be associated with known determinants of bone mineral density and fracture by using the GTEX database.^{50 58} Although direct pleiotropic effects that influence the outcome independently of the risk factor violate mendelian randomisation assumptions (red arrow in fig 1), vertical pleiotropy does not. Vertical pleiotropy is defined as the association of a single nucleotide polymorphism with more than one phenotype in the same biological pathway.⁴³ For example, genetically lowered calcium at CASR could lead to increased parathyroid hormone,⁵⁹ which itself might influence bone mineral density and fracture. But since calcium directly influences parathyroid hormone, which influences the outcome, this is not a violation of mendelian randomisation assumptions.

Comparison of genetically derived effects with pharmacological effects of calcium supplementation

We modelled the effect of a one standard deviation genetically derived increase in serum calcium on estimated bone mineral density and fracture. We compared this one standard deviation increase with the magnitude of increase in serum calcium levels after calcium supplementation. To do so, we compared data from a recent randomized crossover trial of calcium supplementation.⁶⁰

Mendelian randomisation

Individual mendelian randomisation estimates from the seven independent serum calcium associated single nucleotide polymorphisms were calculated by using the Wald method.⁶¹ We meta-analysed individual mendelian randomisation estimates by using both inverse-variance weighted and a random effects models using R and the package MendelianRandomization and RStudio.⁶²⁻⁶⁴ The estimated associations of genetically predicted serum calcium with estimated bone mineral density and odds of fracture were expressed with respect to one standard deviation increase in serum calcium levels, which is equivalent to 0.51 mg/dL or 0.13 mmol/L. This standard deviation equivalence was derived from serum calcium's pooled variance calculation involving the 30 cohorts reported in O'Seaghdha and colleagues and included in the serum calcium genome-wide association study.⁴⁴

Sensitivity analyses

To explore potential pleiotropic effects, we carried out three sensitivity meta-analyses: simple and

weighted median and mendelian-randomisation-Egger regression methods using the R package MendelianRandomization.⁶³ Simple and weighted median meta-analyses provide estimations that are robust to the inclusion of up to 50% invalid instruments in a mendelian randomisation analysis.⁶⁵ The intercept estimate from mendelian-randomisation-Egger regression analysis provides a useful estimation of directional horizontal pleiotropy, that is, the magnitude and direction of the effect of the single nucleotide polymorphisms on the outcome not mediated through the exposure.⁶⁶

In addition to the primary analysis, we performed two additional sensitivity analyses. Firstly, we repeated our analysis including rs17711722 near *VKORC1L1*. rs17711722 is associated with serum calcium levels at a genome-wide significant level, but it was not included in the main findings because it did not meet replication criteria.⁴⁴ Lastly, we performed an additional analysis by excluding rs1801725 (*CASR*), which contributed the most weight in the inverse-variance weighted primary meta-analysis (*CASR* single nucleotide polymorphism explained 0.49% of the serum calcium variance). This single nucleotide polymorphism is in LD with single nucleotide polymorphism rs73186030, which has been associated with parathyroid hormone levels.⁵⁹ The rationale was to test whether, in its absence, the estimated calcium on bone mineral density and on fracture effects were similar to those from primary analyses or if they were mostly driven by *CASR*.

In addition, to assess whether Asian ancestry influenced our fracture mendelian randomisation results, we performed the fracture genome-wide association study meta-analysis removing the HKOS (Southern Chinese) cohort and repeated the fracture mendelian randomisation.

Patient and public involvement

No patients or member of the general public were directly involved in the design, recruitment, or conduct of the study. After publication, dissemination of the results will be sought across different countries involving respective patient organisations, the general public, and other stakeholders; typically, across social media, scientific meetings, and media interviews.

Results

Single nucleotide polymorphism selection

Table 1 shows that the single nucleotide polymorphisms in eight loci previously identified to be associated with serum calcium were rs1801725 in *CASR* ($P=9\times 10^{-86}$), rs1550532 in *DGKD* ($P=8\times 10^{-11}$), rs780094 in *GCKR* ($P=1\times 10^{-10}$), rs7336933 near *VWA8* and *DGKH* ($P=9\times 10^{-10}$), rs10491003 nearby *GATA3* ($P=5\times 10^{-9}$), rs7481584 in *CARS* ($P=1\times 10^{-10}$), rs1570669 near *CYP24A1* ($P=9\times 10^{-12}$), and rs17711722 near *VKORC1L1* ($P=8\times 10^{-9}$). The selected single nucleotide polymorphisms collectively explained 0.77% of the variance in total serum calcium levels, which is sufficient to influence the risk of coronary artery disease.²²

Known biology at associated loci

Among these single nucleotide polymorphisms, rs1801725 (*CASR*) and rs1570669 (*CYP24A1*) are located nearby genes whose functions are involved in calcium homeostasis.²² The most strongly associated calcium locus includes *CASR*, a calcium sensing receptor. *CASR* encodes a protein whose main function is to capture small changes in circulating calcium concentrations and consequently modify parathyroid hormone secretion and renal cation handling.⁶⁷ *CYP24A1* encodes an enzyme that plays a role in calcium homeostasis and the metabolism of the active form of vitamin D.⁶⁸ The diacylglycerol kinase genes *DGKD* and *DGKH* genes have recently been implicated in calcium signaling.⁶⁹ *GATA3* and the *CARS* locus are reportedly associated with hypocalcemia in the hypoparathyroidism, sensorineural deafness, and renal dysplasia (hypoparathyroidism, deafness, and renal dysplasia syndrome) and Beckwith-Wiedemann syndromes, respectively.⁷⁰ The remaining single nucleotide polymorphism, rs17711722, used for sensitivity analysis is located in *VKORC1L1*, is also associated with calcium homeostasis.⁴⁵ Thus, all loci associated with calcium levels contained genes with plausible biological effects on calcium levels.

Pleiotropy evaluation

Single nucleotide polymorphisms rs1801725 in *CASR*, rs7336933 near *VWA8* and *DGKH*, rs10491003 nearby *GATA3*, rs7481584 in *CARS*, and rs1570669 near *CYP24A1* were not associated with any phenotypes other than serum calcium in the Phenoscanner and MRBase databases. Besides its association with calcium levels, rs1550532 in *DGKD* showed evidence of association with bilirubin levels.⁷¹⁻⁷³ Bilirubin, however, is not associated with bone mineral density or with fracture.⁷⁴ Single nucleotide polymorphism rs17711722 (*VKORC1L1*) showed genome-wide level associations with corneal structure and central corneal thickness,⁷⁵ yet these phenotypes are not known to be related to a calcium-independent effect on estimated bone mineral density and fracture. Single nucleotide polymorphism rs780094 in *GCKR* had genome-wide level associations with triglycerides levels, cholesterol, waist circumference, and several other lipid-related phenotypes.^{73 76-78} Waist circumference is highly associated with weight and BMI, which are known determinants of lower extremity bone density and potentially heel bone mineral density.⁷⁹ Thus, the highly pleiotropic nature of rs780094 in *GCKR* represents a calcium-independent effect on our outcomes of interest and was removed from all subsequent analyses.

Association of calcium levels modifying single nucleotide polymorphisms with estimated risk of bone mineral density and fracture

Table 1 shows that the summary statistics for the association between seven calcium increasing single nucleotide polymorphisms and estimated bone mineral density and fracture odds were directly

Table 3 | Mendelian randomisation results for effect of serum calcium on estimated bone mineral density (eBMD) and fracture risk

Method	Main analysis				Sensitivity Analyses							
	6 serum calcium associated SNPs				Including rs17711722 (VKORC1L1)				Excluding rs1801725 (CASR)			
	Mean eBMD (95% CI)	P value	Fracture odds ratio (95% CI)	P value	Mean eBMD (95% CI)	P value	Fracture odds ratio (95% CI)	P value	Mean eBMD	P value	Fracture odds ratio (95% CI)	P value
IVW	0.003 (−0.059 to 0.066)	0.92	1.01 (0.89 to 1.15)	0.85	0.011 (−0.05 to 0.073)	0.72	1.01 (0.90 to 1.13)	0.91	−0.049 (−0.144 to 0.047)	0.32	1.12 (0.92 to 1.36)	0.25
Simple median	0.009 (−0.052 to 0.070)	−0.76	1.11 (0.93 to 1.33)	0.24	0.023 (−0.035 to 0.081)	0.44	1.10 (0.91 to 1.32)	0.32	−0.004 (−0.093 to 0.085)	0.93	1.13 (0.89 to 1.42)	0.32
Weighted median	0.030 (−0.006 to 0.067)	−0.10	0.99 (0.89 to 1.11)	0.91	0.031 (−0.005 to 0.068)	0.09	0.98 (0.88 to 1.10)	0.76	−0.009 (−0.093 to 0.076)	0.84	1.12 (0.90 to 1.41)	0.31
MR-Egger intercept	−0.003 (−0.006 to 0.001)	−0.11	1.00 (1.00 to 1.01)	0.39	−0.001 (−0.005 to 0.002)	0.49	1.00 (1.00 to 1.01)	0.52	−0.005 (−0.019 to 0.008)	0.45	0.98 (0.96 to 1.01)	0.14

Results are expressed per one standard deviation (0.51 mg/dL or 0.13 mmol/L) increase in serum calcium concentration. Estimated bone mineral density (eBMD) is expressed in g/cm².

IVW=inverse-variance weighted meta-analysis

obtained without the use of proxy single nucleotide polymorphisms from their respective studies.⁵⁵ None of the seven calcium single nucleotide polymorphisms had genome-wide significant associations with either estimated bone mineral density or odds of fracture (all $P>0.08$).

Comparison of genetically derived effects with pharmacological effects of calcium supplementation

A previous crossover randomised controlled trial showed that 500 mg of calcium citrate in a fasting state lead to a maximal increase in total serum calcium levels by approximately 0.07 mmol/L, four hours after administration.⁶⁰ The mendelian randomisation analyses here represent a change in total serum calcium of one standard deviation, which is 0.13 mmol/L. Therefore, the effects of total serum calcium presented include the anticipated effects of calcium supplementation.

Mendelian randomisation analysis: serum calcium on estimated bone mineral density

Table 3 and figure 2 show that when performing mendelian randomisation analyses, a one standard deviation (that is, 0.51 mg/dL or 0.13 mmol/L) increase in serum calcium concentration was not associated with a clinically relevant change in estimated bone mineral density (change per standard deviation increase in serum calcium 0.003 g/cm², 95% confidence interval −0.059 to 0.066; $P=0.92$). The mean and standard deviation of estimated bone mineral density are 0.54 g/cm² and 0.12 g/cm², respectively.

Figure 2 lists the individual level randomisation estimates of the single nucleotide polymorphisms used in the inverse variance weighted analysis. Mendelian randomisation estimates as determined by rs7481584 (CASR −0.19 g/cm², 95% confidence interval −0.30 to −0.08; $P=0.001$) and rs1570669 (CYP24A1 −0.13 g/cm², −0.24 to −0.02; $P=0.02$) showed a statistically significant decrease in estimated bone mineral density per standard deviation increase in serum calcium. However, only the former remained statistically significant after Bonferroni correction for multiple hypothesis testing involving six tests, that is, $0.05/6=8.3 \times 10^{-3}$. Mendelian randomisation estimates as determined by the remaining single nucleotide

polymorphisms showed a lack of association between a standard deviation increase in serum calcium and estimated bone mineral density.

Table 3 shows that the sensitivity meta-analyses with six single nucleotide polymorphisms involving simple median (0.009 g/cm², 95% confidence interval −0.052 to 0.070; $P=0.76$) and weighted median estimation (0.030 g/cm², −0.006 to 0.067; $P=0.10$) supported the inverse-variance weighted primary analysis. The mendelian-randomisation-Egger regression intercept, which provides an approximate estimation of directional pleiotropic effects on estimated bone mineral density through pathways independent of serum calcium, showed no significant evidence for such effects (−0.003 g/cm², −0.006 to 0.001; $P=0.11$).

Table 3 shows that the inclusion of an additional serum calcium increasing single nucleotide polymorphism (rs17711722, VKORC1L1) to the primary analysis also indicated that a one standard deviation increase in serum calcium was not associated with clinically relevant change in estimated bone mineral density of 0.011 g/cm² (95% confidence interval −0.050 to 0.073; $P=0.72$).

To assess the degree to which our primary mendelian randomisation inverse-variance weighted estimate would change by removing the single nucleotide polymorphism that provided most weight to the inverse-variance weighted analysis, we ran an additional sensitivity analysis excluding rs1801725 (CASR). Results were, as expected, less precise but did not differ materially from the results of the primary analyses (eg, inverse-variance weighted estimate −0.049 g/cm², 95% confidence interval −0.144 to 0.047; $P=0.32$; table 3).

Mendelian randomisation analysis: serum calcium association with fracture

We estimated the effect of a genetically average increased serum calcium on odds of fracture by implementing a random effects model and inverse-variance weighted method, which included six calcium-increasing alleles described in table 1. Figure 3 and table 3 show that a one standard deviation increase in serum calcium concentration was not associated with odds of fracture (odds ratio 1.01, 95% confidence interval 0.89 to 1.15; $P=0.85$).

Fig 2 | Serum calcium effects on estimated bone mineral density (eBMD). Two sample mendelian randomisation: individual and inverse-variance weighted (IVW) results

Figure 3 shows the estimates from individual level single nucleotide polymorphism mendelian randomisation analysis. Mendelian randomisation estimates after Bonferroni correction for multiple hypothesis testing involving six tests did not show a change in odds of fracture per one standard deviation increase in serum calcium.

Table 3 shows that sensitivity meta-analyses with six single nucleotide polymorphisms involving simple median (odds ratio 1.11, 95% confidence interval 0.93 to 1.33; $P=0.24$) and weighted median estimation (0.99, 0.89 to 1.11; $P=0.91$) supported inverse-variance weighted primary analysis results. Mendelian-randomisation-Egger intercept regression results (1.00, 1.00 to 1.01; $P=0.39$) provided no evidence of directional pleiotropic effects on fracture odds through pathways independent of serum calcium.

The inclusion of an additional serum calcium increasing single nucleotide polymorphism (rs1-7711722, *VKORC1L1*) to our primary analysis with one standard deviation increase in serum calcium (inverse-variance weighted odds ratio 1.01, 95% confidence interval 0.90 to 1.13; $P=0.91$).

We also assessed the degree to which our primary mendelian randomisation inverse-variance weighted estimate would change by removing the single nucleotide polymorphism that provided highest

weight to the inverse-variance weighted analysis, that is, rs1801725 (*CASR*). Again, the results were less precise, but were not materially different from the primary inverse-variance weighted estimate (inverse-variance weighted odds ratio 1.12, 95% confidence interval 0.92 to 1.36; $P=0.25$).

Finally, to assess whether the presence of a cohort of Asian ancestry (HKOS) in our fracture genome-wide association study could affect our fracture mendelian randomisation results, we performed the fracture genome-wide association study meta-analysis after removing the HKOS (Southern Chinese) cohort and repeated the fracture mendelian randomisation analysis. As observed in table 4 and figure 4, our instrumental variables' summary statistics and mendelian randomisation results were virtually identical to those obtained in our primary analysis. The genomic inflation factor (λ) without and with the inclusion of HKOS cohort remained unchanged at 1.025. Therefore, inclusion of the HKOS cohort of Southern Chinese ancestry did not affect our results.

Discussion

This mendelian randomisation study showed that a standard deviation increase in lifelong serum calcium levels was not associated with increased estimated bone mineral density or reduced risk of fracture in individuals with normal calcium levels. The magnitude of a one standard deviation increase in genetically predicted serum calcium includes the increase in serum calcium that would be anticipated after calcium supplementation.^{60 80} Assuming a linear effect between calcium levels and the studied outcomes, this suggests that widespread efforts to use calcium supplements in the general population for long periods of time are unlikely to have any substantial effect on bone health outcomes. Further, we have recently shown that genetically determined increase in serum calcium derived from the same instruments (that is, single nucleotide polymorphisms) is associated with a clinically relevant increase in the risk of coronary artery disease.²² Thus, the cardiovascular risks of

Table 4 | Summary statistics for fracture single nucleotide polymorphisms (SNPs) influencing serum calcium including and excluding HKOS cohort of Asian descent

Nearby gene	Chr	Associated SNP	Calcium increasing allele	Allele freq	Fracture GWAS		Allele freq	Fracture GWAS excluding HKOS cohort	
					Odds ratio (95% CI)	P value		Odds ratio (95% CI)	P value
DGKD	2	rs1550532	C	0.32	1.004 (0.992 to 1.016)	0.51	0.32	1.004 (0.992 to 1.017)	0.53
CASR	3	rs1801725	T	0.13	0.993 (0.976 to 1.011)	0.45	0.13	0.993 (0.976 to 1.011)	0.44
GATA3	10	rs10491003	T	0.09	1.013 (0.993 to 1.034)	0.21	0.09	1.013 (0.993 to 1.034)	0.21
CARS	11	rs7481584	G	0.71	1.003 (0.990 to 1.016)	0.62	0.71	1.004 (0.991 to 1.017)	0.59
DGKH, VWA8	13	rs7336933	G	0.85	1.018 (1.000 to 1.033)	0.05	0.85	1.017 (1.000 to 1.033)	0.05
CYP24A1	20	rs1570669	G	0.34	0.993 (0.981 to 1.005)	0.25	0.34	0.993 (0.980 to 1.005)	0.24
VKORC1L1	7	rs17711722	T	0.45	0.998 (0.986 to 1.010)	0.68	0.45	0.998 (0.986 to 1.010)	0.71

GWAS=genome-wide association study

Fig 3 | Serum calcium effects on odds of fracture. Two sample mendelian randomisation: individual and inverse variance weighted (IVW) results

increasing serum calcium in the general population are unlikely to be offset by beneficial effects on bone density and fracture.

Calcium is vital to many biological processes, and its serum concentration is tightly regulated. Net calcium excretion must be replaced, but the amount of calcium needed is debated. What is not well understood is whether increases in serum calcium amongst individuals who have a normal varied diet and normal calcium levels lead to a decrease in the risk of fracture. As outlined above, there is conflicting observational epidemiological evidence that calcium supplementation does not reduce the risk of fracture, yet such studies could be prone to bias since the individuals most likely to use calcium supplements are those more likely to be at a higher risk of fracture.⁸¹ The mendelian randomisation approach employed here overcomes this potential confounding by relying on the random assignment of alleles at conception, thereby preventing associations with such potentially confounding factors.

One way to improve the quality of evidence in medical research is to employ the principles of triangulation of different sources of evidence. If results are consistent across different types of study designs, and these different types of designs have different sources of potential bias, then the results can be combined in the framework of triangulation to provide a higher standard of evidence.^{82 83} Of importance, our recent mendelian randomisation analysis that

instrumented lactase persistence in adults as a surrogate for consumption of calcium-bearing dairy products found evidence of no protective effect of sustained dairy intake on the risk of fracture,^{52 84} which is supported by our present results. Further, our mendelian randomisation findings are consistent with those of the aforementioned observational studies, numerous randomised controlled trials, and randomised controlled trial meta-analysis.³³⁻³⁸

Strengths of this study include the large sample size for estimated bone mineral density and fracture.^{50 52} Further, multiple sensitivity analyses, including removing the single nucleotide polymorphisms most strongly associated with calcium levels, led to similar findings. Despite multiple sensitivity analyses, we did not identify pleiotropic effects of the single nucleotide polymorphisms on the skeletal outcomes, independent of serum calcium, as a potential source of bias. Of relevance, the serum calcium genome-wide association study from where we obtained serum calcium instruments was composed of population based cohorts that, on average, are normocalcemic, thus our reference population is normocalcemic adults. Hence, our study conclusions are applicable to normocalcemic adult populations similar to those in the general population, not to subpopulations, for example, with extremely low calcium level, which could benefit from calcium supplementation.

Limitations

These findings cannot provide insight into the effects of hypocalcemia and its correction on estimated bone mineral density and the risk of fractures. We have assumed a linear effect between calcium levels and the studied outcomes and tested these effects on individuals from the general population, who on average, have normal serum calcium levels. Thus, these findings can only provide insight into the effect of further increases in serum calcium levels in eucalcemic individuals. Most individuals studied for bone mineral density and fracture outcomes did not have osteoporosis as defined by bone mineral density measurement. Thus, the effects of genetically increased calcium in such individuals should be tested separately.

Most randomised controlled trials for prevention of fracture have used calcium and vitamin D supplements in conjunction with fracture preventive therapies and it is not clear whether giving such drugs in the absence of calcium supplements would provide the same protective benefits as were shown in these randomised controlled trials. Nonetheless, many randomised controlled trials for fracture prevention have given calcium and vitamin D supplements in the control arm of the study. Further, a recent randomised controlled trial of zoledronate showed marked reductions in the risk of fracture without the use of calcium.⁴⁰

Regarding study populations, there is no overlap between the fracture genome-wide association study and the estimated bone mineral density genome-wide association study. Fracture genome-wide association

Fig 4 | Serum calcium effects on odds of fracture excluding cohort with Asian descent: sensitivity analysis. Two sample mendelian randomisation: individual and inverse variance weighted (IVW) results

study overlapped in a 2.8% with the calcium genome-wide association study population. However, we do not expect a substantial impact given the low degree of overlap.⁸⁵ Confounding by ancestry, also known as population stratification, can bias mendelian randomisation studies. A method to overcome such confounding is to limit the study to people of the same ancestry. Although most individuals in the fracture study were of European ancestry, ~1% of individuals were of Asian descent. However, removal of this Asian cohort did not impact on our results.

Our study provides insights into serum calcium levels and not tissue level concentrations. While these different compartments of calcium homeostasis might have different effects on the risk of fracture, calcium supplementation acts on the skeleton by first influencing serum calcium. Thus, the results presented here can provide insight into the expected effects of calcium supplementation by serum calcium. Further, the mendelian randomisation estimate for fracture, a binary outcome, was expressed as an odds ratio, which is a non-collapsible measure, yet this estimator still provides a valid test of the null hypothesis.⁸⁶ Canalization, which is the sum of compensatory feedback mechanisms returning a physiological system to homeostasis, can bias mendelian randomisation results towards the null. However, it is plausible the same mechanisms which maintain calcium homeostasis would act in a similar fashion on serum calcium raised by supplementation and by genetic effects. Further, the genetic predisposition to increased serum calcium used in our study is of sufficient biological and clinical relevance because of its association with increased risk of coronary artery disease.²²

Conclusions

A genetic predisposition to increased serum calcium, amongst individuals with normal calcium levels, was not associated with increased estimated bone mineral density or decreased risk of fracture. The degree to which lifelong genetically derived increased serum calcium mimics the effect of long term calcium supplementation is not known. Since genetically elevated serum calcium is strongly associated with an increased risk of coronary artery disease, widespread calcium supplementation in the general population does not appear to have a favourable risk-benefit profile.

AUTHOR AFFILIATIONS

¹Lady Davis Institute, Jewish General Hospital, McGill University, Montréal, Québec, Canada

²Department of Epidemiology, Biostatistics and Occupational Health, McGill University, 3755 Côte Ste-Catherine Road, Suite H-413, Montréal, Québec, H3T 1E2, Canada

³Department of Human Genetics, McGill University, Montréal, Québec, Canada

⁴Department of Internal Medicine, Erasmus MC, University Medical Center, Rotterdam, Netherlands

⁵Department of Epidemiology, Erasmus MC, University Medical Center, Rotterdam, Netherlands

⁶Unit of Nutritional Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

⁷Department of Surgical Sciences, Uppsala University, Uppsala, Sweden

We thank the individuals who participated in the studies that made this study possible.

Contributors: AC, SZ, and JBR were involved in the design of the study. AC and SZ carried out statistical analysis and wrote the first draft of the manuscript with JBR's support. All authors participated in further drafts and approved final manuscript. The corresponding author confirms that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted. AC is the guarantor.

Funding: JBR and AC are supported by CIHR and FRQS. The funding sources had no role in: a. the design or conduct of the study, b. the collection, management, analysis, and interpretation of the data, or c. preparation, review, or approval of the manuscript. The Richards research group is supported by the Canadian Institutes of Health Research (CIHR), the Lady Davis Institute of the Jewish General Hospital, the Canadian Foundation for Innovation and the Fonds de Recherche Québec Santé (FRQS). JBR is supported by a FRQS Clinical Research Scholarship. TwinsUK is funded by the Wellcome Trust, Medical Research Council, European Union, the National Institute for Health Research (NIHR)-funded BioResource, Clinical Research Facility and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership with King's College London. AC and JAM are funded by CIHR's Frederick Banting and Charles Best Canada Graduate Scholarships Doctoral Award and AC was funded by FRQS's Doctoral Award. SZ is funded by a CIHR Fellowship. This research and the Genetic Factors for Osteoporosis (GEFOS) consortium have been funded by the European Commission (HEALTH-F2-2008-201865-GEFOS). KT and FR are supported by the Netherlands Scientific Organization (NWO) and ZonMW Project number: NW O/ZONMW-VIDI-0 16-136-367. This research has been conducted using the UK Biobank Resource (project number: 24268) and GEFOS.

Competing interest statement: All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Ethical approval: No separate ethical approval was required due to the use of publicly available summary data.

Data sharing: No additional data are available.

The manuscript's guarantor (AC) affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned ((and, if relevant, registered)) have been explained.

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

- 1 Riggs BL, Melton LJ3rd. The worldwide problem of osteoporosis: insights afforded by epidemiology. *Bone* 1995;17(Suppl):505S-11S. doi:10.1016/8756-3282(95)00258-4
- 2 Center JR, Nguyen TV, Schneider D, Sambrook PN, Eisman JA. Mortality after all major types of osteoporotic fracture in men and women: an observational study. *Lancet* 1999;353:878-82. doi:10.1016/S0140-6736(98)09075-8
- 3 Papaioannou A, Morin S, Cheung AM, et al, Scientific Advisory Council of Osteoporosis Canada. 2010 clinical practice guidelines for the diagnosis and management of osteoporosis in Canada: summary. *CMAJ* 2010;182:1864-73. doi:10.1503/cmaj.100771
- 4 Compston J, Cooper A, Cooper C, et al, National Osteoporosis Guideline Group (NOGG). UK clinical guideline for the prevention and treatment of osteoporosis. *Arch Osteoporos* 2017;12:43. doi:10.1007/s11657-017-0324-5
- 5 Cosman F, de Beur SJ, LeBoff MS, et al, National Osteoporosis Foundation. Clinician's Guide to Prevention and Treatment of Osteoporosis. *Osteoporos Int* 2014;25:2359-81. doi:10.1007/s00198-014-2794-2
- 6 Kantor ED, Rehm CD, Du M, White E, Giovannucci EL. Trends in Dietary Supplement Use Among US Adults From 1999-2012. *JAMA* 2016;316:1464-74. doi:10.1001/jama.2016.14403
- 7 Ross AC, Taylor CL, Yaktine AL, et al, eds. *Dietary Reference Intakes for Calcium and Vitamin D*. 2011.

- 8 Rooney MR, Michos ED, Hootman KC, Harnack L, Lutsey PL. Trends in calcium supplementation, National Health and Nutrition Examination Survey (NHANES) 1999-2014. *Bone* 2018;111:23-7. doi:10.1016/j.bone.2018.03.007
- 9 Bailey RL, Dodd KW, Goldman JA, et al. Estimation of total usual calcium and vitamin D intakes in the United States. *J Nutr* 2010;140:817-22. doi:10.3945/jn.109.118539
- 10 Reid IR, Birstow SM, Bolland MJ. Calcium and Cardiovascular Disease. *Endocrinol Metab (Seoul)* 2017;32:339-49. doi:10.3803/EnM.2017.32.3.339
- 11 Bolland MJ, Grey A, Reid IR. Calcium supplements and cardiovascular risk: 5 years on. *Ther Adv Drug Saf* 2013;4:199-210. doi:10.1177/2042098613499790
- 12 Reid IR, Gamble GD, Bolland MJ. Circulating calcium concentrations, vascular disease and mortality: a systematic review. *J Intern Med* 2016;279:524-40. doi:10.1111/joim.12464
- 13 Foley RN, Collins AJ, Ishani A, Kalra PA. Calcium-phosphate levels and cardiovascular disease in community-dwelling adults: the Atherosclerosis Risk in Communities (ARIC) Study. *Am Heart J* 2008;156:556-63. doi:10.1016/j.ahj.2008.05.016
- 14 Rohrmann S, Garmo H, Malmström H, et al. Association between serum calcium concentration and risk of incident and fatal cardiovascular disease in the prospective AMORIS study. *Atherosclerosis* 2016;251:85-93. doi:10.1016/j.atherosclerosis.2016.06.004
- 15 Bolland MJ, Avenell A, Baron JA, et al. Effect of calcium supplements on risk of myocardial infarction and cardiovascular events: meta-analysis. *BMJ* 2010;341:c3691. doi:10.1136/bmj.c3691
- 16 Bolland MJ, Grey A, Avenell A, Gamble GD, Reid IR. Calcium supplements with or without vitamin D and risk of cardiovascular events: reanalysis of the Women's Health Initiative limited access dataset and meta-analysis. *BMJ* 2011;342:d2040. doi:10.1136/bmj.d2040
- 17 Wang L, Manson JE, Song Y, Sesso HD. Systematic review: Vitamin D and calcium supplementation in prevention of cardiovascular events. *Ann Intern Med* 2010;152:315-23. doi:10.7326/0003-4819-152-5-201003020-00010
- 18 Chung M, Tang AM, Fu Z, Wang DD, Newberry SJ. Calcium Intake and Cardiovascular Disease Risk: An Updated Systematic Review and Meta-analysis. *Ann Intern Med* 2016;165:856-66. doi:10.7326/M16-1165
- 19 Lewis JR, Radavelli-Bagatini S, Rejnmark L, et al. The effects of calcium supplementation on verified coronary heart disease hospitalization and death in postmenopausal women: a collaborative meta-analysis of randomized controlled trials. *J Bone Miner Res* 2015;30:165-75. doi:10.1002/jbmr.2311
- 20 Smith GD, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol* 2003;32:1-22. doi:10.1093/ije/dyg070
- 21 Hemani G, Bowden J, Davey Smith G. Evaluating the potential role of pleiotropy in Mendelian randomization studies. *Hum Mol Genet* 2018;27(R2):R195-208. doi:10.1093/hmg/ddy163
- 22 Larsson SC, Burgess S, Michaëlsson K. Association of Genetic Variants Related to Serum Calcium Levels With Coronary Artery Disease and Myocardial Infarction. *JAMA* 2017;318:371-80. doi:10.1001/jama.2017.8981
- 23 Fischer PR, Thacher TD, Pettifor JM. Pediatric vitamin D and calcium nutrition in developing countries. *Rev Endocr Metab Disord* 2008;9:181-92. doi:10.1007/s1154-008-9085-1
- 24 Munns CF, Shaw N, Kiely M, et al. Global Consensus Recommendations on Prevention and Management of Nutritional Rickets. *J Clin Endocrinol Metab* 2016;101:394-415. doi:10.1210/jc.2015-2175
- 25 Chapman T, Sugar N, Done S, Marasigan J, Wambold N, Feldman K. Fractures in infants and toddlers with rickets. *Pediatr Radiol* 2010;40:1184-9. doi:10.1007/s00247-009-1470-8
- 26 Gifre L, Peris P, Monegal A, Martinez de Osaba MJ, Alvarez L, Guañabens N. Osteomalacia revisited : a report on 28 cases. *Clin Rheumatol* 2011;30:639-45. doi:10.1007/s10067-010-1587-z
- 27 Bhan A, Rao AD, Rao DS. Osteomalacia as a result of vitamin D deficiency. *Endocrinol Metab Clin North Am* 2010;39:321-31. doi:10.1016/j.ecl.2010.02.001
- 28 Basha B, Rao DS, Han ZH, Parfitt AM. Osteomalacia due to vitamin D depletion: a neglected consequence of intestinal malabsorption. *Am J Med* 2000;108:296-300. doi:10.1016/S0002-9343(99)00460-X
- 29 Rosen CJ, Compston JE, Lian JB, Lips P, van Schoor NM, Bravenboer N. Vitamin D-related disorders. In: *ASBMR primer on the metabolic bone diseases and disorders of mineral metabolism*. John Wiley & Sons, 2009.
- 30 Bhambr R, Naik V, Malhotra N, et al. Changes in bone mineral density following treatment of osteomalacia. *J Clin Densitom* 2006;9:120-7. doi:10.1016/j.jocd.2005.11.001
- 31 Allen SC, Raut S. Biochemical recovery time scales in elderly patients with osteomalacia. *J R Soc Med* 2004;97:527-30. doi:10.1177/014107680409701104
- 32 Thacher TD, Fischer PR, Pettifor JM, et al. A comparison of calcium, vitamin D, or both for nutritional rickets in Nigerian children. *N Engl J Med* 1999;341:563-8. doi:10.1056/NEJM199908193410803
- 33 Moyer VAUS. Preventive Services Task Force*. Vitamin D and calcium supplementation to prevent fractures in adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2013;158:691-6. doi:10.7326/0003-4819-158-6-201303190-00588
- 34 Jackson RD, LaCroix AZ, Gass M, et al. Women's Health Initiative Investigators. Calcium plus vitamin D supplementation and the risk of fractures. *N Engl J Med* 2006;354:669-83. doi:10.1056/NEJMoa055218
- 35 Grant AM, Avenell A, Campbell MK, et al. RECORD Trial Group. Oral vitamin D3 and calcium for secondary prevention of low-trauma fractures in elderly people (Randomised Evaluation of Calcium Or vitamin D, RECORD): a randomised placebo-controlled trial. *Lancet* 2005;365:1621-8. doi:10.1016/S0140-6736(05)63013-9
- 36 Porthouse J, Cockayne S, King C, et al. Randomised controlled trial of calcium and supplementation with cholecalciferol (vitamin D3) for prevention of fractures in primary care. *BMJ* 2005;330:1003. doi:10.1136/bmj.330.7498.1003
- 37 Salovaara K, Tuppurainen M, Kärkkäinen M, et al. Effect of vitamin D(3) and calcium on fracture risk in 65- to 71-year-old women: a population-based 3-year randomized, controlled trial-the OSTPRE-FPS. *J Bone Miner Res* 2010;25:1487-95. doi:10.1002/jbmr.48
- 38 Bolland MLW, Tai V, Bastin S, Gamble G, Grey A, Reid I. Systematic review of calcium intake and risk of fracture. *BMJ* 2015. doi:10.1136/bmj.h4580
- 39 Bischoff-Ferrari HA, Dawson-Hughes B, Baron JA, et al. Calcium intake and hip fracture risk in men and women: a meta-analysis of prospective cohort studies and randomized controlled trials. *Am J Clin Nutr* 2007;86:1780-90. doi:10.1093/ajcn/86.6.1780
- 40 Reid IR, Horne AM, Mihov B, et al. Fracture prevention with zoledronate in older women with osteopenia. *N Engl J Med* 2018;379:2407-16. doi:10.1056/NEJMoa1808082
- 41 Bonnick S, Brody S, Kaiser F, et al. Treatment with alendronate plus calcium, alendronate alone, or calcium alone for postmenopausal low bone mineral density. *Curr Med Res Opin* 2007;23:1341-9. doi:10.1185/030079907X188035
- 42 Dastani Z, Hivert MF, Timpson N, et al. DIAGRAM+ Consortium, MAGIC Consortium, GLGC Investigators, MuTHER Consortium, DIAGRAM Consortium, GIANT Consortium, Global B Pgen Consortium, Procardis Consortium, MAGIC investigators, GLGC Consortium. Novel loci for adiponectin levels and their influence on type 2 diabetes and metabolic traits: a multi-ethnic meta-analysis of 45,891 individuals. *PLoS Genet* 2012;8:e1002607. doi:10.1371/journal.pgen.1002607
- 43 Holmes MV, Ala-Korpela M, Smith GD. Mendelian randomization in cardiometabolic disease: challenges in evaluating causality. *Nat Rev Cardiol* 2017;14:577-90. doi:10.1038/nrcardio.2017.78
- 44 O'Seaghdha CM, Wu H, Yang Q, et al. SUNLIGHT Consortium, GEFOS Consortium. Meta-analysis of genome-wide association studies identifies six new loci for serum calcium concentrations. *PLoS Genet* 2013;9:e1003796. doi:10.1371/journal.pgen.1003796
- 45 Maresz K. Proper Calcium Use: Vitamin K2 as a Promoter of Bone and Cardiovascular Health. *Integr Med (Encinitas)* 2015;14:34-9.
- 46 Bügel S. Vitamin K and bone health. *Proc Nutr Soc* 2003;62:839-43. doi:10.1079/PNS2003305
- 47 Morris JA, Kemp JP, Youten SE, et al. 23andMe Research Team. An atlas of genetic influences on osteoporosis in humans and mice. *Nat Genet* 2019;51:258-66. doi:10.1038/s41588-018-0302-x
- 48 McCloskey EV, Kanis JA, Odén A, et al. Predictive ability of heel quantitative ultrasound for incident fractures: an individual-level meta-analysis. *Osteoporos Int* 2015;26:1979-87. doi:10.1007/s00198-015-3072-7
- 49 Bauer DC, Glüer CC, Cauley JA, et al. Study of Osteoporotic Fractures Research Group. Broadband ultrasound attenuation predicts fractures strongly and independently of densitometry in older women. A prospective study. *Arch Intern Med* 1997;157:629-34. doi:10.1001/archinte.1997.00440270067006
- 50 Morris JA, Kemp JP, Youten SE, et al. 23andMe Research Team. An atlas of genetic influences on osteoporosis in humans and mice. *Nat Genet* 2019;51:258-66.
- 51 McCarthy S, Das S, Kretzschmar W, et al. Haplotype Reference Consortium. A reference panel of 64,976 haplotypes for genotype imputation. *Nat Genet* 2016;48:1279-83. doi:10.1038/ng.3643
- 52 Trajanoska K, Morris JA, Oei L, et al. GEFOS/GENOMOS consortium and the 23andMe research team. Assessment of the genetic and clinical determinants of fracture risk: genome wide association and mendelian randomisation study. *BMJ* 2018;362:k3225. doi:10.1136/bmj.k3225

- 53 Loh PR, Tucker G, Bulik-Sullivan BK, et al. Efficient Bayesian mixed-model analysis increases association power in large cohorts. *Nat Genet* 2015;47:284-90. doi:10.1038/ng.3190
- 54 Loh P-R. BOLT-LMM v2.3.2 User Manual 2018 [updated March 10, 2018. Available from: <https://data.broadinstitute.org/alkesgroup/BOLT-LMM/>.
- 55 Kemp JP, Morris JA, Medina-Gomez C, et al. Identification of 153 new loci associated with heel bone mineral density and functional involvement of GPC6 in osteoporosis. *Nat Genet* 2017;49:1468-75. doi:10.1038/ng.3949
- 56 Willer CJ, Li Y, Abecasis GR, et al. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 2010;26:2190-1. doi:10.1093/bioinformatics/btq340
- 57 Staley JR, Blackshaw J, Kamat MA, et al. PhenoScanner: a database of human genotype-phenotype associations. *Bioinformatics* 2016;32:3207-9. doi:10.1093/bioinformatics/btw373
- 58 Consortium GTGEx Consortium. The Genotype-Tissue Expression (GTEx) project. *Nat Genet* 2013;45:580-5. doi:10.1038/ng.2653
- 59 Robinson-Cohen C, Lutsey PL, Kleber ME, et al. Genetic Variants Associated with Circulating Parathyroid Hormone. *J Am Soc Nephrol* 2017;28:1553-65. doi:10.1681/ASN.2016010069
- 60 Bristow SM, Gamble GD, Stewart A, Kalluru R, Horne AM, Reid IR. Acute effects of calcium citrate with or without a meal, calcium-fortified juice and a dairy product meal on serum calcium and phosphate: a randomised cross-over trial. *Br J Nutr* 2015;113:1585-94. doi:10.1017/S000711451500080X
- 61 Baiocchi M, Cheng J, Small DS. Instrumental variable methods for causal inference. *Stat Med* 2014;33:2297-340. doi:10.1002/sim.6128
- 62 R: A language and environment for statistical computing [program]. Vienna, Austria, 2018. <https://www.gbif.org/en/tool/81287/r-a-language-and-environment-for-statistical-computing>
- 63 Yavorska OO, Burgess S. MendelianRandomization: an R package for performing Mendelian randomization analyses using summarized data. *Int J Epidemiol* 2017;46:1734-9. doi:10.1093/ije/dyx034
- 64 Team R. RStudio: Integrated Development for R. *RStudio, Inc, Boston, MA* 2015
- 65 Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet Epidemiol* 2016;40:304-14. doi:10.1002/gepi.21965
- 66 Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol* 2015;44:512-25. doi:10.1093/ije/dyv080
- 67 Hendy GN, D'Souza-Li L, Yang B, Canaff L, Cole DE. Mutations of the calcium-sensing receptor (CASR) in familial hypocalciuric hypercalcemia, neonatal severe hyperparathyroidism, and autosomal dominant hypocalcemia. *Hum Mutat* 2000;16:281-96. doi:10.1002/1098-1004(200010)16:4<281::AID-HUMU1>3.0.CO;2-A
- 68 Liu PT, Stenger S, Li H, et al. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* 2006;311:1770-3. doi:10.1126/science.1123933
- 69 Okada Y, Imendra KG, Miyazaki T, Hotokezaka H, Fujiyama R, Toda K. High extracellular Ca²⁺ stimulates Ca²⁺-activated Cl⁻ currents in frog parathyroid cells through the mediation of arachidonic acid cascade. *PLoS One* 2011;6:e19158. doi:10.1371/journal.pone.0019158
- 70 Van Esch H, Groenen P, Nesbit MA, et al. GATA3 haplo-insufficiency causes human HDR syndrome. *Nature* 2000;406:419-22. doi:10.1038/35019088
- 71 Kang TW, Kim HJ, Ju H, et al. Genome-wide association of serum bilirubin levels in Korean population. *Hum Mol Genet* 2010;19:3672-8. doi:10.1093/hmg/ddq281
- 72 Johnson AD, Kavousi M, Smith AV, et al. Genome-wide association meta-analysis for total serum bilirubin levels. *Hum Mol Genet* 2009;18:2700-10. doi:10.1093/hmg/ddp202
- 73 Shin SY, Fauman EB, Petersen AK, et al. Multiple Tissue Human Expression Resource (MuTHER) Consortium. An atlas of genetic influences on human blood metabolites. *Nat Genet* 2014;46:543-50. doi:10.1038/ng.2982
- 74 Smith DL, Shire NJ, Watts NB, Schmitter T, Szabo G, Zucker SD. Hyperbilirubinemia is not a major contributing factor to altered bone mineral density in patients with chronic liver disease. *J Clin Densitom* 2006;9:105-13. doi:10.1016/j.jocd.2005.10.001
- 75 Lu Y, Vitart V, Burdon KP, et al. NEIGHBOR Consortium. Genome-wide association analyses identify multiple loci associated with central corneal thickness and keratoconus. *Nat Genet* 2013;45:155-63. doi:10.1038/ng.2506
- 76 Willer CJ, Schmidt EM, Sengupta S, et al. Global Lipids Genetics Consortium. Discovery and refinement of loci associated with lipid levels. *Nat Genet* 2013;45:1274-83. doi:10.1038/ng.2797
- 77 Speliotes EK, Yerges-Armstrong LM, Wu J, et al. NASH CRN, GIANT Consortium, MAGIC Investigators, GOLD Consortium. Genome-wide association analysis identifies variants associated with nonalcoholic fatty liver disease that have distinct effects on metabolic traits. *PLoS Genet* 2011;7:e1001324. doi:10.1371/journal.pgen.1001324
- 78 Kraja AT, Vaidya D, Pankow JS, et al. A bivariate genome-wide approach to metabolic syndrome: STAMPEED consortium. *Diabetes* 2011;60:1329-39. doi:10.2337/db10-1011
- 79 Moayyeri A, Adams JE, Adler RA, et al. Quantitative ultrasound of the heel and fracture risk assessment: an updated meta-analysis. *Osteoporos Int* 2012;23:143-53. doi:10.1007/s00198-011-1817-5
- 80 Wang H, Bua P, Capodice J. A comparative study of calcium absorption following a single serving administration of calcium carbonate powder versus calcium citrate tablets in healthy premenopausal women. *Food Nutr Res* 2014;58. doi:10.3402/fnr.v58.23229
- 81 Wang Y, Huang Z, Yi B. Calcium and Vitamin D Supplements and Fractures in Community-Dwelling Adults. *JAMA* 2018;319:2042. doi:10.1001/jama.2018.3919
- 82 Lawlor DA, Tilling K, Davey Smith G. Triangulation in aetiological epidemiology. *Int J Epidemiol* 2016;45:1866-86. doi:10.1093/ije/dyw314
- 83 Munafò MR, Davey Smith G. Robust research needs many lines of evidence. *Nature* 2018;553:399-401. doi:10.1038/d41586-018-01023-3
- 84 Ding M, Huang T, Bergholdt HK, Nordestgaard BG, Ellervik C, Qi L. CHARGE Consortium. Dairy consumption, systolic blood pressure, and risk of hypertension: Mendelian randomization study. *BMJ* 2017;356:j1000. doi:10.1136/bmj.j1000
- 85 Burgess S, Davies NM, Thompson SG. Bias due to participant overlap in two-sample Mendelian randomization. *Genet Epidemiol* 2016;40:597-608. doi:10.1002/gepi.21998
- 86 Burgess S, Small DS, Thompson SG. A review of instrumental variable estimators for Mendelian randomization. *Stat Methods Med Res* 2017;26:2333-55. doi:10.1177/0962280215597579