

Assessing coverage of interventions for reproductive, maternal, newborn, child, and adolescent health and nutrition

Progress has been made in priority interventions, but we need new measurement systems that include the whole life course and give better assessment of equity of coverage, argue

Jennifer Requejo and colleagues

Embedded within the framework of the United Nations sustainable development goals (SDG) for 2030 is the principle of equity, with the aim of reaching universal health coverage. Soon after the framework was adopted in 2015, the Every Woman Every Child global strategy for women's, children's and adolescents' health was launched.^{1,2} The global strategy translates the SDG agenda into concrete guidance on how to accelerate progress in women's, children's and adolescents' health through a multisectoral approach. It includes a monitoring framework with 60 indicators to help countries and their partners promote accountability in ending preventable deaths (survive), ensuring health and wellbeing (thrive), and expanding enabling environments, so that all women, children, and adolescents can reach their potential (transform).³ Previous assessments show mixed progress, with some indicators advancing more rapidly

than others but with pervasive inequities between and within countries.⁴

The Countdown to 2030 initiative also regularly tracks progress in the countries experiencing the highest burdens of maternal and child mortality. Countdown's 2017 report,⁵ together with those from Every Woman Every Child, highlight laudable reductions in maternal and child mortality over the past two decades, but many settings will need to increase efforts to achieve the 2030 goals. Coverage of essential health interventions is unacceptably low in many contexts and among specific populations; considerable policy and programmatic work is needed to shore up primary healthcare systems to make universal health coverage a reality.⁵

We combine the countdown and global strategy indicators to take stock of progress in all 138 low and middle income countries plus Panama, which was reclassified as a high income country in 2018 but remains a priority country in Countdown to 2030.⁶ We examine how well the world is doing in reaching every woman, child, and adolescent with effective health interventions, how far we need to go to achieve the SDGs, and identify gaps in the data. We also propose revisions to the Countdown chart of key indicators so that it better reflects the dimensions of survive, thrive, and transform in the Every Woman Every Child strategy and the interconnecting links between reproductive, maternal, newborn, child, and adolescent health and nutrition.

Our assessment is based on the common evaluation framework underlying the countdown and global strategy analyses (see web supplement). This framework posits that health outcomes are determined by the ability of healthcare systems to deliver high quality services to all, which, in turn, depends on supportive policies and sufficient resources, including financial,

supplies and equipment, and healthcare workers. The framework also considers contextual factors, such as humanitarian and environmental crises, women's social status, and other political and economic factors that influence access to services, and the independent effects of education and other life opportunities on health.

Understanding gaps and success

Multiple analytical lenses are needed to understand whether women, children, and adolescents worldwide are receiving effective health interventions. Snapshots of current status tell us where we are and how far we need to go to reach global goals. However, it's important to visualise trends to determine whether progress is accelerating, stagnating, or even reversing. Countdown's indicator list helps determine which specific interventions are reaching women, children, and adolescents better than others. Another approach involves ranking countries and regions to identify those that are performing best and those that are lagging behind, providing insight into which countries and regions need constructive action to direct resources and support.

Where are we now?

Countdown's continuum of care chart shows a subset of prioritised indicators of intervention coverage across the dimensions of family planning, pregnancy, childbirth, postnatal care, and childhood, with selected crosscutting indicators for water and sanitation. These indicators reflect population level data, showing how well countries are doing in reaching those in need. Figure 1 shows progress on this chart for all low and middle income countries for 2014-18.^{7,8} Summary data (table 1) show that we are far from achieving universal coverage for many interventions, with larger gaps for family planning services, breastfeeding, and treatment of childhood

KEY MESSAGES

- Despite substantial progress in reducing maternal, newborn, and child mortality worldwide, inequities persist
- Countries in sub-Saharan Africa are lagging most behind
- Coverage of interventions is higher for those that are well resourced, can take place at planned times (such as preventive services), and do not depend on a functioning healthcare system
- The indicators for monitoring progress need to be revised to include proved interventions for older children, adolescents, and adult women
- Further disaggregation of intervention coverage by equity measures is important to better understand who is being left behind




Fig 1 | Median national coverage for selected interventions for women's and child health using most recent WHO and Unicef data for each country (2014 or later; see web supplement for further details).^{7,8} Bars represent the global median and the dots represent country estimates (DTP3=diphtheria-tetanus-pertussis)

illnesses. Over 80% coverage has been achieved for long established immunisation interventions, skilled attendant at delivery, and safe drinking water, yet coverage is below 50% for interventions that require substantial behaviour change (eg, breastfeeding) or have received less consistent political commitment and resources (eg, oral rehydration salts for diarrhoea).

Have we progressed, stumbled, or flatlined?

We evaluated change in coverage over the past 10 years, focusing on countries with data available for 2009-13 and 2014-18 (table 2, fig 2). We included only countries with at least one estimate for each time interval. Thirteen out of the 16 interventions examined show positive change over the past 10 years, especially treatment of pregnant women with HIV, which has benefited from substantial global investment. Three interventions showed no improvement or a decline: diphtheria-tetanus-pertussis (DTP3) vaccination, measles vaccination, and number of children aged 12 to 15 months who are still breastfed. Because rotavirus vaccine is relatively new, immunisation increased substantially between the two periods as countries implemented the policy. However, universal coverage has not been achieved, and coverage levels are around the same as for DTP3 and measles.

The data suggest that immunisation coverage is flatlining. Many countries

experience challenges over vaccines costs, vaccine supply chains, and difficulties reaching the most disadvantaged children in remote or rural areas, or in slums. In some countries parental willingness to vaccinate is also a problem. Although considerable attention has been devoted to understanding vaccine hesitancy in high income settings, most unvaccinated children live in areas without reliable access to them.⁹

Further scrutiny is needed to understand the factors driving positive trends in low and middle income countries as well as those that are inhibiting progress so that appropriate action can be taken. Interventions with high national coverage can start to fall behind as focus shifts elsewhere. Gains need to be sustained through close partnerships between the global community, governments, and local civil society groups.

Best and worst performing countries

Indices including standardised indicators are useful for comparing overall country performance towards specific development goals. Such comparisons can help to identify which countries are progressing and which are falling behind and should trigger additional national and sub-national analyses to understand the factors driving success and failure.

We used the composite coverage index (CCI)¹⁰ to rank countries with available data on every indicator included in the index during 2010-18 (45 countries lack

available data, most of which are middle income countries in Europe, North America, Oceania, and Latin America and the Caribbean). The CCI is an established summary measure for intervention coverage that correlates with key indicators of health status such as child mortality and stunting. It is calculated from data provided by UN member states, so the quality of the measure is only as strong as the underlying data.

The index has four components (reproductive health, maternal health, child immunisation, and treatment of childhood illness), is limited to interventions delivered through healthcare systems, and has eight coverage indicators: access to modern family planning methods, four or more antenatal care visits, skilled attendant at birth, three doses of DTP vaccine, BCG vaccination, first dose of measles vaccine, oral rehydration salts for diarrhoea, and care seeking for acute respiratory infections (see web supplement for details on how the index is calculated).

Figure 3 shows a heatmap of the CCI for the five best and five worst performing countries (a heat map for all low and middle income countries is given in the web supplement). Indicators for the best performers are mostly green, but all have slightly lower coverage for treating childhood illness. The indicators for the lowest performers, in contrast, are mostly red and orange, with some performing better for immunisations. A comprehensive

Table 1 | Coverage of priority indicators for women's and children's health in low and middle income countries* 2014-18⁷

| Indicator | No of countries with data | Coverage | | | Countries with lowest value | Countries with highest value |
|--|---------------------------|---|-------------|-------------|---|---|
| | | Median (interquartile range) coverage (%) | Minimum (%) | Maximum (%) | | |
| Pre-pregnancy | | | | | | |
| Demand for family planning satisfied with modern methods | 77 | 56 (42-74) | 5 | 90 | Albania | North Korea |
| Pregnancy | | | | | | |
| Antenatal care (4+ visits) | 74 | 73 (53-89) | 18 | 98 | Afghanistan | Cuba |
| Treatment of pregnant women living with HIV | 93 | 79 (46-93) | 5 | 99 | Sudan | Benin, Bolivia, Cuba, Grenada, Jamaica, Mozambique, Mauritius, Malawi, Malaysia, Namibia, Romania, Rwanda, Sao Tome and Principe, Suriname, Saint Vincent and the Grenadines, Zambia |
| Neonatal tetanus protection | 99 | 90 (85-94) | 60 | 99 | Central African Republic, Nigeria | Dominican Republic, Eritrea, Guyana, Honduras, Sri Lanka, Maldives, Sao Tome and Principe |
| Birth | | | | | | |
| Skilled attendant at delivery | 109 | 92 (74-99) | 20 | 100 | Chad | Turkmenistan, Uzbekistan |
| Postnatal | | | | | | |
| Postnatal visit for mothers | 68 | 71 (50-90) | 1 | 100 | Colombia | Turkmenistan |
| Postnatal visit for babies | 61 | 64 (32-91) | 4 | 100 | Chad | Turkmenistan |
| Early initiation of breastfeeding | 68 | 54 (41-67) | 20 | 90 | Pakistan | Sri Lanka |
| Infancy | | | | | | |
| Exclusive breastfeeding (<6 months) | 68 | 46 (35-58) | 0 | 87 | Chad | Rwanda |
| Continued breastfeeding (year 1) | 67 | 80 (60-90) | 25 | 98 | Serbia | Nepal |
| DTP3 immunisation | 137 | 90 (81-96) | 25 | 99 | Equatorial Guinea | Albania, China, Cuba, Fiji, Iran, Sri Lanka, Morocco, Maldives, Mongolia, Malaysia, Turkmenistan |
| Measles immunisation (first dose) | 137 | 89 (78-96) | 30 | 99 | Equatorial Guinea | China, Cape Verde, Cuba, Eritrea, Iran, Kazakhstan, Sri Lanka, Morocco, Maldives, Mongolia, Mauritius, Nicaragua, Nauru, Rwanda, Turkmenistan, Tanzania, Saint Vincent and the Grenadines |
| Rotavirus immunisation | 73 | 85 (74-92) | 2 | 99 | Philippines | Fiji, Morocco |
| Childhood | | | | | | |
| Vitamin A supplementation (two doses) | 66 | 62 (26-90) | 0 | 99 | Burkina Faso, Central African Republic, Gabon, Mauritania | Benin, Bangladesh, Morocco, Uzbekistan, Vietnam, Zambia |
| Care seeking for pneumonia symptoms | 66 | 63 (49-77) | 24 | 93 | Nigeria | Cuba |
| Diarrhoea: oral rehydration treatment | 64 | 44 (34-55) | 16 | 84 | Cameroon | Eswatini |
| Environment | | | | | | |
| Population using basic drinking water services | 138 | 91 (69-97) | 39 | 100 | Chad | Romania |
| Population using basic sanitation services | 135 | 76 (39-91) | 7 | 100 | Ethiopia | Libya, Uzbekistan |

*Includes Panama, which was reclassified as a high income country in 2018. Data from WHO and Unicef on immunisation rates, July 2019; WHO and Unicef Joint Monitoring Programme for Water Supply and Sanitation, June 2019; Joint United Nations Programme on HIV/AIDS, July 2019; and Unicef (www.data.unicef.org) and WHO global databases (www.who.int/gho/database/en/), May 2019 (based on demographic and health surveys, multiple indicator cluster surveys, and other national surveys)

approach is clearly needed for these countries to increase access to care for women and children.

The best performers are spread across different world regions, but apart from Yemen the lowest performers are all in Africa, suggesting that greater attention needs to be directed there. Comparing these countries' performance on the CCI to mortality statistics provides support for the role of this evaluation framework and the role of high intervention coverage in improving survival. The highest performers have all achieved considerable reductions

in maternal and child mortality over the past 10 years.^{11 12} The poorest performers, in contrast, have made less progress in improving maternal and child survival; all but Equatorial Guinea have levels of stunting of at least 30%, and four of the five (Central African Republic, Chad, South Sudan, and Yemen) are affected by conflict or are fragile states^{13 14}

Where do we go from here?

Global monitoring involves regular tracking of sentinel indicators to assess progress towards fixed goals and targets. But global

monitoring is also an iterative process, staying on track with changes in the evidence on effective interventions, implementation strategies, demographic and epidemiological trends, and country priorities. Thus, indicators monitored through global initiatives need to be regularly reviewed and updated.

The SDG and Every Woman Every Child strategy frameworks use a life course approach to development, which has been reflected in strategic plans, programming, and research agendas. The large reduction in child mortality¹¹ has enabled a shift from a focus on reducing deaths from infectious

Table 2 | Changes in median national coverage of selected interventions in low and middle income countries with available data for 2009-13 and 2014-18, ordered by proportion of the gap closed

| Stage | Indicator | No of countries | 2009-13 median | 2014-18 median | Change (% points) | % of gap closed |
|---------------|--|-----------------|----------------|----------------|-------------------|-----------------|
| Pregnancy | Treatment of pregnant women with HIV | 93 | 10 | 79 | 69 | 77 |
| Infancy | Rotavirus immunisation | 34 | 69 | 87 | 18 | 58 |
| Postnatal | Postnatal visit for babies | 31 | 30 | 68 | 38 | 54 |
| Pregnancy | Neonatal tetanus protection | 99 | 85 | 90 | 5 | 33 |
| Pregnancy | ≥4 antenatal contacts | 63 | 60 | 73 | 13 | 32 |
| Birth | Skilled attendant delivery | 95 | 90 | 93 | 3 | 30 |
| Postnatal | Postnatal visit for mothers | 38 | 44 | 59 | 15 | 27 |
| Environment | Population using basic drinking water services | 138 | 88 | 91 | 3 | 25 |
| Infancy | Exclusive breastfeeding (<6 months) | 52 | 38 | 46 | 8 | 13 |
| Childhood | Care seeking for pneumonia symptoms | 54 | 57 | 62 | 5 | 12 |
| Pre-pregnancy | Demand for family planning satisfied with modern methods | 62 | 51 | 55 | 4 | 8 |
| Postnatal | Early initiation of breastfeeding | 53 | 50 | 54 | 4 | 8 |
| Childhood | Oral replacement therapy for diarrhoea | 56 | 38 | 42 | 4 | 6 |
| Infancy | Continued breastfeeding to 1 year | 51 | 81 | 80 | -1 | -5 |
| Infancy | DTP3 immunisation | 137 | 91 | 90 | -1 | -11 |
| Infancy | Measles immunisation (first dose) | 137 | 91 | 89 | -2 | -22 |

Most recent data point for the 2014-18 period and earliest data point for the 2009-2013 period. For countries that had more than one survey during 2009-13, we selected the survey that was closest to the midpoint of 2011. If surveys were equidistant in time to the midpoint, we chose the earlier survey. See web supplement for dates and information on how the percentage coverage gap closed was calculated.

causes and malnutrition in young children to tackling health and development needs throughout the first two decades of life.

Improvements in maternal mortality from direct causes means that indirect causes such as reduced fertility and women having their first pregnancies at older ages are becoming more important.¹² The focus of action is therefore shifting from improving access to intrapartum and emergency obstetric care to include women's experiences of care, the quality of available services, short and long term forms of maternal morbidity and

disability, and women's empowerment, social status, and rights.

Available evidence also shows huge variations in countries' progress on the epidemiological, demographic, and nutritional transitions and hence "one size fits all" approaches to development will not work. Poverty reduction is not a straight trajectory, and success can be precarious. Conflict, global recessions, in-country economic crises, environmental disasters, and epidemics can all cause massive destruction, particularly if resilience and

safety nets are not built into health and other systems. Several countries were downgraded, for example, in the 2019 World Bank income classifications, with implications for the resources they have for social protection, health, and other welfare programmes.⁶ Hence, the international community and country governments must remain vigilant and rapidly respond to deteriorating situations as part of the pledge to achieve sustainable development.

The challenge in selecting core indicators for global monitoring is to keep the number




Fig 2 | Median national coverage of selected interventions for women's and child health for 2009-13 and 2014-18, among countries with data for both periods (DTP3=diphtheria-tetanus-pertussis)




Fig 3 | Heatmap of the composite coverage index (CCI) for the five best and worst performing low and middle income countries with available data, 2010-18, showing values for constituent indicators and four components of the index

small so that reporting burdens on countries are manageable yet comprehensive enough to be relevant and spur action. We cannot lose sight of the fact that the leading killers of young children stubbornly remain pneumonia, diarrhoea, and malaria (in endemic areas), often underpinned by undernutrition. But the indicator set needs to encompass children aged 5-9 years, along with other emerging priorities relating to chronic diseases, disabilities, injuries, violence, and child development.

Adolescent health has garnered considerable attention in recent years.¹⁵ However, only a few adolescent specific indicators have been included in global monitoring frameworks, mainly around reproductive and maternal health, and most are not included as part of routine data collection activities, which hampers accountability.¹⁶

In response to evidence documenting an increasing proportion of child deaths in the neonatal period, numerous initiatives have aimed to improve monitoring around maternal and newborn health, including health systems, community, and broader social, political, and environmental determinants, and the design and implementation of effective programmes.^{17,18} This has also prompted greater focus on the quality of care provided before, during, and immediately after delivery and motivated the development of guidance on assessing health facility

readiness to provide high quality care and health workers' performance.¹⁹

The importance of a life course approach has become internationally recognised, as has the salience of viewing women's, children's, and adolescents' health as interlinked and having intergenerational effects. However, data collection efforts are a few steps behind, with data gaps and measurement challenges persisting. Even for those indicators that cover well established interventions, many countries are missing data from the past five years (tables 1 and 2). Most of the 139 countries have modelled data for established interventions such as immunisations, water and sanitation, and HIV. The number of countries with household survey based data range from 61 (for postnatal care for babies) to 109 (for skilled attendant at birth). This wide range in data availability is due to a combination of factors—some of the interventions are new or have never been included in country programmes or data collection processes (eg, rotavirus vaccine, postnatal care for babies) but many of the 139 countries have not conducted a household survey in the past five years. As countries develop and move from low income to middle income status, they often decide to carry out their own national household surveys, which do not always follow standardised methods (such as the international demographic and health surveys). These surveys generate

non-comparable data, limiting inclusion in global databases. Conflict or other circumstances can also preclude nationally representative household surveys, and some countries have not provided the resources to carry out surveys.

Global stewardship is now needed on several fronts:

- To increase data collection efforts and to generate more comparable data
- To increase investments in countries' health information systems and promote better use of existing data
- To coordinate existing initiatives on updating core lists of indicators to monitor progress towards the 2030 goals
- To refine and move forward the measurement agenda, and
- To push for accountability so that every woman, child, and adolescent gets the care they deserve.

Promising efforts are under way to close data gaps for women, children, and adolescents.^{16,20} WHO and Unicef have engaged experts to reach consensus on how to define and measure effective coverage in order to better capture the potential effect health services can have on health and nutrition outcomes.²¹ Many of the indicators included in global monitoring frameworks and in the continuum of care chart (fig 1) capture information on service contacts and do not provide information on quality of care, which is desperately needed.



This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>.



A consultative process for revising the continuum of care chart should incorporate all these changes on the global landscape, technical advancements in measurement, and country realities. Both the Countdown and Every Woman Every Child indicator lists were derived from lengthy consultative processes involving academia, civil society, UN agencies, healthcare professionals, and countries' ministries of health.^{3,5} A similar process, soon to be instituted, will allow reporting on key indicators of intervention coverage at the midpoint of the SDG period, enabling assessment of whether global progress is on track. This consultative process should be informed by the ongoing work of relevant global accountability and monitoring initiatives.²²⁻²⁴ The process should examine how best to obtain data for missing or under-represented areas such as children aged 5-9, adolescents aged 10-14, adolescent indicators beyond reproductive and maternal health, environmental indicators beyond water and sanitation, and a wider list of reproductive and sexual health indicators as part of health outcomes for adult women.

The revised chart should still be viewed as embedded within the common evaluation framework, serving as a starting point for further analyses focusing on key drivers of coverage such as equity, policy and legislative frameworks, and contextual and health systems factors. A companion chart on quality of care measures is also being developed.²⁵ Unicef and WHO, in partnership with Countdown to 2030, the UN Population Fund (UNFPA), and other UN agencies will lead this consultative process.

Conclusion

Our analyses show that work is needed to achieve universal coverage of important health interventions and to prevent interventions with high coverage levels, such as immunisations, from backsliding. Indicator lists and reporting processes must evolve so that they remain relevant to guide national action and programmatic responses. We have suggested a consultative process for revising Countdown's continuum of care chart to bring it up to date with new thinking on life course approaches and to ensure a manageable set of indicators for reporting on the survive, thrive, and transform dimensions of the Every Woman Every Child strategy. We believe greater investment in data collection is needed so that all countries can report on a core set of indicators, enabling meaningful assessments of global progress.

Finally, further disaggregation of intervention coverage by key markers of equity and the application of newer techniques such as geospatial analyses are essential to improve our understanding of who is being left behind. This information is the starting point for designing strategies to reach all women and children, and for holding all to account for successful implementation.

Competing interests: We have read and understood BMJ policy on declaration of interests and have no relevant interests to declare. The authors alone are responsible for the views expressed in this article, which does not necessarily represent the views, decisions, or policies of the institutions with which the authors are affiliated. Funding from USAID and DFID was used to help support some analysis, graphic design, and to organise references.

Contributors and sources: JR, TD, DC, DJ, ABM, RG, KS conceived the outline of this paper and drafted the first version, LP and AC conducted analysis of data to create the CoC and CCI. DC, DJ, ABM, RG, KS, ACM, LS, JPM, and AB reviewed and revised drafts of the manuscript, and JR and TD created the final version. The data used for the analyses come from publicly available Unicef and WHO global databases, much of which comes from the Demographic and Health Survey (<https://dhsprogram.com>) and Multiple Indicator Cluster Survey (<https://mics.unicef.org/>) programmes.

Data availability: Data are available from the authors on reasonable request, including the underlying data files used to generate the figures and the associated code.

Provenance and peer review: Commissioned; externally peer reviewed.

This article is part of a series proposed by Countdown to 2030 for Women's, Children's and Adolescents' Health and the Partnership for Maternal, Newborn and Child Health (PMNCH) hosted by the World Health Organization and commissioned by *The BMJ*, which peer reviewed, edited, and made the decisions to publish. Open access fees are funded by the Bill and Melinda Gates Foundation and PMNCH.

Jennifer Requejo, senior adviser¹

Theresa Diaz, coordinator²

Lois Park, researcher^{3,4}

Doris Chou, medical officer²

Allysa Choudhury, statistics adviser¹

Regina Guthold, scientist²

Debra Jackson, professor^{1,5}

Ann-Beth Moller, technical officer²

Jean-Pierre Monet, technical officer⁶

Allisyn C Moran, scientist²

Lale Say, coordinator²

Kathleen L Strong, scientist²

Anshu Banerjee, chair^{2,7}

¹Unicef, New York, USA

²World Health Organization, Geneva, Switzerland

³Johns Hopkins University, Baltimore, USA

⁴University of Southern California, Los Angeles, USA

⁵University of Western Cape, Cape Town, South Africa

⁶UNFPA, New York, USA

⁷United Nations H6+ Technical Group, New York, USA,

Correspondence to: J Requejo
jrequejo@unicef.org

- 1 United Nations General Assembly. Transforming our world: the 2030 agenda for sustainable development. New York, 2015.
- 2 Every Woman Every Child, Executive Office of the United Nations Secretary-General. Global Strategy for Women's, Children's and Adolescents' Health (2016-2030). New York, 2015.
- 3 Every Woman Every Child, Executive Office of the United Nations Secretary-General. Indicator and monitoring framework for the global strategy for women's, children's and adolescents' health (2016-2030). New York, 2016.
- 4 World Health Organization, Joint United Nations Programme on HIV/AIDS, United Nations Population Fund, United Nations Children's Fund, UN Women, World Bank Group. Survive, thrive, transform. Global strategy for women's, children's and adolescents' health: 2018 report on progress towards 2030 targets. WHO, 2018.
- 5 Countdown to 2030 Collaboration. Countdown to 2030: tracking progress towards universal coverage for reproductive, maternal, newborn, and child health. *Lancet* 2018;391:1538-48. doi:10.1016/S0140-6736(18)30104-1
- 6 World Bank. Classifying countries by income. 2018. <http://datatopics.worldbank.org/world-development-indicators/stories/the-classification-of-countries-by-income.html>
- 7 United Nations Children's Fund. Unicef data. 2019. <https://data.unicef.org/>
- 8 World Health Organization. Global Health Observatory data repository. 2019. <https://www.who.int/gho/database/en/>
- 9 World Health Organization, United Nations Children's Fund. Progress and challenges with achieving universal immunization coverage: 2018 WHO/Unicef estimates of national immunization coverage. 2019.
- 10 Wehrmeister FC, Restrepo-Mendez MC, Franca GV, Victora CG, Barros AJ. Summary indices for monitoring universal coverage in maternal and child health care. *Bull World Health Organ* 2016;94:903-12. doi:10.2471/BLT.16.173138
- 11 UN Interagency Group for Child Mortality Estimation. Levels and trends in child mortality: report 2018. Estimates developed by the UN interagency group for child mortality estimation. New York, 2018.
- 12 Alkema L, Chou D, Hogan D, et al, UN Maternal Mortality Estimation Inter-Agency Group collaborators and technical advisory group. Global, regional, and national levels and trends in maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a systematic analysis by the UN maternal mortality estimation interagency group. *Lancet* 2016;387:462-74. doi:10.1016/S0140-6736(15)00838-7
- 13 UN Children's Fund, WHO, International Bank for Reconstruction and Development, World Bank. Levels and trends in child malnutrition: key findings of the 2019 edition of the joint child malnutrition estimates. WHO, 2019.
- 14 World Bank Group. *Fragile and conflict affected situations*. World Bank, 2019.
- 15 World Health Organization. Global reference list of health indicators for adolescents (aged 10-19 years). WHO, 2015.
- 16 Guthold R, Moller AB, Azzopardi P, et al. The global action for measurement of adolescent health

- (GAMA) initiative-rethinking adolescent metrics. *J Adolesc Health* 2019;64:697-9. doi:10.1016/j.jadohealth.2019.03.008
- 17 World Health Organization, United Nations Children's Fund. Every newborn action plan: country progress tracking report. WHO, 2016.
 - 18 Moran AC, Jolivet RR, Chou D, et al. A common monitoring framework for ending preventable maternal mortality, 2015-2030: phase I of a multi-step process. *BMC Pregnancy Childbirth* 2016;16:250. doi:10.1186/s12884-016-1035-4
 - 19 World Health Organization. *Standards for improving quality of maternal and newborn care in health facilities*. WHO, 2016.
 - 20 Moller AB, Newby H, Hanson C, et al. Measures matter: a scoping review of maternal and newborn indicators. *PLoS One* 2018;13:e0204763. doi:10.1371/journal.pone.0204763
 - 21 Amouzou A, Leslie HH, Ram M, et al. Advances in the measurement of coverage for RMNCH and nutrition: from contact to effective coverage. *BMJ Glob Health* 2019;4(Suppl 4):e001297. doi:10.1136/bmjgh-2018-001297
 - 22 World Health Organization, United Nations Children's Fund, World Bank. *Nurturing care for early childhood development: a framework for helping children survive and thrive to transform health and human potential*. 2018.
 - 23 Coll-Seck A, Clark H, Bahl R, Peterson S, Costello A, Lucas T. Framing an agenda for children thriving in the SDG era: a WHO-Unicef-Lancet Commission on Child Health and Wellbeing. *Lancet* 2019;393:109-12. doi:10.1016/S0140-6736(18)32821-6
 - 24 World Health Organization. Technical advisory groups on measurement. <https://www.who.int/data/maternal-newborn-child-adolescent>
 - 25 World Health Organization. Quality of care for maternal and newborn health—a monitoring framework for network countries. WHO, 2019.

Web supplement: Additional information on indicator lists for the Every Woman Every Child global strategy and Countdown to 2030, the evaluation framework underlying the analyses, the methods used to prepare tables and figures, plus supplemental figures.

Cite this as: *BMJ* 2020;368:l6915
<http://dx.doi.org/10.1136/bmj.l6915>