¹ Department of Primary Care and Public Health, Imperial College London, London, UK

² UCL Great Ormond Street Institute of Child Health, London, UK

Correspondence to: S Saxena
s.saxena@imperial.ac.uk

Cite this as: *BMJ* 2020;369:m2392

<http://dx.doi.org/10.1136/bmj.m2392>

Published: 16 June 2020

Routine vaccination during covid-19 pandemic response

Falls in uptake must be reversed quickly

Sonia Saxena,¹ Helen Skirrow,¹ Helen Bedford²

Routine vaccination of pregnant women and children must remain a priority during the covid-19 pandemic response. The UK and other countries could experience outbreaks of measles and other vaccine preventable diseases if uptake of routine vaccinations falls, undoing decades of progress. Vaccination has been one of the most effective interventions in driving down infant mortality to historically low levels worldwide. Complete, timely vaccination protects and confers benefits across the whole of childhood. However, this protection is greatly reduced if vaccine schedules are delayed or incomplete.¹

In the UK, almost 95% of infants have completed their primary vaccinations by their first birthday,² but uptake of later childhood vaccines and seasonal flu and whooping cough vaccines recommended in pregnancy, varies from year to year.^{3,4} Uptake is lower in inner cities, in deprived postcodes, and among some minority ethnic groups.³ Disease epidemics can disrupt healthcare and cause resurgence of vaccine preventable infections.⁵ In 2019, the UK lost its measles “elimination status” because of outbreaks resulting from historical low vaccine uptake.⁶ Children risk becoming unseen victims of the covid-19 pandemic.

Flawed messaging

Health professionals are concerned about the effect on children’s health of the upheaval of NHS routine care.⁷ The number of MMR (measles, mumps, and rubella) vaccines delivered in England dropped by 20% during the first three weeks of the lockdown,⁸ and smaller falls were reported in infant vaccines in Scotland.⁹ Although vaccination services are on the priority list for primary care, this message has been lost among the dominant message that everyone should stay at home and avoid burdening the NHS throughout April 2020.

Children’s visits to emergency departments fell by over 90% during April 2020.¹⁰ Parents have expressed concerns about overburdening the NHS and fear of exposure to covid-19 when attending for vaccination. Of 752 health visitors surveyed by the Institute of Health Visiting in May 2020, over 60% reported contact with families who had considered cancelling or postponing their child’s vaccinations (Institute of Health Visiting, personal communication).

The public is now being urged to “Stay alert, protect the NHS, save lives,” which provides little assurance for families that the NHS is open for routine care. Messages are tailored to those with an excellent command of English and may not reach families such as recent migrants and those with poor health or language literacy or little access to digital technology. Over 75% of pregnant women admitted to hospital

in the UK with covid-19 were black or Asian,¹² and the risk of dying from covid-19 during hospital admission is twofold to fourfold higher among black, Asian, and minority ethnic groups than among white people.¹² Hence, pre-existing inequalities in uptake may widen if parents from minority ethnic groups feel more vulnerable and avoid healthcare settings. Families concerned about vaccine safety will be susceptible to strongly voiced opinion and misinformation in the media.¹³ Many new parents will also lack support and information usually available from antenatal and postnatal support groups.

Time for action

General practitioners, practice nurses, midwives, and health visitors are well positioned to counsel prospective and new parents about vaccination. In areas of low uptake, contacting parents and pregnant women directly may be valuable, as conversations with a trusted healthcare professional can influence vaccination decisions.¹⁴ Vaccine uptake in primary care is highest when parents feel safe and are given timely information that vaccines are due, and when healthcare staff have systems to ensure robust call and recall.¹⁵ Primary care has risen to the challenges posed by covid-19 by enforcing triage to ensure safe distancing and by introducing hygiene measures and protective equipment to reduce infection risk. Some general practices have adopted drive-in vaccine clinics to reduce face-to-face contact with parents.¹⁶

Urgent action is required to maintain vaccination rates and limit preventable infections. We need clearer government messaging that reaches all groups and more support in the community to inform and reassure pregnant women and new parents about the importance and safety of attending primary care for vaccines during the pandemic. This needs to be accompanied by investment to support primary care teams to protect themselves and patients and to access, store, and deliver vaccines during disruptions to services and supplies. Primary care teams also need to follow up missed vaccinations quickly by strengthening reminder systems and providing opportunistic vaccination. Finally, the government must acknowledge that its policies have been blind to the serious health disparities uncovered by covid-19.¹⁷ It must invest in stronger public health surveillance to monitor areas and populations in real time to enable rapid local public health response to any drop in vaccine uptake, particularly among vulnerable groups.

The current pandemic is a reminder of the ever present threat of infectious disease. We should not miss this opportunity to strengthen public trust in primary care and public health. We hope that a

covid-19 vaccine will enable us to end containment measures safely, if such a vaccine can be developed. In the meantime, the health of our children and the whole population depends on high uptake of routine vaccinations.

Competing interests: *The BMJ* has judged that there are no disqualifying financial ties to commercial companies. The authors declare the following other interests: SS is supported by the NIHR School for Public Health Research Programme and NW London Applied Research Collaboration. SS holds funding from the Daily Mile Foundation. HS is supported by the IMPRINT network funded by UK's Global Challenges Research Fund (GCRF). HB is a member of the health improvement committee of Royal College of Paediatrics and Child Health.

Provenance and peer review: Commissioned; externally peer reviewed.

We thank two new parents for their helpful advice.

- 1 Cecil E, Bottle A, Ma R, et al. Impact of preventive primary care on children's unplanned hospital admissions: a population-based birth cohort study of UK children 2000-2013. *BMC Med* 2018;16:151. doi: 10.1186/s12916-018-1142-3 pmid: 30220255
- 2 Public Health England. Quarterly vaccination coverage statistics for children aged up to five years in the UK (COVER programme): October to December 2019. Health protection report Vol 14 No 7, 27 Mar 2020. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/875855/hpr0720_AA_COVER.pdf.
- 3 Byrne L, Ward C, White JM, Amirthalingam G, Edelstein M. Predictors of coverage of the national maternal pertussis and infant rotavirus vaccination programmes in England. *Epidemiol Infect* 2018;146:197-206. doi: 10.1017/S0950268817002497 pmid: 29239285
- 4 Walton S, Cortina-Borja M, Dezateux C, et al. Measuring the timeliness of childhood vaccinations: using cohort data and routine health records to evaluate quality of immunisation services. *Vaccine* 2017;35:7166-73. doi: 10.1016/j.vaccine.2017.10.085 pmid: 29132992
- 5 Suk JE, Paez Jimenez A, Kourouma M, et al. Post-Ebola measles outbreak in Lola, Guinea, January-June 2015. *Emerg Infect Dis* 2016;22:1106-8. doi: 10.3201/eid2206.151652 pmid: 27191621
- 6 Public Health England. Measles in England. Public health matters. 2019. <https://publichealthmatters.blog.gov.uk/2019/08/19/measles-in-england/>.
- 7 Pollard AVR, Saxena S, Leach J, et al. Protecting children [letter to editor]. *Times* 2020 Apr 14. <https://www.thetimes.co.uk/article/times-letters-social-care-and-the-treatment-of-older-people-bcm7189n6>.
- 8 McDonald HI, Tessier E, White JM, et al. Early impact of the coronavirus disease (COVID-19) pandemic and physical distancing measures on routine childhood vaccinations in England, January to April 2020. *Euro Surveill* 2020;25. doi: 10.2807/1560-7917.ES.2020.25.19.2000848 pmid: 32431288
- 9 Public Health Scotland. Uptake of first dose of 6-in-1 vaccine (offered to children at 8 weeks of age): Scotland. Scottish immunisation and recall system. 2020. <https://scotland.shinyapps.io/phs-covid-wider-impact/>.
- 10 Public Health England. Emergency department syndromic surveillance system: England, week 16. 2020. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/880759/EDSSSBulletin2020wk16.pdf.
- 11 Knight M, Bunch K, Vousden N, et al. UK Obstetric Surveillance System SARS-CoV-2 Infection in Pregnancy Collaborative Group. Characteristics and outcomes of pregnant women admitted to hospital with confirmed SARS-CoV-2 infection in UK: national population based cohort study. *BMJ* 2020;369:m2107. doi: 10.1136/bmj.m2107 pmid: 32513659
- 12 Public Health England. Disparities in the risk and outcomes of covid-19. 2020. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/891116/disparities_review.pdf.
- 13 Depoux A, Martin S, Karafillakis E, Preet R, Wilder-Smith A, Larson H. The pandemic of social media panic travels faster than the COVID-19 outbreak. *J Travel Med* 2020;27. doi: 10.1093/jtm/taaa031 pmid: 32125413
- 14 Leask J, Kinnersley P, Jackson C, Cheater F, Bedford H, Rowles G. Communicating with parents about vaccination: a framework for health professionals. *BMC Pediatr* 2012;12:154. doi: 10.1186/1471-2431-12-154 pmid: 22998654
- 15 Williams N, Woodward H, Majeed A, Saxena S. Primary care strategies to improve childhood immunisation uptake in developed countries: systematic review. *JRSM Short Rep* 2011;2:81. doi: 10.1258/shorts.2011.011112 pmid: 22046500
- 16 Hussain D. Childhood immunisations: how a practice reduced face-to-face time. *Nursing in Practice* 2020 Apr 30. <https://www.nursinginpractice.com/covid-19/childhood-immunisations-how-practice-reduced-face-face-time>.
- 17 Salway S, Holman D, Lee C, et al. Transforming the health system for the UK's multiethnic population. *BMJ* 2020;368:m268. doi: 10.1136/bmj.m268 pmid: 32047065

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.