Portsmouth Hospitals NHS Trust

drparthakar@gmail.com Follow Partha

on Twitter: @parthakar

Cite this as: *BMJ* 2020;370:m3034

<http://dx.doi.org/10.1136/bmj.m3034>

Published: 06 August 2020

THE BOTTOM LINE

Partha Kar: Our approach to tackling obesity needs rethinking

Partha Kar *consultant in diabetes and endocrinology*

Obesity is a focus of political and media discussion, since multiple datasets have shown that being overweight puts people at greater risk of serious illness from covid-19. But, if we want to improve the way we manage obesity, we need better research on what works in weight management—and a rethink of our approach to service provision.

NICE's most recent guidance on identifying, assessing, and managing obesity was published in 2014¹: in the six years since, has there been no new evidence that would warrant an update? At that time, a lot of NICE's recommendations were based on thin evidence, and NICE clearly accepted in its guidance that there were gaps in the research that needed covering. So, why hasn't the research community or the National Institute for Health Research done the relevant investigations? The available evidence on diets from randomised controlled trials is equivocal at best and confusing at worst, owing to inherent bias, non-randomisation, and small participant numbers—especially when you try to look at long term impact.

I firmly believe that the evidence base for the “tiers” used in obesity management, and in the NICE guidance, needs reviewing. For the uninitiated, there are four tiers: the first two (which cover health promotion services, primary care, and lifestyle interventions) fall under the ambit of government, local authority, and public health; the next two (specialist weight management services and bariatric surgery) are the responsibility of local clinical commissioning groups.

Tier 3 (specialist weight management services) deserves particular attention for review, especially around its evidence base. One meta-analysis suggests that specialist weight management services provide benefit, although with caveats around data from single centre studies and variability in centre staffing.²

There's also a question as to whether participants would have benefited from bariatric surgery earlier or whether innovations in medicine may have now opened up new treatment options, and whether such a review needs to be done by a specialist physician or by any trained prescriber.

If we want to be serious about obesity management, we need more focus on the societal changes needed to tackle it, rather than a simplistic narrative of lifestyle change and motivation, with the subtle (or sometimes overt) stigma attached to terminology and the implication of “not trying hard enough.” We should collect data on a national basis, do the necessary research, overhaul the archaic tier system, and try to have an open mind about all diets. The

roles of NICE and Public Health England should be to evaluate all available data around dietary interventions, whether they be low calorie or low carb, and assess the impact not only on weight loss but on sustainability and long term impact—as the data suggest that this is a major issue with dietary interventions.³

Two tiers

A fresh approach should involve two tiers only:

- One focused on dietary changes (post review of existing data with emphasis on weight loss sustainability), exercise, and psychological support;
- The other, an amalgamation of medicine and surgery with psychological support where appropriate, with a clear remit for the government to tackle the societal adaptations needed for wider uptake.

The key would be to offer the individual the choice, rather than shoehorning everyone into an imaginary magic bullet. Speaking about the perfect diet as a clinician—which in most cases is from a position of privilege—may sound good, but it achieves little on a population basis, as datasets continue to show the impact of deprivation on obesity.

We're a few months away from the 50th anniversary of Julian Tudor Hart's inverse care law paper.⁴ It wouldn't be amiss to reopen that paper and apply the principles of the inverse care law to restructuring obesity services.

Competing interests: www.bmj.com/about-bmj/freelance-contributors.

Provenance and peer review: Commissioned; not externally peer reviewed.

- 1 National Institute for Health and Care Excellence. Obesity: identification, assessment and management. Clinical guideline [CG189]. 27 Nov 2014.
- 2 Alkharaji M, Anyanwagu U, Donnelly R, Idris I. Tier 3 specialist weight management service and pre-bariatric multicomponent weight management programmes for adults with obesity living in the UK: A systematic review. *Endocrinol Diabetes Metab* 2018;2:e00042. <https://onlinelibrary.wiley.com/doi/full/10.1002/edm2.42>.
- 3 Hall KD, Kahan S. Maintenance of lost weight and long-term management of obesity. *Med Clin North Am* 2018;102:183-97. doi: 10.1016/j.mcna.2017.08.012. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5764193/>.
- 4 Tudor Hart J. The inverse care law. *Lancet* 1971;1:405-12. doi: 10.1016/S0140-6736(71)92410-X pmid: 4100731