Department of Metabolism, Digestion and Reproduction, Imperial College London, Chelsea and Westminster Hospital Campus, London, UK

v.male@imperial.ac.uk Cite this as: *BMJ* \${year};374:n2211 http://dx.doi.org/10.1136/bmj.n2211 Published:

## Menstrual changes after covid-19 vaccination

## A link is plausible and should be investigated

Victoria Male *lecturer* in reproductive immunology

Common side effects of covid-19 vaccination listed by the UK's Medicines and Healthcare Products Regulatory Agency (MHRA) include a sore arm, fever, fatigue, and myalgia.¹ Changes to periods and unexpected vaginal bleeding are not listed, but primary care clinicians and those working in reproductive health are increasingly approached by people who have experienced these events shortly after vaccination. More than 30 000 reports of these events had been made to MHRA's yellow card surveillance scheme for adverse drug reactions by 2 September 2021, across all covid-19 vaccines currently offered.¹

Most people who report a change to their period after vaccination find that it returns to normal the following cycle and, importantly, there is no evidence that covid-19 vaccination adversely affects fertility. In clinical trials, unintended pregnancies occurred at similar rates in vaccinated and unvaccinated groups. In assisted reproduction clinics, fertility measures and pregnancy rates are similar in vaccinated and unvaccinated patients. 3-6

MHRA states that evaluation of yellow card reports does not support a link between changes to menstrual periods and covid-19 vaccines since the number of reports is low relative to both the number of people vaccinated and the prevalence of menstrual disorders generally. However, the way in which yellow card data are collected makes firm conclusions difficult. Approaches better equipped to compare rates of menstrual variation in vaccinated versus unvaccinated populations are needed, and the US National Institutes of Health has made \$1.67m (£1.2m; €1.4m) available to encourage this important research. 8

Menstrual changes have been reported after both mRNA and adenovirus vectored covid-19 vaccines, suggesting that, if there is a connection, it is likely to be a result of the immune response to vaccination rather than a specific vaccine component. Vaccination against human papillomavirus (HPV) has also been associated with menstrual changes. Indeed, the menstrual cycle can be affected by immune activation in response to various stimuli, including viral infection: in one study of menstruating women, around a quarter of those infected with SARS-CoV-2 experienced menstrual disruption. 10

Biologically plausible mechanisms linking immune stimulation with menstrual changes include immunological influences on the hormones driving the menstrual cycle<sup>11</sup> or effects mediated by immune cells in the lining of the uterus, which are involved in the cyclical build-up and breakdown of this tissue.<sup>12</sup> Research exploring a possible association

between covid-19 vaccines and menstrual changes may also help understand the mechanism.

Although reported changes to the menstrual cycle after vaccination are short lived, robust research into this possible adverse reaction remains critical to the overall success of the vaccination programme. Vaccine hesitancy among young women is largely driven by false claims that covid-19 vaccines could harm their chances of future pregnancy. Failing to thoroughly investigate reports of menstrual changes after vaccination is likely to fuel these fears. If a link between vaccination and menstrual changes is confirmed, this information will allow people to plan for potentially altered cycles. Clear and trusted information is particularly important for those who rely on being able to predict their menstrual cycles to either achieve or avoid pregnancy.

We are still awaiting definitive evidence, but in the interim how should clinicians counsel those who have experienced these effects? Initially, they should be encouraged to report any changes to periods or unexpected vaginal bleeding to the MHRA's yellow card scheme. This will provide more complete data to facilitate research into any link and signal to patients that their concerns about vaccine safety are taken seriously, building trust. In terms of management, the Royal College of Obstetricians and Gynaecologists and the MHRA recommend that anyone reporting a change in periods persisting over several cycles, or new vaginal bleeding after the menopause, should be managed according to the usual clinical guidelines for these conditions.<sup>7</sup>

One important lesson is that the effects of medical interventions on menstruation should not be an afterthought in future research. Clinical trials provide the ideal setting in which to differentiate between menstrual changes caused by interventions from those that occur anyway, but participants are unlikely to report changes to periods unless specifically asked. Information about menstrual cycles and other vaginal bleeding should be actively solicited in future clinical trials, including trials of covid-19 vaccines.

Competing interests: The BMJ has judged that there are no disqualifying financial ties to commercial companies. The author declares the following other interests: research funding from the Wellcome Trust and research charity Borne; payments to act as an external examiner for the University of Cambridge and the University of Leeds; and royalties received for my contribution to Immunology 9th edition (Elsevier). Further details of The BMJ policy on financial interests is here: https://www.bmj.com/sites/default/files/attachments/resources/2016/03/16-current-bmj-education-col-form.pdf.

Provenance and peer review: Commissioned; not externally peer reviewed.

Medicines and Healthcare Products Regulatory Agency. Coronavirus vaccine—weekly summary of yellow card reporting. 2021. https://www.gov.uk/government/publications/coronavirus-covid-19-vaccine-adverse-reactions/coronavirus-vaccine-summary-of-yellow-card-reporting#annex-1-vaccine-analysis-print

- Male V. Are covid-19 vaccines safe in pregnancy? Nat Rev Immunol 2021;21:200-1. doi: 10.1038/s41577-021-00525-y pmid: 33658707
- 3 Morris RS. SARS-CoV-2 spike protein seropositivity from vaccination or infection does not cause sterility. F S Rep 2021. doi: 10.1016/j.xfre.2021.05.010. pmid: 34095871
- Orvieto R, Noach-Hirsh M, Segev-Zahav A, Haas J, Nahum R, Aizer A. Does mRNA SARS-CoV-2 vaccine influence patients' performance during IVF-ET cycle? *Reprod Biol Endocrinol* 2021;19:69. doi: 10.1186/s12958-021-00757-6 pmid: 33985514
- 5 Bentov Y, Beharier O, Moav-Zafrir A, et al. Ovarian follicular function is not altered by SARS-Cov-2 infection or BNT162b2 mRNA Covid-19 vaccination. *medRxiv* 2021:2021.04.09.21255195. [Preprint.] doi: 10.1101/2021.04.09.21255195
- 6 Safrai M, Rottenstreich A, Herzberg S, Imbar T, Reubinoff B, Ben-Meir A. Stopping the misinformation: BNT162b2 COVID-19 vaccine has no negative effect on women's fertility. medRxiv 2021;2021.05.30.21258079 [Preprint.] doi: 10.1101/2021.05.30.21258079
- Medicines and Healthcare Products Regulatory Agency. COVID-19 vaccines: updates for August 2021. https://www.gov.uk/drug-safety-update/covid-19-vaccines-updates-for-august-2021
- 8 Eunice Kennedy Shriver National Institute of Child Health and Human Development. Item of interest: NIH funds studies to assess potential effects of COVID-19 vaccination on menstruation. 2021.https://www.nichd.nih.gov/newsroom/news/083021-COVID-19-vaccination-menstruation
- 9 Suzuki S, Hosono A. No association between HPV vaccine and reported post-vaccination symptoms in Japanese young women: Results of the Nagoya study. *Papillomavirus Res* 2018;5:96-103. doi: 10.1016/j.pvr.2018.02.002 pmid: 29481964
- Li K, Chen G, Hou H, etal. Analysis of sex hormones and menstruation in COVID-19 women of child-bearing age. *Reprod Biomed Online* 2021;42:260-7. doi: 10.1016/j.rbmo.2020.09.020 pmid: 33288478
- 11 Karagiannis A, Harsoulis F. Gonadal dysfunction in systemic diseases. Eur J Endocrinol 2005;152:501-13. doi: 10.1530/eje.1.01886 pmid: 15817904
- Monin L, Whettlock EM, Male V. Immune responses in the human female reproductive tract. Immunology 2020;160:106-15. doi: 10.1111/imm.13136 pmid: 31630394
- Speed B. Young women are the unlikely new face of covid-19 vaccine resistance. i News 2021 Jan 6. https://inews.co.uk/news/health/coronavirus-latest-experts-debunk-vaccine-fertility-myths-women-819783
- 14 Royal College of Obstetricians and Gynaecologists. RCOG responds to reports that COVID-19 vaccine affects periods. 2021. https://www.rcog.org.uk/en/news/rcog-responds-to-reports-that-covid-19-vaccine-affects-periods/

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.