



NHS Lanarkshire and Edinburgh Napier
University, Scotland, UK

stephanie.dancer@lanarkshire.scot.nhs.uk

Cite this as: *BMJ* 2022;377:e01408

<http://dx.doi.org/10.1136/bmj.01408>

Published: 29 June 2022

Airborne SARS-CoV-2

Time for an indoor air revolution

Stephanie J Dancer *consultant microbiologist*

Debate over the exact mode of transmission of SARS-CoV-2 has been intense.¹ This is entirely reasonable, given that the mechanism of spread determines preventive and potentially lifesaving policies. But the choice between respiratory aerosol or droplet settled on short range droplets, which neatly circumvented any risk outside the fabled 2m zone.¹ This choice gave rise to social distancing, hand and surface hygiene, and masks, but not to improved indoor air quality.

And so the debate smoulders on, as Duval and colleagues (doi:10.1136/bmj-2021-068743) report from their linked systematic review supporting the role of long distance airborne transmission of SARS-CoV-2.² The review examined covid-19 transmission events in a variety of indoor community settings ranging from fitness facilities, offices, buses, and restaurants to choir venues and a church, but not hospitals, hospices, or care homes.²⁻⁸ The inclusion of care home outbreaks might have strengthened overall findings, along with more recent studies detailing nosocomial clusters among vaccinated healthcare workers.⁹⁻¹⁰

Study selection was, of necessity, somewhat labile, because any outbreak inferring even the slightest possibility of contact or fomite transmission would have been excluded. This might explain the omission of notable community outbreaks,¹¹⁻¹³ including those where the virus almost certainly spread through sanitation systems in high rise flats.¹⁴⁻¹⁵ This opportunistic transmission route is reminiscent of the notorious Amoy Gardens outbreak of SARS-1 in Hong Kong.¹⁶

SARS-CoV-2 survives in faeces, urine, and waste water, and aerosol transmission through interconnected sanitary drains (just as for SARS-1) needs further exploration.¹⁷ Viral spread in toilet facilities might not feature in the literature, but that does not mean the risk should be ignored. Faecal aerosols might have played an important role in transmission during the covid-19 pandemic, especially as diarrhoea is common among infected patients and viral shedding persists in stool despite negative respiratory sample results.¹⁷⁻¹⁸

Arguably, Duval and colleagues' review should also have mentioned studies reporting aerosol transmission of SARS-CoV-2 between animals.¹⁹⁻²¹ Given that similar studies on humans would never obtain ethical approval, these investigations—which virtually all support long distance aerosol spread, skilfully emulate the original work on tubercle transmission from the early 1960s.²²⁻²³ This work was eventually accepted by the scientific community as evidence for airborne transmission of tuberculosis in humans—despite the fact that *Mycobacterium*

tuberculosis has never been successfully cultured from air. It is hoped that SARS-CoV-2 and its proclivity for airborne transmission will be accepted a little quicker than it was for tuberculosis. Influenza might have to wait.²²

Of course, some argue that reliance on observational events is poor science. But a role clearly exists for detailed epidemiology in respiratory outbreaks, simply because it provides empirical validation that aerosol transmission occurs, and in fact occurs extensively.²² As Duval and colleagues surmise, there is a need to develop a new framework for evidence synthesis of outbreak investigations.²⁻²⁴⁻²⁵ Either that, or more than a century of detailed epidemiological work identifying the cause of disease outbreaks and tracking the spread of notable pathogens must be ignored. After all, who would choose to inhabit the “control” environment in a randomised trial examining the protective effect of fresh air during an influenza outbreak?²⁶ It is laudable to seek solid scientific evidence, but when a disease spreads so rapidly, we really should not have to wait for randomised evidence that might never materialise.²⁷

Just as the world woke up to a pandemic, a small group of determined scientists (including this author) appealed for consideration of airborne spread.²⁸ Their advice was summarily dismissed.¹ And so the group—in common with the pioneers of tuberculosis transmission—“provided an ingredient that scientists seldom mention: a mission to convince unbelievers.”²⁹

Now, the evidence presented in Duval and colleagues' review, tenuous as it is, validates the premise that tiny respiratory particles containing SARS-CoV-2 freely transmit throughout inadequately ventilated environments. That this small group of scientists have (almost) won their argument is of small consolation to those still experiencing the effects of covid-19. But through persistence and escalating independent evidence, better indoor air quality can be entertained for everyone in the future.³⁰ It is hoped that public health leaders will develop practical guidance and “tilt” people and places closer to safety.³¹ Now, indeed, is the time for an indoor air revolution.³²⁻³³

Competing interests: I have read and understood the BMJ Group policy on declaration of interests and declare the following interests: none.

Provenance and peer review: Commissioned; not externally peer reviewed.

- Jimenez JL, Marr LC, Randall K, et al. What were the historical reasons for the resistance to recognizing airborne transmission during the COVID-19 pandemic? *Indoor Air* 2022 (in press).
- Duval D, Palmer JC, Tudge I, et al. Long distance airborne transmission of SARS-CoV-2: rapid systematic review. *BMJ* 2022;377:e068743.
- Lednicky JA, Lauzard M, Fan ZH, et al. Viable SARS-CoV-2 in the air of a hospital room with COVID-19 patients. *Int J Infect Dis* 2020;100:476-82. doi: 10.1016/j.ijid.2020.09.025 pmid: 32949774

- 4 Cheng VC, Fung KS, Siu GK, et al. Nosocomial outbreak of Coronavirus Disease 2019 by possible airborne transmission leading to a superspreading event. *Clin Infect Dis* 2021;73:e1356-64. doi: 10.1093/cid/ciab313 pmid: 33851214
- 5 Nissen K, Krambrich J, Akaberi D, et al. Long-distance airborne dispersal of SARS-CoV-2 in COVID-19 wards. *Sci Rep* 2020;10:19589. doi: 10.1038/s41598-020-76442-2 pmid: 33177563
- 6 Goldberg L, Levinsky Y, Marcus N, et al. SARS-CoV-2 infection among health care workers despite the use of surgical masks and physical distancing—the role of airborne transmission. *Open Forum Infect Dis* 2021;8:ofab036. doi: 10.1093/ofid/ofab036 pmid: 33732749
- 7 Feathers L, Hinde T, Bale T, et al. Outbreak of SARS-CoV-2 at a hospice: terminated after the implementation of enhanced aerosol infection control measures. *Interface Focus* 2022;12:20210066. doi: 10.1098/rsfs.2021.0066 pmid: 35261730
- 8 de Man P, Paltansing S, Ong DSY, Vaessen N, van Nielen G, Koeleman JGM. Outbreak of Coronavirus Disease 2019 (COVID-19) in a nursing home associated with aerosol transmission as a result of inadequate ventilation. *Clin Infect Dis* 2021;73:170-1. doi: 10.1093/cid/ciaa1270 pmid: 32857130
- 9 Park SY, Kim TH, Lee E, et al. A SARS-CoV-2 outbreak associated with vaccine breakthrough in an acute care hospital. *Am J Infect Control* 2022;50:196-6553(22)00436-9. doi: 10.1016/j.ajic.2022.05.010 pmid: 35605754
- 10 Lim WY, Tan GSE, Htun HL, et al. First nosocomial cluster of COVID-19 due to the Delta variant in a major acute care hospital in Singapore: investigations and outbreak response. *J Hosp Infect* 2022;122:27-34. doi: 10.1016/j.jhin.2021.12.011 pmid: 34942201
- 11 Wallace M, Hagan L, Curran KG, et al. COVID-19 in correctional and detention facilities—United States, February–April 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:587-90. doi: 10.15585/mmwr.mm6919e1 pmid: 32407300
- 12 Pauser J, Schwarz C, Morgan J, Jantsch J, Brem M. SARS-CoV-2 transmission during an indoor professional sporting event. *Sci Rep* 2021;11:20723. doi: 10.1038/s41598-021-99997-0 pmid: 34671096
- 13 Park SY, Kim YM, Yi S, et al. Coronavirus Disease Outbreak in Call Center, South Korea. *Emerg Infect Dis* 2020;26:1666-70. doi: 10.3201/eid2608.201274 pmid: 32324530
- 14 Kang M, Wei J, Yuan J, et al. Probable evidence of fecal aerosol transmission of SARS-CoV-2 in a high-rise building. *Ann Intern Med* 2020;173:974-80. doi: 10.7326/M20-0928 pmid: 32870707
- 15 Wang Q, Li Y, Lung DC, et al. Aerosol transmission of SARS-CoV-2 due to the chimney effect in two high-rise housing drainage stacks. *J Hazard Mater* 2022;421:126799. doi: 10.1016/j.jhazmat.2021.126799 pmid: 34396958
- 16 Yu IT, Li Y, Wong TW, et al. Evidence of airborne transmission of the severe acute respiratory syndrome virus. *N Engl J Med* 2004;350:1731-9. doi: 10.1056/NEJMoa032867 pmid: 15102999
- 17 Dancer SJ, Li Y, Hart A, Tang JW, Jones DL. What is the risk of acquiring SARS-CoV-2 from the use of public toilets? *Sci Total Environ* 2021;792:148341. doi: 10.1016/j.scitotenv.2021.148341 pmid: 34146809
- 18 Chen Y, Chen L, Deng Q, et al. The presence of SARS-CoV-2 RNA in the feces of COVID-19 patients. *J Med Virol* 2020;92:833-40. doi: 10.1002/jmv.25825 pmid: 32243607
- 19 Kutter JS, de Meulder D, Bestebroer TM, et al. SARS-CoV and SARS-CoV-2 are transmitted through the air between ferrets over more than one meter distance. *Nat Commun* 2021;12:1653. doi: 10.1038/s41467-021-21918-6 pmid: 33712573
- 20 Port JR, Yinda CK, Avanzato VA, et al. 2021. Increased aerosol transmission for B.1.1.7 (alpha variant) over lineage A variant of SARS-CoV-2. *bioRxiv* (doi: 10.1101/2021.07.26.453518).
- 21 Hawks SA, Prussin AJ, 2nd Kuchinsky SC, Pan J, Marr LC, Duggal NK. Infectious SARS-CoV-2 is emitted in aerosol particles. *mBio* 2021;12:e0252721. doi: 10.1128/mBio.02527-21 pmid: 34663099
- 22 Tellier R. COVID-19: the case for aerosol transmission. *Interface Focus* 2022;12:20210072. doi: 10.1098/rsfs.2021.0072 pmid: 35261731
- 23 Riley RL, Mills CC, O'Grady F, Sultan LU, Wittstadt F, Shivpuri DN. Infectiousness of air from a tuberculosis ward. Ultraviolet irradiation of infected air: comparative infectiousness of different patients. *Am Rev Respir Dis* 1962;85:511-25. pmid: 14492300
- 24 Dancer SJ, Inkster T. One size does not fit all: why infection prevention is difficult to randomize or control. *J Hosp Infect* 2022;123:182-3. doi: 10.1016/j.jhin.2022.02.008 pmid: 35196558
- 25 Stone SP, Cooper BS, Kibbler CC, et al. The ORION statement: guidelines for transparent reporting of outbreak reports and intervention studies of nosocomial infection. *Lancet Infect Dis* 2007;7:282-8. doi: 10.1016/S1473-3099(07)70082-8 pmid: 17376385
- 26 Hobday RA, Cason JW. The open-air treatment of pandemic influenza. *Am J Public Health* 2009;99(Suppl 2):S236-42. doi: 10.2105/AJPH.2008.134627. pmid: 19461112
- 27 Lewis D. Why the WHO took two years to say COVID is airborne. *Nature* April 6, 2022. <https://www.nature.com/articles/d41586-022-00925-7>
- 28 Morawska L, Milton DK. It Is Time to Address Airborne Transmission of Coronavirus Disease 2019 (COVID-19). *Clin Infect Dis* 2020;71:2311-3. doi: 10.1093/cid/ciaa939 pmid: 32628269
- 29 Riley RL. What nobody needs to know about airborne infection. *Am J Respir Crit Care Med* 2001;163:7-8. doi: 10.1164/ajrccm.163.1.hh11-00 pmid: 11208616
- 30 Morawska L, Tang JW, Bahnfleth W, et al. How can airborne transmission of COVID-19 indoors be minimised? *Environ Int* 2020;142:105832. doi: 10.1016/j.envint.2020.105832 pmid: 32521345
- 31 Lynch RM, Goring R. Practical steps to improve air flow in long-term care resident rooms to reduce COVID-19 infection risk. *J Am Med Dir Assoc* 2020;21:893-4. doi: 10.1016/j.jamda.2020.04.001 pmid: 32389592
- 32 Morawska L, Allen J, Bahnfleth W, et al. A paradigm shift to combat indoor respiratory infection. *Science* 2021;372:689-91. doi: 10.1126/science.abg2025 pmid: 33986171
- 33 Dancer SJ, Bluysen PM, Li Y, Tang JW. Why don't we just open the windows? *BMJ* 2021;375:n2895. doi: 10.1136/bmj.n2895 pmid: 34836876

This article is made freely available for personal use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.