nected, formed a perfectly solid mass; whereas the centre of the loops, towards the circumference of the tumour, being but loosely connected by thin cellular tissue, had, in many places, left inter-laminar spaces, which had subsequently become filled with fluid. The cysts thus formed were of various sizes, and few in number; their cavities were perfectly smooth, and not subdivided by bands of cellular tissue.

There is another class of Fibrous Tumours, which appear to have been formed by the union of several smaller ones by means of loose cellular tissue, in the meshes of which, fluid has been thrown out, and has subsequently formed cavities or cysts. In these cases, the cavities, generally speaking, are not of any great size, and the mass looks like a large œdematous Fibrous Tumour. When punctured, but little fluid escapes from them; but, if incisions be made in several places, the greater part of the fluid will drain off in a few hours, leaving a coarse cavernous-looking tissue.

The Fibro-Cystic Tumour described in the case of M. G., cannot, I think, be looked upon as belonging to either of the three varieties already mentioned. From its earliest formation, the tumour was most probably of a fibro-cystic nature, and exactly resembled that form of disease so commonly observed in the ovary, which is composed of unilocular cysts, of various sizes, closely aggregated together, with more or less delicate fibro-serous walls; in fact, so close was the resemblance, that for some time the mass was taken for a diseased ovary. Portions of it also closely resembled the simple cystic disease of the testis, as well as one form of fibro-cystic disease which I have met with in a large tumour of the breast. The accumulations of solid fibrous tissue observed in the case of M. G., were scattered, and small in comparison to the size of the tumour; whereas the mass of it was made up of cysts, which existed in every part, even close to the root.

ON THE CAUSES, CONSEQUENCES, AND TREATMENT OF INFLAMMATION OF THE VEINS:

WITH CASES ILLUSTRATING THE EFFECTS OF PURULENT INFECTION OF
THE BLOOD.

By HENRY LEE, Esq., F.R.C.S. Eng., Assistant-Surgeon to King's College
Hospital, etc.

PART II.

(Concluded from page 240.)

VIII. The experiments cited in the first part of this essay, illustrate the power possessed by the blood of preventing certain foreign substances from circulating with it. They shew that pus, in particular, has a tendency to coagulate the blood; and that by this means, when introduced into the vessels, its progress is arrested in some part of the circulating system. This fact, which, taken by itself, might appear of little consequence, assumes considerable importance when considered as one of the inherent properties of the blood, at all times ready, under favourable circumstances, to be called into action in the living body.

The conditions under which pus will determine the coagulation of the blood, and those under which it will circulate in the living vessels, require to be accurately ascertained, in order to reconcile the discordant evidence which we at present have upon this point.

Dr. Sédillot,¹ in a work recently published, mentions, that a great number of cases are met with, in which pus is poured into the general circulation without meeting with any obstruction, and states that, in such instances, he can detect the globules of pus in different parts of the circulating system. He even affirms that he can recognise a disease caused by purulent infection, by examining, under a microscope, a portion of the blood abstracted from the body.

M. Dance, and, since his time, equally accurate observers, have, on the other hand, failed to detect the characters of pus in the blood, even when that fluid had been injected into the veins of living animals. The results of these different observations may perhaps be reconciled, by considering the influence exercised upon the globules of pus by the blood, before its coagulating power has been impaired. This subject appears not to have hitherto occupied the attention of pathologists.

In all the cases quoted by Dr. Sédillot, in which he detected the globules of pus in the blood, the patients had died of the disease; but in the researches instituted by M. Dance and others, the experiments were made upon animals in perfect health. In the latter, the pus cannot enter the circulation, as has already been shewn, or can only do so after the blood has partially or entirely coagulated round it, and the coagulum has subsequently become broken up.

In the act of coagulation under these circumstances, the appearances of the globules of pus are changed,—these being perhaps mechanically compressed by the contraction of the fibrine,—so that the most experienced eye can no longer recognise them.

Pus, mixed with healthy recently drawn blood, out of the body, will entirely lose its characters in this way; and as the coagulation, under similar circumstances, is by no means retarded in the living vessels, we may, without fear of contradiction, affirm, that globules of pus cannot be detected when introduced into the vessels in small quantities, and mixed with healthy blood.

In cases where, from long-continued disease and the repeated introduction of vitiated fluids into the circulation, the blood has lost its power, there appears no reason to doubt the correctness of Dr. Sédillot's observations; and it is probable that pus-globules may then circulate with those of the blood.

In experiments upon animals, it has always been found that the power of the constitution, in resisting the effects of the injection of pus into the veins, was much greater at the first than at any subsequent operation. This circumstance would appear to associate itself directly with the observations now made, and to afford another illustration of the power of healthy blood in resisting the entrance of some foreign matters into the system.

From the consideration of these facts, and of the experiments previously recorded, it becomes evident, that the introduction of pus into the

¹ De l'Infection Purulente, p. 399.

system through an injured or inflamed vein, can rarely be the first step towards purulent infection of the system. Some change must previously have passed in the blood, by which its coagulating power is impaired, or some unusual mechanical means must have been employed, before the pus can find its way in the course of the circulation. The contradictory statements which have been made by those who have injected pus into the veins, may thus be reconciled, by taking into account the power exercised by the blood in the experiments which have been made. There can be little doubt that, while, in some instances, a portion of the pus has been forced into the general circulation, in the great majority of cases it has been detained in the vein into which it was first introduced, and has never become part of the circulating fluid. We accordingly find some experimenters recording the secondary diseases which they observed, while in other hands these appearances were not produced.

Dr. Sédillot¹ has attempted to prove that the globules, or solid parts of pus, must be introduced into the system, in order to produce well-marked indications of purulent infection. But this hypothesis would not only appear to be at variance with the oft-repeated experiments of MM. Gaspard and Cruveilhier, in which similar effects were produced by the injection of mercury and of putrid fluids, but would also leave unexplained the mode of the introduction of these globules, where there is evidence of the disease originating through the lymphatic system. The changes which all substances undergo in their passage through the absorbent glands, would at once forbid the idea that globules of pus could be thus introduced unchanged into the circulation; and yet we have direct evidence (Case xxix) of irritating fluids being conveyed in this way into the system, and leading to the formation of secondary abscesses.

Another class of cases, in which there would be difficulty in admitting the doctrine of the introduction of pus in substance into the circulation, presents itself, where, in the primary affections (as in Case vi), no evidence can be obtained of the original lesion having suppurated. The fluids effused in such cases may be serum, lymph, or blood, mixed in different proportions; and yet the constitutional symptoms will be exactly similar to those which follow the formation of pus in other cases. There may exist, both in the primary and in the secondary affection, every intermediate gradation between the healthy secretion of a part, and the formation of pure pus, or pus mixed with blood or lymph, without any of the essential characters of the disease being absent. An inflamed bursa, or a punctured wound, without the formation of pus, (Cases iv and v), may give rise to symptoms as severe, and consequences as fatal, as any that arise from the direct introduction of pus into the system. The secondary affections, in such cases, may run their course

¹ In deducing general conclusions from experiments upon animals, it must be borne in mind, that in them suppuration is induced with great difficulty. Many of the appearances produced by the injection of putrid fluids (as in Experiment xiv) would, in man, probably have terminated in suppuration. Dr. Sédillot has nevertheless established the fact, that, generally speaking, a different class of post-mortem appearances may be expected from the introduction of decomposed serum, to those produced from fluids containing solid particles.

and prove as speedily fatal, as where well-formed purulent deposits have taken place. The most severe constitutional symptoms will sometimes be followed by the effusion of bloody fluid only, in one of the serous cavities (Case xxx). It would be unphilosophical, even were it practicable, to refer such cases to a different disease, merely because the accidental circumstance of the formation of pus is wanting. The origin of the affection in such instances may be as well-marked, the poison can often be traced as distinctly into the system, and the secondary disease may be as clearly connected with the primary, as in any case where pus has been originally formed. In some cases again, the constitutional symptoms which accompany, or are followed by, effusions into distant parts of the body, begin before sufficient time has elapsed to allow the supposition that pus can have been fully formed at the original seat of injury. Such instances occasionally, although rarely, present themselves in extensive burns and scalds, occurring in enfeebled habits, and after amputation of the limbs in scrofulous children.

In nearly all cases, when the origin of the constitutional disease cannot be traced to the introduction of diseased fluid into the system through an open vein, it will be found that the part primarily injured has wanted the degree of vigour, requisite to establish and maintain healthy adhesive inflammation.

In the first part of this paper,¹ I have endeavoured to show that, where lymph is effused around a poisoned wound, the virus will find its way less easily along the absorbent vessels, than when no such effusion has taken place; and that when, in such a wound, the effusion of lymph is checked or prevented, as by the administration of mercury, a larger proportion of cases will indicate an affection of the lymphatic system, than when the natural process has not been interfered with. There can be little doubt, that the same principle may be observed with regard to ordinary wounds. The number of cases in which the absorbents inflame, will be in inverse proportion to the number of those in which the original wounds are circumscribed by healthy adhesive inflammation. In Case xxvii, it is mentioned, that the surface of a muscle, implicated in the original lesion, was as cleanly dissected as if with a scalpel, thus showing the total absence of any surrounding effusion of lymph. The absence of, or defect in, the process of adhesion may thus be associated with inflammation of the absorbents, as the want of "union by first intention" has been shown to be connected with inflammation of the veins (Section ii.)

The minuteness of the absorbent vessels, and the changes which their contents undergo in their glands, prevent any unhealthy fluids from being as readily recognised in them as in the veins. But when the progress of inflammation can be traced along these vessels from a wound, towards the centre of the circulation, marked, as it often is, at intervals, by the formation of abscesses, we cannot doubt that an irritating fluid has found its way along their canals: and when the constitutional symptoms, which arise at the same time, terminate in the formation of purulent deposits (as in Case xxix), we cannot but admit that the absorbent vessels are the direct means by which, in such cases,

¹ LONDON JOURNAL OF MEDICINE, vol. i, p. 799.

diseased secretions are poured into the blood, and the system becomes infected. It would, therefore, appear that there are two principal conditions, under which local disease may produce a general infection of the system by the direct introduction of vitiated fluids into the blood. The first of these is connected with defective union in injured veins; the second is associated with want of healthy adhesion in inflamed lymphatics.

The period of invasion of the attack differs in some degree in the different classes of cases, but it is generally marked with great precision: even when apparent recovery has been followed by a second attack, the occurrence has in each instance been accurately noted by the sudden appearance of constitutional symptoms (see Case XXXVII).

When one of the large veins has been originally affected, the period which elapses before symptoms of infection of the system manifest themselves, is comparatively short (Case I). In cases occurring after child-birth, it is usually longer, extending to the end of the second week. After surgical operations or accidents involving some portion of bone, the access of the disease will be marked by a rigor during the third or fourth week; and finally, when the absorbent system is primarily affected, the period of the occurrence of the constitutional symptoms may be much farther removed from that of the original injury (if any such existed), and is by no means so accurately defined.

At the time of the occurrence of the general disturbance of the system, the local injury or wound will generally put on an unhealthy appearance. The skin in the immediate neighbourhood will sometimes assume a dull brownish-red appearance, which will gradually fade into the colour of the surrounding parts. This symptom will usually commence near the termination of those vessels, which are derived from the same trunk as those which supply the injured part. When the original injury is complicated with a wound upon the surface of the body, it will usually become dry and glazed, and the blush upon the skin will commence in its neighbourhood, or a short distance from it, and will usually extend towards the centre of the circulation, without presenting any very defined margin: occasionally it will extend, in the form of erratic erysipelas, over a large part of the body.¹

IX. The commencement of constitutional disease, after direct infection of the blood, is marked by a sudden change in the manner and appearance of the patient; a severe rigor is usually the most prominent symptom, and is followed by much febrile excitement, or by extreme depression; a very peculiar heat of skin (Case XXII) will sometimes be present, while, at other times, the surface will be covered by a profuse clammy perspiration. The rigor may be repeated at irregular intervals, but occasionally it will recur about the same hour for three or four days in succession (Case XXXVII), and in a few instances it will not be observed at all.

Great depression frequently accompanies even the first stages of this disease, indicated by a want of tone in the pulse, by an extremely listless

¹ In a case of fracture of the femur into the knee-joint, I saw a dark ill-defined erysipelatous blush extend from the affected limb to the body, and thence to the head. Purulent deposits formed in various parts of the body, of which the patient died.

manner, and sometimes by a tendency to syncope (Case 1). The countenance becomes anxious, the tongue dry and brown in the centre, and red at the edges, or, in other instances, it presents a coating of a pasty yellowish-white colour; a dusky yellow hue frequently pervades the skin, and sometimes the conjunctivæ of the eyes. This may or may not depend upon an accompanying affection of the liver. The pulse varies much in frequency in different cases, and at different times in the same case: generally it is very rapid, especially when accompanied with much heat of skin.

The pain is sometimes severe, and may be referred exactly to the spot which subsequent examination shows to have been the seat of secondary inflammation; at other times it is not confined to any particular situation, but consists of general ill-defined feelings of short duration, and recurring at irregular intervals. The peculiarity of such sensations is best expressed by the terms applied to them by the patients themselves. "Catching pains all over", "soreness of the stomach", and "thrilling in the blood", not unfrequently accompany this disease.

Vomiting may occur, either as a symptom of constitutional disturbance, or as indicative of inflammation of an abdominal organ (Case xxvi). In the latter case, it is extremely obstinate, and the fluid ejected is generally of a green colour. Diarrhœa is a symptom of frequent occurrence, and appears to exercise a considerable influence on the course of the disease. Its appearance will not unfrequently be accompanied by relief of the other symptoms (Case III); when it occurs, it is generally profuse, and little under the control of medicine, but, if checked, may be followed by a sudden change for the worse in the condition of the patient.

The intellect is seldom affected during the first stages of the complaint; but subsequently, in severe cases, restlessness, delirium, and coma, seldom fail to succeed each other. These symptoms are all peculiar, both in regard to the rapidity with which they make their appearance, and also the sudden manner in which they occasionally disappear. The disease may seem, within a few hours, to leave a part which it has first attacked, and to fall upon a different organ in some remote part of the body.

X. The post-mortem appearances observed in those who die in consequence of the introduction of vitiated fluids into the blood, cannot, for the most part, be distinguished from similar changes produced by other causes; yet there are some effects which are peculiar, and may be directly associated with the reception of foreign matter into the circulation. The most characteristic circumstance, attending the extension of disease to different organs of the body through the medium of the blood, is that several parts of these organs, or even different organs, will be simultaneously attacked. The disease will appear at once in various spots, which will become rapidly disorganized, while the surrounding textures will remain unaltered, either in structure or colour. The appearances observed upon dissection will vary according to the part attacked, and the stage of development in which the disease is found.

The lungs are the organs in which the successive changes may best be observed. When puriform fluid has entered the circulation,

the first appearance produced in the structure of the lungs, is that of one or more congested or dilated veins¹ of very small diameter. This will be followed by a well defined spot, of much darker colour than the surrounding texture (see Plate, facing p. 624.) Several of these spots will probably appear at the same time, and each one of them will soon become surrounded by a hard spherical patch of purple congestion. Effusion of lymph will now take place, commencing in the centre of each affected portion, and gradually extending towards its circumference. If the disease continue, each spot will suppurate, and the different parts will become softened and broken down, in the same order in which they were previously solidified.

The liver frequently becomes the seat of secondary inflammation. In the early stage, brownish-red spots may be observed scattered through its substance. These, as they extend, assume a bluish or slate-colour; and the structure of the liver thus affected is found to have lost its consistence, and to be very easily broken down by pressure. Every part affected here, as in the lungs, proceeds rapidly to suppuration; and the usual appearance presented after death, is that of several small circumscribed abscesses, around which the structure of the liver has been condensed only to a very small extent. It sometimes happens, that the larger veins in the liver become inflamed. These vessels, being held open by the firm structure of the part, are not so readily obliterated as in other situations; and it consequently happens, that the lymph and pus poured into them become irregularly mixed with the blood, more or less perfectly coagulated, which they contain: a very peculiar mottled appearance, resembling granite, is thus occasionally produced.

Affections of the spleen, produced by the introduction of foreign matter into the blood, are probably not so readily recognised as similar affections in the lungs and liver. For, although the spleen is often found to be diseased in those who die from infection of the blood, yet it is comparatively seldom that secondary abscesses have been found in it. In the accompanying table, containing twenty-three cases, some morbid appearance, not recognised as peculiarly the result of secondary inflammation, was observed in the spleen in no less than eight instances. So large a proportion of cases renders it probable, that the alterations observed have more than an accidental connexion with the disease of which the patient died, although they presented no characters which could be said to be peculiar to that disease. In well-marked cases of secondary affections of the spleen, one or more well defined, but frequently irregular indurations, of a chocolate colour, may be recognised; such patches are usually seen soon after the commencement of the complaint, and very soon become softened or broken down. The rapidity with which they lose their original character, may probably account for their being comparatively seldom observed in post-mortem examinations.

Deposits of lymph are sometimes met with in the kidneys; but these are of small extent, of a light colour, and resemble lymph deposited in consequence of ordinary inflammation. The patches of congestion, so characteristic of this disease in other organs, are not here observed.

¹ For the knowledge of this fact, I am indebted to Mr. Cæsar Hawkins, of St. George's Hospital.

This may depend upon the peculiar disposition of the capillary system of the kidney. The blood has to pass through the Malpighian tufts, and may be purified, or altered in character, before it reaches the proper venous system of the organ.

In cases where purulent infection of the blood has been purposely produced, portions of the kidney will not unfrequently be found inflamed and firmer than natural; but, if the origin of the disease were not known, these appearances could not be distinguished from those produced by inflammation of the kidney from other causes.

The skin is liable to be affected in three different forms. (See Cases I, II, III, IX, X, and XXXIV.) The first of these occurs very rarely, and consists of small deposits of matter in the structure, or upon the surface of the skin, resembling in many respects the pustules of small-pox. The second form is also of rare occurrence, and consists of small congested spots on the surface of the skin. These are generally of a dark purple hue, but I have seen one case in which they were of a bright red colour. In this instance, a secondary abscess had formed in the knee-joint, and some pustules appeared upon the skin in the neighbourhood. A fortnight before the death of the patient, a number of small bright red spots made their appearance in different parts of the thigh and upper part of the leg; some of these were three or four lines in diameter, while others were so small as not to be seen without attention; they appeared in accurately defined spots, of a brighter colour than the mucous membrane of the lips, and continued unchanged in appearance till death. The third form presents itself much more frequently than either of the others, although it has not hitherto much attracted the attention of pathologists, in connexion with purulent or other infection of the blood. It commences very suddenly, and frequently without any particular attention being directed to the part. A large circular patch of congestion, livid or purple in the centre, but becoming of a lighter colour towards the circumference, will form, usually upon some part of the lower extremities. The skin of the calf of the leg is perhaps more frequently attacked than that of any other part. In the centre of the congested portion, mortification very rapidly takes place, and is indicated by the part assuming a black or dull leaden colour. In some cases, it would be difficult to say where the mortification ceases, and the congestion begins; but in other instances, there is a distinct line of demarcation formed: a zone of bright red congestion will then occasionally surround the mortified part.

Some modifications of this third form of affection of the skin may be met with occasionally in the course of the disease. Blotches assuming a livid or dusky red appearance (which gradually fades into the colour of the surrounding skin), will present themselves in different parts (Case XXXVI). In some instances, the superficial portions only of the skin are destroyed, and the parts beneath appear comparatively unaffected; small circumscribed portions of the outer layer of the skin will exfoliate, and the subjacent parts will heal without suppuration, by a process similar to that of scabbing (Case III).

It is remarkable in this disease, that the most vascular parts are those which soonest lose their vitality. Thus, in the case last referred to, portions of the surface of the skin perished, while the deeper layers

recovered ; again, it is not unusual to observe the whole thickness of the skin destroyed, without any corresponding affection of the cellular membrane beneath. The quantity of blood sent to a part would thus appear to favour mortification in this disease. The reason of this peculiarity will be considered in the next section.

In two of the instances recorded in the Appendix (Cases xxii and xxvi), the lining membrane of the rectum was found of a very dark colour, and in one it had assumed a greenish appearance. This discoloration was at first looked upon as some accidental complication, or as depending upon previous disease. But M. Gaspard has noticed a similar condition, after the artificial introduction of putrid fluid into the blood.¹ In one of the experiments referred to, the mucous membrane of the intestines was everywhere healthy, except in the *rectum* and *duodenum*. In the former situation, the rugæ were prominent, and of a violet colour ; in the latter, the membrane was of the colour of pale lees of wine. From the coincidence thus observed, we are led to believe that the same condition which produces congestion in the skin, may produce an analogous affection of the mucous membrane. Nor must we omit to note, in connexion with this subject, the fact of the mucous membrane of the vagina being occasionally found of a dark purple colour in those who die of puerperal affections.

In the cellular membrane, serum, lymph, and pus, may be deposited, mixed with each other in various proportions. The surrounding vascularity, in these cases, is unusually small, and the lymph effused not properly organized ; there is, consequently, no natural boundary to the disorganizing process, and the fluid secreted becomes infiltrated in the surrounding parts.

When the muscular structure is affected, suppuration takes place with great rapidity ; portions of muscles may be found quite soft, and sometimes pultaceous, in circumscribed patches, around which the fibre is perfectly healthy. Pus is occasionally deposited on the exterior of muscles ; and it will be then smeared over the surface, and rather infiltrated in the cellular tissue, than contained in a cyst. In the interior of muscles, there is the same absence of the natural limit to the inflammation ; but, owing to the more compact structure of the part, the deposits of matter generally remain circumscribed.

The brain and its membranes frequently present diseased appearances in those who die from secondary inflammation ; these, for the most part, may be altogether independent of any peculiar effects of the disease ; but, in some cases, it appears probable that they are not altogether unconnected with it. In one of the accompanying cases, the *pons Varolii* and *medulla oblongata* were found of a pink colour, in consequence of congestion, where the system had become contaminated by the absorption of diseased secretion ; and, in another, a layer of purulent lymph was found within the cavity of the arachnoid, accompanied by marks of inflammatory action in the fourth, and in one of the lateral ventricles.

The serous membranes are peculiarly liable to be attacked by secondary inflammation ; and, when affected, suppurate with the greatest readi-

¹ Journal de Physiologie, t. iv, p. 45.

ness. They generally exhibit but a slight degree of vascularity, and sometimes scarcely appear more injected than in their natural condition. In the peritoneal cavity, large quantities of unorganised lymph are frequently poured out, mixed with turbid serum or pus. The synovial membranes of joints, when affected, appear to run directly into suppuration, and will become distended with pus in a very short space of time. The pleura, on the other hand, will seldom suppurate at first; but lymph will be deposited upon its surface, and its cavity will contain turbid serum, occasionally mixed with blood.

XI. In the first sections, the changes produced in the blood, both in and out of the body, by the admixture of purulent or diseased secretions, were considered; and, in the last section, the post-mortem appearances observed in the different organs of those who have died in consequence of secondary inflammations, have been described. It now remains to connect these two series of observations, and to trace the relation that they bear to each other.

The most direct way in which diseased blood produces disease in the parts to which it is conveyed, is by communicating to them its own condition. Even foreign substances, which have no natural connexion with the body, may, in this way, be conveyed in the blood, and deposited in the organs of the body. Hunter relates a case,¹ in which a house-painter, who had been paralytic in his hands and legs for a considerable time, had his thigh broken, and died, about three weeks after, of the accident: "On examining the body, after death, the muscles, particularly those of the arms, had lost their natural colour; but, instead of being ligamentous and semitransparent, as happens in common paralysis, they were opaque, resembling exactly in appearance parts steeped in a solution of Goulard's extract. From this case it appears, that the lead had been evidently carried along with the blood into the muscles themselves. The blood can thus receive and retain extraneous matter capable of destroying the solids."

If foreign matter may, in this way, be conveyed to different parts of the frame, and there produce its chemical effects, it will readily be admitted, that a mechanical or vital action commenced in the blood,² may be continued in it when moved to a different part of the body.

In those who die of secondary inflammations, the result of such actions may frequently be traced from the seat of the primary injury even to the heart itself; and, as it has already been shown that contaminated blood will communicate morbid action to the vessels in which it is contained (section III), there no longer remains any difficulty in accounting for the sudden way in which this disease may fall upon a particular organ, or the unexpected manner in which the symptoms may shift from one part of the body to another. The conditions of the blood, which may be observed upon dissection, in this disease, so far as they tend to illustrate the present subject, may be included in two general expressions: 1st, those in which the blood has had a preternatural tendency to coagulate impressed upon it; and, 2nd, those in

¹ Op. cit. p. 99.

² It may appear unusual to speak of action going on in the blood; but, in so doing, we only illustrate the principle with which we commenced, viz., that "the blood has the power of action within itself."

which its coagulating power has, to a greater or less extent, been impaired. In one case, the blood is generally found of a dark colour, with firm, and sometimes adherent coagula in the vessels; in the other, it is thin and fluid.

When a preternatural tendency to coagulate has been impressed upon the blood, it will lodge in different parts of the vascular system, in situations which are most favourable to such an action; when, on the contrary, it has lost its coagulating power, extensive effusions may be expected, and the symptoms included under the term "gangrenous diathesis" will generally manifest themselves.

As the aggregate diameter of the vessels in the body decreases, the blood in them flows with greater rapidity and force; and when near the heart, it is placed in circumstances unfavourable to coagulation, in consequence of the rapid motion there communicated to it; and we accordingly find that, although adherent coagula may be formed in the veins leading from the seat of injury, and may be traced thence through other larger vessels, they will usually terminate abruptly, when these open into the vena cava. But when the blood arrives in the cavities of the heart, the tendency to coagulate may again manifest itself. Diseased coagula, presenting a "mottled appearance, partly brown, and partly of a dirty yellow colour", or "dark coloured, and partly composed of a yellowish grey opaque substance", or "with portions inelastic, and of an opaque yellow colour", will be found entangled among the projecting fibres of the auricles and ventricles. In the arteries, the blood is in the most unfavourable circumstances for coagulation during life, in consequence of the comparative smallness of their diameters, and the succession of impulses communicated to their contents; but even here, diseased and adherent coagula may be found in cases of secondary inflammations.

M. Cruveilhier¹ relates an instance in which, after child-birth, the pulmonary artery was found filled with coagula following its divisions. The principal clot had lost its colour, and was *adherent* to the sides of the vessel, and contained in its centre puriform fluid.

In another case, following the operation for necrosis, the following post-mortem appearances were observed. The left lung presented, in several places, patches of red hepatization, perfectly defined, and resembling so many spots of lobular inflammation. When cut into, these patches presented several puncta of puriform fluid. The *veins* of the lung contained fibrinous-looking coagula blocking up their cavities; and in the centre of these coagula was a whitish purulent looking fluid.

The diseased condition of the blood may thus be traced visibly from the original wound, through the larger vessels, to the heart, and again from the heart to the capillary system. The disposition to coagulate, once impressed upon the blood, is not destroyed by that fluid being conveyed to a different part of the body: the action may be retarded by motion in, as well as out of the body, but will nevertheless occur when it is placed under more favourable circumstances.

These circumstances, in the living vessels, are when the blood becomes separated into small quantities, and when it moves slowly

¹ Op. cit. p. 669.

along the capillaries. The blood will then coagulate in circumscribed patches,¹ as illustrated in the first characteristic marks of secondary disease which have already been mentioned as occurring in the lungs, the liver, the spleen, and the skin. The accompanying plate is taken from the lung of a donkey in which purulent fluid had been made to circulate with the blood. The stagnation of the blood, when the pus was first introduced, was mechanically prevented, and the livid spots produced by its subsequent coagulation in the capillaries of the lungs has been very faithfully represented. (See Experiment No. VI.)

"Besides the disposition for coagulation," observes Mr. Hunter, "the blood has, under certain circumstances, a disposition for the separation of the red globules, and probably of all its parts; for I have reason to believe, that a disposition for a separation of the red part and coagulation, are not the same thing, but arise from two different principles. This is always observable in bleeding; for if we tie up an arm and do not bleed immediately, the first blood that flows from the orifice, or that which has stagnated for some time in the veins, will soonest separate into its three constituent parts: this circumstance exposes more of the coagulating lymph at the top, which is supposed by the ignorant to indicate more inflammation, while the next quantity taken suspends its red parts in the lymph, and gives the idea that the first small quantity had been of such service at the time of its flowing, as to have altered for the better the whole mass of blood. Rest, therefore, may be regarded as one of the immediate causes of the separation." *Hunter*, p. 29.

This disposition of the blood to separate into its constituent parts is evinced in a very marked degree in one class of secondary affections. Extensive effusions of serum, lymph, and pus, mixed in different proportions, will take place in the serous cavities of the body, and become infiltrated in the cellular membrane, accompanied with very little vascular action in the parts. The colouring matter of the blood will also sometimes become effused with its other parts; but when this is the case, they will be found to have lost altogether the power of coagulation: in this respect presenting a direct contrast to the blood effused from a healthy wounded vessel. The lymph deposited will be found lying in unorganized flakes, wanting its usual adhesive properties, and very slightly attached to parts, presenting little or no increased vascularity. The rapid manner in which these effusions take place, shows that they are separated from the blood without undergoing any very elaborate process. In this condition of the system, any organ upon which the disease falls, may rapidly become disorganized and will readily mortify, and, after death, a tendency to rapid decomposition will be manifested. The veins on the surface of the body may frequently be traced as dark blue lines, as though the skin covering them were stained by the colouring matter of the blood. The lungs and other organs may, under these circumstances, be found in every grade of disorganization, till they present all the characteristics of gangrene: even the peculiar fetor which accompanies mortification of the lung, will, in some instances, be pre-

¹ The term ecchymosis does not appear appropriate to the discoloured condition of parts observed in the commencement of this disease; the blood is not at first extravasated from the vessels, as the term frequently employed would indicate.

PLATE SHÉWING THE FIRST MORBID APPEARANCE PRODUCED
BY THE STAGNATION OF VITIATED BLOOD IN THE CAPILLARIES OF THE LUNGS.

Surface of Lung with Pleura removed.

Section of Lung.

sent. In this condition may also be observed a tendency to the formation of petechial spots in different parts; and even the organs which do not appear to have been the peculiar seat of the disease, will be found to have lost their consistency, and to break down upon comparatively slight pressure.

Mr. Hunter found, that in proportion as the blood retained the power of coagulation, it had the power of resisting putrefaction; and conversely, we observe that, in this class of cases, the deficiency of the former is accompanied in a marked manner with the absence of the latter.

The two conditions of the blood which have now been mentioned, appear to bear a direct relation to the two classes of post-mortem appearances, observed in cases of secondary inflammations: the first being generally connected with congestion of different organs during the first stages of the disease, the second with extensive effusions, accompanied with comparatively little vascularity.

XII. The treatment of secondary inflammations naturally divides itself into local and constitutional, both as regards the primary lesion and the subsequent affections. The circumstances which interfere with union by the first intention in veins, have been shown to be the same as those which precede the formation of purulent deposits, in a large class of cases. Whatever then tends to favour the healthy reparation of a wounded vein, may be regarded as affording security against any subsequent disease; but the chief point in the local treatment is, perhaps, to prevent any accidental circumstances from interfering with the natural process of repair. When the powers of the constitution are enfeebled, even the natural motions of a part may interfere with recovery, and rest sometimes becomes an important object in the treatment. How necessary this is after child-birth, when the divided veins are being closed, every one who has attended such cases practically knows.

Again, after bleeding, the arm will inflame in a much greater proportion of cases, when the patient is obliged to follow his usual occupation, or when, from accidental circumstances (as from the pain experienced in Case i), the arms are kept in motion. It has occurred to me, to see the symptoms of purulent deposits set in, on the day following prolonged attempts to bring fractured portions of bone into position. In all such cases, any external violence (as in Experiment No. vi), or even the motion of the body, as in Dr. Davis's case (section iv) may loosen the coagula formed, either between the wounded edges, or in the cavities of veins. In the treatment of the local injury, a valuable hint has been left us by Mr. Hunter, connected directly with the consideration of the pathology of the disease. "The way in which sore arms after bleeding come on, shows plainly that they arise from the wound not healing by the first intention"; and he recommends that the two sides of the vein should be approximated by a compress, until union of the divided edges has taken place.

It has been shown upon very high authority, that the sides of a vein do not unite after venesection (as has sometimes been presumed to be Mr. Hunter's opinion); but that the divided edges only of the vessel are agglutinated by the coagulum, which "serves as a bed to the new

membrane." But the case is different, should this first attempt at union fail; the sides of the vessel may then become united, and its cavity for a time obstructed (section IV). The approximation of the sides of the veins would materially facilitate this action, which is the natural security, under the circumstances, against the admission of foreign matter. When an abscess is suspected to have formed in a vein, a similar mode of treatment is recommended by Hunter; the compression, in this case, being placed between the inflamed part and the centre of the circulation. In the pathological museum of the College of Surgeons, one of Mr. Hunter's preparations (No. 1728) exhibits such a case, where, from the imperfect union of a vein, the contents of the abscess had become mixed with the blood.

As the process of reparation has been variously described by authors, so the different theories propounded have led to different kinds of treatment. At the Veterinary College, even within the last few years, it was publicly taught, that a coagulum in a vein was a foreign substance, and ought to be removed; and the jugular vein in horses which had been bled, was sometimes slit up for several inches, in order to remove the coagula which formed in successive portions of its course.

That a coagulum in a vein may be an irritating substance has been fully proved (sections I and II); but the irritation depends upon the accidental admixture of foreign matter: and the process that consequently takes place requires to be carefully distinguished from the natural mode of union by the first intention.

To remove coagula which have formed round purulent secretion (if such could be recognised), might be to remove a cause of irritation; but to remove them in ordinary cases, is to remove the very means prepared by nature for the restoration and safety of the part. In operations involving large vessels, the local conditions which may influence the actions in the veins, appear not to have been fully determined; for while some surgeons regard the tying or cutting of a vein as a serious operation, others are in the constant habit of doing it with impunity. In operations upon hæmorrhoidal tumours, the veins involved are of some size, and in two of the instances recorded in the Appendix, a ligature applied to them was followed by purulent deposits. In the usual operation of passing a needle armed with a double ligature through the base of such a tumour, the hæmorrhoidal veins are necessarily sometimes wounded, and there is danger that, in tying the ligatures, the sides of a wounded vein may be drawn asunder. The vessel may thus be held open, and be in a similar condition to the vessels contained in bony structures.

A safe way of performing this operation, when admissible, is to destroy a portion of the mucous membrane with strong nitric acid. The blood in the vessels then becomes charred, and their cavities obstructed, till they are permanently closed by adhesive inflammation. Every means of treating a local injury which tends to produce healthy union or adhesion, may be considered in some sort as a preventive mode of treatment. Position, topical applications, bandages, and temperature, may all have their influence in producing these healthy actions; but, as the wounds which precede purulent deposits are generally characterized by feeble powers, those conditions which tend to invigorate

the parts are principally indicated. "When action is greater than strength, whatever has the tendency to raise the power above irritability should be used: the object of this practice consists in bringing the strength of the constitution and parts as near upon a par with the action as possible, by which means, a kindly resolution or suppuration may take place, according as the parts are capable of acting." The irritability of a wound frequently appears nothing else than a series of attempts to bring about an action, which it has not the power to accomplish: as soon as that is fulfilled, which the necessity of the parts demands for their healthy condition, irritation will cease. In such cases, everything that will confer strength to carry out the intended action, will prevent inflammation. But no local applications will be sufficient to produce this effect, unless the powers of the constitution are supported at the same time.

A heavy man received a wound in the back of his head from an iron spike; he was kept very low, complaining occasionally of want of food. Repeated hæmorrhage took place at intervals of a day or two, which no local applications could suppress, and he ultimately died from loss of blood. The wound was found to extend through the bone into the lateral sinus, which contained only some fluid blood. No inflammatory action had taken place within the skull.

The constitutional treatment of wounds, liable to be succeeded by purulent deposits, is most important; for during the healing of the primary wound, the system may be influenced by remedies, which may be subsequently quite useless. It is evident, observes M. Cruveilhier,¹ that the treatment of phlebitis ought to be concentrated on the first period of the disease, viz., that of the coagulation of the blood; for as soon as pus has become mixed with the circulating blood, medicine is generally of no avail. When there are indications of the extension of inflammation along a vein, the mode of treatment usually adopted in this country has been the administration of calomel and opium; and, in France, general bleeding, but especially the application of large numbers of leeches. "We may subdue inflammation of veins, whatever their situation, by general blood-letting, and especially by local bleeding, repeatedly sufficiently often and in sufficient quantities!"² It is true that patients recover after such treatment; but the published records of cases not unfrequently terminate with such a description as the following: "The patient was repeatedly bled, and with apparent relief every time, the blood being extremely sizzly. *Two days, however, previous to death, the vital principle was so exhausted as to need the use of cordials!!*"

If the explanation of the mode of adhesion in veins already given be correct, neither the propriety of bleeding, nor of the administration of mercury, as a rule, derives much support from a consideration of the pathology of the disease. Careful comparative experience is required, before either of these plans can be adopted as a recognised mode of treatment. Satisfactory evidence in favour of the employment of these remedies is still wanting; and, indeed, the numerical evidence which we have in some instances, tends to indicate an opposite plan of treatment.

¹ Op. cit. p. 662.

VOL. II.

² Ibid.

At a period when puerperal fever was rife, forty cases, attacked with some form of the disease, "were treated without any bleeding or leeching, or without any attempt to induce the constitutional effects of mercury; and of these, only two died." "In irritable habits, when the inflammation becomes more diffused," says Mr. Hunter, "bleeding should be performed with great caution: even a quick, hard pulse, and sizzly blood, are not always to be depended upon as sure indications of bleeding being the proper method of the resolution of the inflammation; more must be taken into the account. The kind of blood is of great consequence to be known; for although it should prove sizzly, yet if it lies squat in the basin, and is not firm in texture, and if the symptoms at the same time are very violent, bleeding must be performed very sparingly, if at all; for I suspect, that under such a state of blood, if the symptoms continue, bleeding is not the proper mode of treatment. If we had medicines which, when given internally, could be taken into the constitution, and were endowed with a power of making the vessels contract, such, I apprehend, would be proper medicines. Bark has certainly this property, and is of singular service, I believe, in every inflammation attended with weakness; and therefore, I conceive, should be oftener given than is commonly done."

In bleeding women suffering from puerperal fever during an epidemic, in which the disease frequently terminated in purulent deposits, I have frequently seen an irregular transparent bluish layer form upon the surface of the blood, almost immediately after the operation; in consequence of this, a repetition of the bleeding was sometimes had recourse to, when the subsequent stages of the disease appeared to indicate that it had not been required. The transparent layer of lymph on the surface of the blood, may, in such instances, be only an indication of its tendency to separate into its different parts, as previously described. Large bleedings, under such circumstances, inasmuch as they tend to diminish the already enfeebled power of coagulation in the blood, may predispose to the infection of the system, and to the formation of purulent deposits. The action of mercury upon the system, as illustrated in Case xxvi, may have the same tendency.

The action of mercury, so far as it can be traced upon the surface of the body, is certainly unfavourable both to union by the first intention, and to adhesive inflammation; and, inasmuch as the safety of the patient depends upon the due performance of these processes after an injury or wound, its effect upon the system must be regarded as of at least doubtful advantage.

When salivation has been induced, serum is poured out, and the texture of the gums is loosened and rendered spongy. When lymph is effused upon the iris, the action of mercury loosens its adhesions, and dissolves its connexions; it cannot, therefore, be supposed that its action upon the system should render the union of divided vessels stronger, or the newly-formed adhesions in wounds firmer. The cases in which purulent deposits usually form, indicate a debilitated state of constitution. They are of much more frequent occurrence in large towns than in the country, and in hospital than in private practice. The de-

¹ FERGUSON (Dr.) On the Diseases of Women. These cases formed one-fifth of the whole number treated.

pressing influences which they give to erysipelas, or puerperal fever, will also predispose to the formation of these abscesses; and as scanty diet, loss of blood, debilitating surgical operations, and over-crowded rooms, have been found among the causes of the former, so may they be looked upon as favouring the production of the latter. The Treatment of Inflammation of the Veins, in which purulent deposits frequently originate, is thus spoken of, from the result of lengthened observation: "All the experience that I have had on the subject, would lead me to believe that, like erysipelas, it has its origin in a low asthenic state of the system, and that those persons are especially liable to it, who have been much lowered by hæmorrhage at the time of an operation, or by too scanty a diet afterwards. An operation is a shock upon the system, making a great demand upon the vital powers. The effects of this shock are often much aggravated by loss of blood, and a very scanty diet actually makes the patient more liable to some kinds of inflammation. Our mode of practice ought to be rather to sustain his powers by allowing him wholesome nourishment, and not to add to the influence of the other depressing causes, the still worse one of starvation."¹ But as there are many circumstances, besides mere debility, which tend to produce a depressing effect upon the vital powers, so tonic medicines, and a generous diet, must not be looked upon as the only preventive modes of treatment. Each case that occurs in practice may present some peculiarity; in one case, the patient will be found to have previously suffered from diabetes; in another, from disease of the spleen; and in a third, from organic disease of the kidney; and, in each of these, a peculiar mode of treatment may be required. The general treatment of this disease, after the formation of purulent deposits has commenced, is probably as little satisfactory as any that come under the notice of the surgeon. The whole character of the affection is characterized by what has been aptly designated as action without power. The excitement of the system will imitate all the acts of genuine inflammation, without any of its healthy results; and loss of energy will appear immediately after, or even arise in conjunction with, the first symptoms of increased action.

In the treatment of such deposits, it must be borne in mind, that they are not the disease, but the effects of a concealed cause, which may develop itself in any part of the system; and that, while the relief of one organ is sought by depletion, additional vigour may be given to the latent evil, so as to render it more ready to develop itself in some other part. When purulent deposits occur, the sudden congestions, which indicate their commencement, are not accompanied by any reparative actions, and the lymph, which is effused in the subsequent stages of their formation, is not so disposed as to circumscribe and limit the inflammation; there is, therefore, no natural process by means of which such collections of fluid can be evacuated; hence, when situated near the surface, they should be opened as soon as they are detected. It will sometimes happen, that, after the symptoms of secondary inflammation of internal organs have commenced, an abscess will present itself near the surface of the body, and relief will be afforded to the part first affected; at other times, an obstinate and violent diarrhœa will precede

¹ BRODIE (Sir B. C.) *Medical Gazette*, vol. xxxvii, p. 642.

recovery. Although the mode of treatment adopted may have little influence either in bringing on, or checking, such salutary actions, it is important to watch their occurrence, and perhaps still more important, to be careful not to mistake a remedial action for a symptom of the disease.

"What treatment," says Cruveilhier, "shall we oppose to purulent infection? To this question experience is as yet dumb, while theory would seem to point to diffusible stimuli and tonics; to ammonia, quinine, and sudorifics; to hot external applications, to the vapour baths, to purgatives, and especially to emetics; to tartarized antimony, in large doses; to vesicatories, and to strong diuretics. Calomel has been extensively employed, to create a fluxion from the intestinal mucous membrane; but all these means have failed as signally in my hands as in those of others; yet, when the injection of putrid matters into the veins of living animals has been followed by abundant and very fetid evacuations, they have usually got well. It is a fundamental fact of pathology, that the intestinal canal is chiefly affected in diseases caused by miasmata. I am certain that diseases resulting from purulent infection would not be stamped with the seal of incurability, and that nature, seconded by art, would triumph in the majority of cases, if the pus, which is incessantly renewed, did not incessantly renew the sources of infection. As soon as constitutional symptoms manifest themselves, neither general nor local bleeding affords any advantage. A portion of the *materies morbi* is, no doubt, abstracted with the blood; but, as it is constantly being reproduced, the constitution is only deprived of the power it would otherwise have of resisting the disease."¹ In accordance with this remark, M. Gaspard found that animals, which recovered after the injection of a certain quantity of pus into their veins, often died when the experiment was repeated. The recovery was usually preceded by black, liquid, and extremely fetid evacuations, which often seemed to afford immediate relief. When such evacuations have taken place in other diseases, the gall-bladder has been found distended with black bile;² and it appears probable that the liver, in these cases, is one of the principal organs through which the cleansing of the system is attempted. If the supply of morbid matter to the system could be checked, mercurial action, in this stage of the disease, might be of service, by enabling the liver, or other organs, to throw off their vitiated secretions. When patients recover from purulent deposits (Cases III and XXVII), they are often left in a debilitated and languid state, in which ordinary tonics exercise little influence. The consequences of the disease appear to hang about the system, long after the cause which gave rise to it has ceased. The pulse will sometimes continue irritable, and there will be a tendency to derangement of the secretions of the skin, bowels, and other organs, accompanied by occasional slight attacks of fever. In this condition, an alterative course of mercury, combined with sarsaparilla, has been found beneficial. These remedies, by stimulating the activity of the excreting organs, may assist the constitution in throwing off the disease; and their mode of action may be the same as in other cases, where the system has been infected by an animal poison.

¹ Op. cit p. 662.

² I have observed this condition after injuries of the spine which proved fatal, by causing inflammation of other parts.

APPENDIX OF CASES.

A. CASES OF SECONDARY INFLAMMATION, ACCOMPANIED BY MORTIFICATION OF THE SKIN.

CASE I. William Ford, 33, an apparently healthy man, was seized with the symptom of strangulated hernia, at Harrow, December 28th. He was bled in both arms; but, as the hernia remained unreduced, he was sent to town, in a state of considerable restlessness and suffering. During his journey, the hernia returned of its own accord. He left quite relieved, and remained well till the 31st, when he had a rigor. On the 1st of February, there was a considerable degree of febrile excitement, and the anterior part of the left arm was red, swollen, and painful upon pressure. The tenderness and pain soon extended up the arm, in the course of the cephalic vein: the redness assumed the character of erysipelas. On the 3rd, he suffered from sickness, the matter vomited being green and acid. He had two severe rigors, followed by great heat of skin, and a quick full pulse. The orifice in the left arm discharged a thin serous fluid. On the 4th, he complained of stiffness in the right arm; also of being very feeble and faint. 5th. Wandered much during the night. The right arm was a good deal swollen about the end of the elbow-joint, and presented a patch of a bright red colour above the condyle. The pulse was weak and tremulous; motions of the hands was occasionally observed. He complained of pain in the ring-finger of the right hand, and in the ball of the right great toe. The skin, in both these situations, had assumed a red appearance. 6th. Was delirious during the greater part of the night; countenance expressive of much anxiety; pulse very weak; tongue covered by a brown dry fur; profuse perspiration; complained of severe pain in the calves of his legs. He died during the following night.

Post-mortem appearances. A small deposit of thick pus was found on the external surface of the left cephalic vein. The blood was coagulated in its cavity, its coats were thickened, and its lining membrane appeared very red. This redness could be traced, though in a less degree, through the whole extent of the vena innominata. On the right side, the cellular tissue, both above and below the elbow-joint, was greatly distended with serum. The cephalic vein of the right arm presented marks of having been inflamed; but not nearly to the same extent as upon the opposite side. The joints in which pain had been experienced during life, were distended with turbid pus; deposits of pus were also found in the anterior mediastinum, and between the œsophagus and trachea. A large gangrenous spot was observed in the skin of the calf of the right leg. The subjacent cellular tissue was much distended with serum.

CASE II. Richard Mason had a small cancerous tumour removed from the lower lip. The operation was performed in the usual way, and the wound appeared to heal by the first intention; a small abscess, however, followed by some ulceration, subsequently appeared in the neighbourhood of the cicatrix. Nine days after the operation, he complained of sore throat and general uneasiness, and three days afterwards he was seized with rigors, followed by cold perspiration and coma.

Post-mortem appearances. A small deposit of matter was found,

situated beneath the fascia of the left thigh. The synovial membrane of the left knee was highly inflamed, and contained a large quantity of pus. On the right side, the skin of the whole inferior extremity presented a dark livid appearance, with the exception of that situated upon the fore part of the thigh. The same dark colour was observed in the muscles of the limb, which were infiltrated with blood and serum. The arteries and veins were discoloured, but in other respects presented nothing remarkable. It was ascertained, that, two years previously, this patient had suffered from diabetes; and upon examining the urine found in the bladder, it was found to contain sugar.

CASE III. Jane Thornton, æt. 32, came under treatment on the 22nd of March. A week previously, her right ankle had become red and painful, and inflammation subsequently extended up the inner side of the leg. When first seen, she was evidently much out of health, although no one organ could be said to be particularly affected. On the 28th of March, she was attacked with severe rigors, and experienced pains in different parts of her limbs: the rigors were repeated for several days in succession. The inflammation of the leg now entirely disappeared, and she complained of pain in the right knee, which was slightly swollen. On the 31st, her bowels became much relaxed; there was great general depression, and much nervous agitation. Both knees were swollen. The general symptoms now became somewhat relieved; but on the 4th April, she was attacked with vomiting, which recurred frequently during the day, and was accompanied with great depression, and severe pain in the epigastrium. On the 5th, the sickness continued, apparently quite uninfluenced by any remedies. She passed considerable quantities of blood by stool: her countenance presented a dusky yellow hue: the pulse was excited, without power, and the sense of depression was greatly increased. On the 7th, the vomiting still continued, and she still passed blood by stool. Some spots of a dark purple colour now made their appearance upon her face. The hands both became slightly swollen; and upon the right one, some small, dark, livid spots, similar to those upon the face, made their appearance. She was much troubled with hiccough. On the 9th, the countenance was very anxious, the complexion more sallow: some more livid spots appeared upon the face and cheeks. The right hand and arm were swollen and painful: some fresh livid spots appeared, upon the knuckles. Complained much of faintness: had extreme debility with occasional hiccough: pulse extremely weak: the surface of the body was covered with cold perspiration. The sickness had entirely ceased, and there was no blood in the motions: the tongue was rather dry in the centre, but tolerably clean. 10th. She wandered slightly during the night, and vomited once: some blood again appeared in the motions; the right hand and arm were less swollen. 11th. The countenance was anxious, the pulse about 90, and intermitting irregularly. The vomiting recurred several times. The livid spots on the right hand had not increased in size, but appeared like distinct small black superficial sloughs of the skin; these all scabbed off, without supuration. 12th. The pain and swelling of the arm had nearly subsided. The bowels acted very freely with the aid of medicine, and she expressed herself much relieved. From this time the patient slowly but gradually improved, with one or two slight intermissions, till the beginning of

May, when she again complained of pain at the inner part of the right arm, above the elbow. Some hardness could here be felt in the course of the basilic vein. Some small collections of matter were now deposited upon the back of the right hand, resembling, in some respects, the eruption of confluent small-pox. On May 11th, she had regained much of her strength, but still felt some pain in the elbow upon motion. She also complained of the joints of one of her fingers. She now left London for change of air.

The following Table is formed of Cases taken consecutively during One Year.

Heads of Cases.	Period of secondary inflammation.	Post-mortem appearances.
<p>CASE IV. Elizth. Mackintosh, æt. 25. Inflamed bursa patellæ; erysipelatous inflammation in the neighbourhood of the right axilla; sudden suppression of the catamenia; rigor; peritonitis; tongue covered with yellowish white coating; sickness; "catching pains" in the epigastric region.</p>	<p>A few days after the appearance of the erysipelatous inflammation; three days before death.</p>	<p>Dark-coloured serum, mixed with shreds of recently effused lymph, in the cavity of the left pleura; large quantities of sero-purulent fluid, with recently effused lymph, in the peritoneal cavity.</p>
<p>CASE V. James Stevens, æt. 46. Punctured wound of the finger whilst opening a rabbit; diffuse cellular inflammation of the right arm; spasmodic and "catching pains", principally referred to the epigastrium; expectoration of bloody fluid.</p>	<p>Twenty-seven days after the injury; about seven days before death.</p>	<p>Bloody fluid in the cavity of the left pleura; pus in the left elbow-joint.</p>
<p>CASE VI. William Collins, æt. 36. Bruise of the patella, caused by the wheel of a carriage; apparent recovery; erysipelatous redness over the left knee; rigor; rapid pulse; hot skin, followed by perspirations, pains in the head, restlessness, delirium.</p>	<p>Twenty days after the accident; three days before his death.</p>	<p>Cavity of the knee-joint containing a quantity of thick grumous fluid, apparently a mixture of blood and synovia; fibrous degeneration of a portion of the cartilage of the patella; turbid serum in the sub-arachnoid cellular tissue; bloody puncta in the brain, larger and more numerous than natural; posterior part of both lungs gorged with blood.</p>
<p>CASE VII. Maria Martin, æt. 39. Caries and necrosis of the tibia, with large ulcer of the leg.</p>	<p>Several months after the occurrence of caries of the tibia; a few days before death.</p>	<p>The skin of the upper part of the leg and the whole of the thigh of a mottled appearance, caused by extensive dark patches of incipient gangrene; the cellular tissue of the limb infiltrated with lymph and pus.</p>
<p>CASE VIII. George Mason, æt. 42. Compound comminuted fracture of the metacarpal bones; inflammation of the absorbents; erysipelatous redness of the skin; secondary abscess in the affected arm; rigidity of the muscles of the tongue; trismus; universal affection of the muscles.</p>	<p>Nine days after the injury; twenty-two days before death.</p>	<p>Increased degree of congestion, both in the grey and white substance of the brain; substance of the pons Varolii and of the medulla oblongata of a pinkish colour, and presenting irregular streaks of increased vascularity; spleen soft, and somewhat congested.</p>

Heads of Cases.	Period of secondary inflammation.	Post-mortem appearances.
CASE IX. Sarah Leg, <i>set.</i> 50. Necrosis of a portion of the tibia, accompanied by a large foul ulcer.	A few days before death.	Effusion of serum and lymph in the cellular tissue, which surrounds the pharynx and cesophagus; inflammation and ulceration of the mucous membrane of the larynx; slight inflammation of the lungs; the spleen of a greyish red colour, more solid and more easily lacerated than natural.
CASE X. Elizabeth Moleno, <i>set.</i> 42. Strangulated femoral hernia of the left side; operation; erysipelatous blush around the wound upon the third day, followed by sickness, cold perspiration, and delirium; several dark patches upon the skin of the right leg.	Eight days after the operation; four days before death.	The lining membrane of the right internal saphenic vein of a dark livid colour throughout, the cavity of the vessel filled with a large quantity of coagulated blood mixed with puriform fluid; pus in the common iliac vein; effusion of serum around the veins of the leg; liver large and congested; mottled degeneration of both kidneys.
CASE XI. Jane Cox, <i>set.</i> 60. Scalp wound; erysipelas of the head and face; transverse fracture of the external malleolus, followed by suppuration of the ankle-joint.	A short time before her death.	Mortification of the skin of the lower part of the leg, ankle, and foot; slight extravasation of blood into the arachnoid cavity, and into the substance of the brain; kidneys coarse in structure, and remarkably soft.
CASE XII. Bartholomew Sullivan, <i>set.</i> 27. Lacerated and contused wound of the leg, followed by diffuse cellular inflammation, and inflammation of the absorbents; delirium; a separate large patch of mortification, surrounded by bright red congestion, appeared in the right groin the day before his death.	Five days after the accident; eight days before his death.	Mortification of the skin and cellular tissue of the right leg; the veins of the limb healthy; spleen of a pale colour, and very soft.
CASE XIII. George Foscutt, <i>set.</i> 24. Fracture of the femur into the knee-joint; rigors; erysipelas of the limb, ill defined and very slow in its progress; coma; abscesses in the leg and thigh; mortification of the skin on the dorsum of the foot, and over the left hip.	Six days after the accident; twenty-two days before his death; pain in the chest the day before his death.	Hepaticization of both lungs, with secondary abscesses in the left one; kidneys soft and coarse in texture, the left presenting a small deposit of apparently tubercular matter; spleen large, pale, and soft.
CASE XIV. William Wright, <i>set.</i> 30. Fracture of the patella; erratic erysipelas; diarrhoea; abscesses in the leg, and in the knee-joint.	Erysipelas appeared a month after the fracture of the patella, and about the same time before his death.	Effusion of lymph in the pleura; secondary abscesses in different stages of formation in both lungs, and deposit of lymph in one kidney.
CASE XV. Henry Bateman, <i>set.</i> 19. Fracture of the fibula; diffuse cellular inflammation of the leg; suppuration in the knee-joint; necrosis of a portion of the fibula.	Three months after the accident; three weeks before his death.	Recent effusion of lymph upon the pleura; incipient secondary abscess in both lungs; the tibia exposed, and its structure of a black colour, and soft.

Heads of Cases.	Period of secondary inflammation.	Post-mortem appearances.
<p>CASE XVI. John Clark, <i>set.</i> 45. Large scalp wound; rigor; followed by paralysis of one side; a portion of bone exposed, of a darkish green colour, and when removed of a putrid odour.</p>	<p>Nineteen days after the accident; three before death.</p>	<p>Effusion of lymph between the dura mater and the bone, and of pus and lymph between the cavity of the arachnoid; pus, mixed with blood, in the superior longitudinal sinus; deposit of lymph in the structure of the pia mater; recently effused lymph in the cavity of the left pleura; secondary abscesses of the left lung.</p>
<p>CASE XVII. Matthew Elmes, <i>set.</i> 37. Injury of the wrist; diffuse cellular inflammation; abscesses in the cellular tissue, and in the wrist-joint; lower extremity of the radius denuded; pains in various parts of the body, especially the head and abdomen.</p>	<p>The tenth day after his admission into the hospital; four days before his death.</p>	<p>Secondary abscesses, in various stages of formation, in both lungs; suppuration between the different bones of the carpus; both kidneys large, coarse in texture, and flabby; the spleen soft, easily lacerated, and of a pale colour.</p>
<p>CASE XVIII. Mary Hopkins, <i>set.</i> 19. Ulceration of the cartilages, followed by suppuration of the knee-joint; amputation; erratic erysipelas over various parts.</p>	<p>Erysipelas appeared a week after amputation; sixteen days before death.</p>	<p>Low inflammation of a portion of the left lung; dark-coloured patches of deposit in the spleen.</p>
<p>CASE XIX. John Wilkinson, <i>set.</i> 56. Compound fracture of the right tibia; inflammation around the wound, with collections of matter; slight delirium; sickness.</p>	<p>Pain in the right side a week after his admission.</p>	<p>Turbid serum in the right pleural cavity; incipient secondary abscesses in both lungs; large cysts in the kidney.</p>
<p>CASE XX. James Bryant, <i>set.</i> 20. Scalp wound, denuding the bone; puffiness of the scalp upon the eighteenth day; rigors, followed by profuse perspirations; restlessness; delirium; projection of the eyeballs.</p>	<p>Twenty-four days after the accident; five days before death.</p>	<p>Bone exposed to the extent of a shilling, of a yellow colour, and with a very dark diploë; effusion of lymph and pus between the dura mater and the bone, extending to the base of the skull, and through the sphenoidal fissures into the orbits; effusion of pus into the arachnoid cavity; incipient secondary abscesses in the lower lobe of the left lung: spleen large, and very soft, mottled; degeneration of both kidneys.</p>
<p>CASE XXI. James Williams, <i>set.</i> 41. Fracture of the lower extremity of the left radius; diffuse cellular inflammation of the arm at the expiration of four weeks; abscesses in the limb, one of which communicated with the fracture; diarrhoea; vomiting; tongue dry and brown.</p>	<p>Ten weeks after the accident; one week before death.</p>	<p>Large cavity containing foul matter, in contact with the sacrum, which was exposed; abscesses between the bones of the left wrist and hand; spleen soft, congested, and grumous.</p>
<p>CASE XXII. John Munday, <i>set.</i> 36. Prolapsus ani; hæmorrhoids; rigor; anxiety of countenance; great heat of skin; pulse 150.</p>	<p>The eighth day after the operation for hæmorrhoids; the fourth before death.</p>	<p>The mucous membrane of the whole of the large intestine of a very dark colour; congested patches of it thrown up into prominent folds; recently effused lymph upon the right pleura; secondary abscesses in both lungs.</p>

Heads of Cases.	Period of secondary inflammation.	Post-mortem appearances.
CASE XXIII. Esther Polley, <i>æt.</i> 50. Lacerated wound of the foot; separation of a small portion of the base of one of the metatarsal bones; pain in chest; rapid pulse; depression, with delirium.	The tenth day after the accident; the third before death.	A double fracture of the fifth metatarsal bone; inflammation of the right pleura; secondary abscesses in right lung.
CASE XXIV. Henry Lacy, <i>æt.</i> 26. Scalp wound, exposing the bone; fracture of the skull; signs of nausea; pains in the head; "soreness of the stomach"; drowsiness and insensibility; paralysis of one side; muscular twitchings; portions of bone removed by the trephine, near the top of the head.	The eighteenth day after the accident; and a week before death.	Yellow matter in the diploë of the parietal bones, in the neighbourhood of the part where the trephine had been applied; effusion of lymph upon the surface of the dura mater; pus and lymph in the posterior half only of the longitudinal sinus; effusion of pus in the cavity of the arachnoid; some bloody serum in both pleural cavities; incipient secondary abscesses in the liver. Vessels on the surface of the brain congested; lateral ventricles distended with fluid; a thick layer of purulent lymph upon the arachnoid membrane at the base of the brain; some slight spots of ecchymosis on the anterior surface of both lungs.
CASE XXV. Thomas Meed, <i>æt.</i> 15. Injury of the leg; small suppurating sinus upon the outside of the limb; erysipelas; diarrhœa; coma.		
CASE XXVI. Thomas Daffey, <i>æt.</i> 42. Hæmorrhoids; operations; rigors; sickness; great abdominal pain; diarrhœa; hiccough. This patient had been salivated previous to the operation for the hæmorrhoids.	The tenth day after the operation; and ninth before death.	Incipient secondary abscesses in the right lung; liver studded, throughout its whole extent, by secondary abscesses; mucous membrane of the rectum of a dark greenish colour; effusion of pus and lymph into the hæmorrhoidal and inferior mesenteric veins; cavity of the left knee-joint distended with pus; spleen soft, pultaceous, and thickly-studded with dark-coloured blotches.

CASE XXVII. George Burton, *æt.* 22, a stout navigator, was first seen September 18th, 1848. He had an enormous slough of the skin and cellular membrane, covering the lower part of the abdomen on the right side. He gave a most imperfect history of himself, and seemed frequently incapable of comprehending the questions which were put to him. The skin was hot and dry, the pulse 130. For several days, he remained in the same apathetic condition. The bowels were particularly obstinate, and the purgative medicines which were administered produced no effect whatever.

When the slough separated, the surface of the external oblique muscle was left as clean as if recently dissected. The skin was undermined for some extent, and no attempt was apparently being made to limit the progress of the disease, by the effusion of lymph. Fresh portions of cellular tissue consequently became affected, and the whole surface ultimately exposed was full six inches in diameter. *Sept.* 25th. Complained of some pain in the chest and upper part of the abdomen; has a constant short cough; perspires very freely. 26th. The pulse has

become weaker, but remains of the same frequency (130). It communicates a peculiar jerking sensation to the finger. 28th. Pulse 96, weaker; slight diarrhoea; he vomited several times during the day. 29th. Restless, with delirium. 30th. Passed another restless night, but became better in the course of the day; his appetite returned. *Aug. 5th.* His appetite again failed; complained of a sense of distension in the abdomen. 8th. An abscess was discovered at the lower and back part of the right leg; the skin over this presented various shades of yellow and brown, giving the appearance of having been extensively bruised. About two ounces of discoloured pus were evacuated, together with a considerable quantity of grumous blood. 10th. Has much improved since the last report; the wound discharges dark semi-coagulated blood. 13th. A second abscess now presented itself in the same leg, and the skin covering it presented the same discoloured appearance as in the first instance. When opened, it discharged dark-coloured semi-coagulated blood with the matter. He now rapidly improved in health, and at length perfectly recovered.

CASE XXVIII. Samuel Todd, æt. 58, fell from a wagon fifteen miles from town, and was brought up in an open cart, during a sharp frost. There was a compound fracture of the left leg. Two days afterwards, he suffered from cellular inflammation around the wound. On the 24th day, he had a slight rigor, and complained of slight stiffness in the right shoulder. He ate and drank well till within two days of his death, which took place on the thirty-fourth day.

Post-mortem appearances. Both lungs contained secondary abscesses, in various stages of formation. The left external iliac and common femoral veins were blocked up with firm coagula, and confined by these was a quantity of purulent-looking fluid; an adherent layer could, for some distance, be peeled off the internal surface of these vessels.

CASE XXIX. James Howard, æt. 33, had a small abscess in the dorsum of the right foot, which was followed by inflammation of the absorbents; abscesses subsequently formed in the right thigh and groin. Two months after the first appearance of the abscess on his foot, he was seized with rigors, vomiting, and profuse perspiration. He complained also of pain in the lower part of the left lung, and it was said that the respiratory murmur was deficient in this situation. He died on the 23rd day from the appearance of the symptoms of the secondary affection.

Post-mortem appearances. The skin and conjunctivæ were of a light yellow colour; there were incisions in the right groin. In the cavity of the right pleura, were some patches of recently effused lymph, of a yellowish colour. In the lower lobe of the right lung, were several patches of secondary deposit; these were found in different stages of their formation, and some of them were of considerable size; some of the pulmonary veins, in the neighbourhood of those which had suppurated, were found to contain a fluid resembling pus. On the left side was a small abscess, under the pleura costalis. The liver was healthy in structure, rather larger than usual. Kidneys healthy. The veins in the groin presented a healthy appearance.

CASE XXX. A gentleman was operated upon for hæmorrhoidal tumours in the usual manner, and died shortly afterwards, with effusion of

bloody serum into one of the pleural cavities. The only peculiarity that could be detected in the appearance of the rectum was, that the blood was still fluid in one of the largest veins. This vein led directly to an ulcer produced by the operation; and, even at its extremity, it contained no appearance of coagulum.

B. CASES IN WHICH VITIATED FLUIDS WERE OBSERVED IN THE VEINS OF THE UTERUS AFTER CHILD-BIRTH.

CASE XXXI.¹ On the second day after a natural labour, a woman had prolonged rigors, which were followed by abdominal pain and much perspiration. On the fourth day, she was affected with syncope and bilious vomiting, accompanied by extreme prostration. On the fifth day, the abdominal pain, which had subsided, re-appeared with agitation and delirium. On the sixth, she had cold sweats, irregular pulse, rapid breathing, and vomiting. She died on the following morning.

Post-mortem appearances. There was a small quantity of limpid serum in the cavity of the peritoneum. Some of the uterine veins were found to contain a turbid fluid. The heart was half filled with brown blood. The lungs were congested, and the other organs natural.

CASE XXXII. A woman, twenty-two years of age, of a nervous temperament, miscarried at the expiration of the first month. She was almost immediately afterwards attacked by rigors and bilious vomiting, accompanied by pains in the loins and in the hypogastrium. On the following day, there was coma, with indications of extreme debility. On the third day, consciousness returned. There was difficult articulation; and moaning. The extremities became cold, the pulse imperceptible, and death occurred upon the same day.

Post-mortem appearances. The peritoneum was slightly injected, and contained a small quantity of reddish serum. The cervix uteri was covered with a layer of pus. Semi-transparent lymph was found in some of the uterine veins. The brain, and other organs, were found healthy.

CASE XXXIII. A feeble girl, 21 years of age, was confined at the expiration of the eighth month. On the fourth day, there was shivering and prolonged fainting. The following morning, she had acute abdominal pain, fever, and diarrhoea. On the seventh day, the symptoms had all subsided, but on the eighth the abdominal pain returned, accompanied by syncope. She died two days afterwards.

Post-mortem appearances. The peritoneum was slightly injected, and contained about a pint of red-coloured serum. The uterus was large, white, and firm, its veins half filled with fluid blood; its lymphatics natural; its inner surface lined with a layer of fetid brown blood, but otherwise healthy; the cervix covered with a grey, thin exudation. The lungs, heart, brain, and other organs, were quite sound.

CASE XXXIV.² Anne Biggs, æt. 39, confined March 18th, 1830, eighth child. On the evening of her confinement, her manner was much excited. On the 19th, she was incoherent, and complained of pain in the calf of the right leg, which was tender on pressure. The pulse being hard, she was bled to eight ounces. On the 28th, the leg was swollen and white; the pain in it much increased: towards evening the

¹ This, and the two following cases, are taken from Tonellé. The author is responsible for the accuracy of all the preceding cases.

² Dr. Ferguson.

calf of the limb became black, while, at the tendo-achillis, the skin was hot, tender, dry, and mottled. The bowels were much opened, the head giddy, the pulse quick and strong. She was again bled, to twenty-six ounces, and twelve leeches were applied to the temple. On the 21st, there was nausea, vomiting, and diarrhoea. On the 23rd, she complained much of the confusion in her head, the leg was tolerably easy, but the upper and inner part of the thigh was very tender. On the 24th, the diarrhoea continued, and there was increased weakness. A hard swelling, about half as large as an egg, appeared at the wrist, and one of the orifices made by venesection was black and painful. She died in the evening.

Post-mortem appearances. All the ventricles of the brain were dilated with serum, and there was a good deal of effusion into the arachnoid and pia mater. The viscera were perfectly healthy, except the heart and spleen: the latter was very large, and on pressing it, a large quantity of dirty red pulpy matter exuded. The lining membrane of the right auricle and ventricle, when washed, had a dark red hue. The femoral vein, just at the ingress of the saphæna, and the superficial vein in the back of the leg, had their coats much thickened, so as to cut like arteries. Their lining membrane was similar to that of the right cavity of the heart. When they were divided, purulent matter, mixed with thin light-coloured blood, escaped. The cellular tissue forming the sheath of the femoral vessels, and on the calf of the leg, shewed marks of recent inflammation; but there was no appearance of pus in these situations. None of the glands in the groin or ham were enlarged. The inferior vena cava appeared healthy.

C. CASES OF PURULENT DEPOSITS, CONNECTED WITH INFLAMMATION OF THE VEINS AFTER DELIVERY, RECORDED BY DR. ROBERT LEE
IN THE MED.-CHIR. TRANSACTIONS.

CASE XXXV. Mrs. Mayhew, æt. 33, was delivered on the 2nd March 1829. On the 5th, there was a discharge of blood from the uterus. From the 6th to the 20th, she made no complaint of uneasiness in any region of the body, though her strength rapidly declined. The countenance was of a dusky yellow hue. The heat of the surface slightly increased; the respiration hurried, particularly on bodily exertion, and the pulse above 130, and feeble; the tongue pale and glossy, with loss of appetite. The lochial discharge had a peculiarly offensive smell. She died upon the 28th of March.

Post-mortem appearances. When the uterus was laid open, there was found to be a portion of the placenta, about the size of a nutmeg, in a putrid state, adhering to its inner surface. The substance of the uterus, to the extent of an inch around this, was of a peculiarly dark colour, almost black, and as soft as a sponge. On cutting into it, about a teaspoonful of purulent matter escaped from the veins, and a small additional quantity was pressed out from them. . . . On opening the capsular ligament of the right knee-joint, about six ounces of thin purulent matter escaped, and the cartilages of the femur and tibia were extensively eroded. There was no appearance of inflammation, however, on the exterior of the capsular ligament. The right wrist was swollen, and the cellular membrane around it was unusually vascular, and infiltrated with serum.

CASE XXXVI. Mrs. Pope, æt. 40, was delivered, on the 26th of Oct.,

of her fourteenth child, and appeared to recover favourably until the 3rd of Nov., when she was suddenly attacked with a severe rigor. This was followed by intense head-ache, vomiting, general soreness of abdomen, and suppression of lochia. Nov. 6th. Great prostration of strength; laborious respiration, with pain at the bottom of the sternum, and frequent hacking cough; pulse 135, extremely feeble; skin hot and dry. . . . Occasional retching and vomiting. Several hard, lumpy cords were found running up in the inside of the thigh, in the direction of the superficial veins. 7th. Delirium; general debility greatly increased; the surface of the body was covered with a yellow suffusion; the middle finger of the left hand was much swollen around the second joint, and the skin covering it was of a dusky red colour.

CASE XXXVII. Mrs. Edwards, æt. 35, was suddenly attacked, three weeks after delivery, with pain in the calf of the right leg, and loss of power in the whole right inferior extremity. On the 5th day from the attack, a considerable swelling, without induration, had taken place from the ham to the foot, and great tenderness was experienced along the inner surface of the thigh to the groin. The extremity became universally swollen, painful, and deprived of all power of motion. The temperature along the inner surface of the limb increased; the integuments were pale and glistening, not pitting upon pressure. The femoral vein, from the groin to the middle of the thigh, was indurated, enlarged, and exquisitely sensible; pulse 80; tongue much loaded; thirst; bowels open. On the 23rd day from the attack, the disease was apparently declining. The femoral vein could no longer be felt, but there was still a sense of tenderness in its course down the thigh, and she experienced considerable uneasiness between the umbilicus and pubes, as well as in the loins. She now, for the first time, began to have rigors, accompanied by a quick pulse, loaded tongue, and thirst. From this period to the 31st day, the swelling of the limb and tenderness in the course of the femoral vessels subsided, but she experienced attacks of acute pain in the umbilical region, loins, and back, which assumed a regular intermittent form. Every afternoon there was a rigor of an hour's duration, followed by increased heat, and profuse perspiration: the skin was hot and dry; pulse 125; tongue brown and parched; bowels open. These febrile attacks gradually declined in severity, and she appeared to recover till the 43rd day, when she had a long and violent fit of cold shivering. The countenance now became expressive of great anxiety, and the pulse extremely weak and feeble. 45th day. Vomiting; pain upon the left side, increased upon taking a deep respiration. 46th day. Another severe and prolonged rigor; skin hot and dry; pulse 140; tongue brown and parched; diarrhoea; respiration hurried, with frequent cough; surface of body presenting a peculiar yellow tinge. The conjunctiva of the right eye now suddenly became of a deep red colour, and so much swollen, that the eyelids could not be closed. On the day following, the left eye also became red and swollen, the surface of the body was cold and clammy, pulse 140, extremely feeble, with great debility, and repeated attacks of vomiting. From this date, the patient lived nearly three weeks, but for the last fortnight the vision in both eyes was entirely destroyed.

Post-mortem appearances. The left pleural cavity contained upwards of two pints of a thin, purulent fluid. The inferior lobe of the left lung

was of a dark colour, and soft in texture. In its centre, about an ounce of thick cream-coloured pus was found deposited in its dark and softened texture. This was not contained in any cyst or membrane, but was infiltrated into the pulmonary tissue. The coats of the vena cava inferior were considerably thickened; its whole cavity was occupied by a coagulum, terminating above in a loose pointed extremity. The left common iliac vein was plugged up, by a continuation of the coagulum from the cava. The coagulum was continued beyond the entrance of the internal iliac, which it completely closed, and terminated in a pointed extremity about the middle of the external iliac. Neither the remainder of the vessel, nor the femoral vein, exhibited any morbid changes. The right common iliac vein was contracted to more than one-half its natural size; it was firm to the touch, of a greyish blue colour; to its internal coat adhered an adventitious membrane of the same colour, containing within it a firm coagulum. The internal iliac was rendered quite impervious by dense, dark-coloured bluish membranes; and, at its entrance into the common iliac, it was converted into a solid cord. The contracted external iliac contained within it a soft yellowish coagulum; its coats were three or four times their natural thickness, and lined with dark-coloured membranous layers. The right femoral vein, from Poupart's ligament to the middle of the thigh, was diminished in size, and almost inseparable from the artery. Its tunics were thickened, and its interior coated with a dense membrane, surrounding a solid purple coagulum strongly adherent to it.

D. CASE, SHEWING THE PERIOD AT WHICH A COAGULUM MAY GIVE WAY
IN A WOUNDED VESSEL.

CASE XXXVIII. George Haydon, ætat. 37, received a wound about half an inch in length over the right radial artery, on March 5th, 1848. The hæmorrhage was arrested by pressure. On the 12th, a small slough formed in the bottom of the wound, the edges of which were inflamed and painful. On the 14th, slight bleeding from the wound occurred, which at first was arrested by the application of cold; but in the evening it recurred in considerable quantity, and again was checked by cold and pressure; during the night, however, profuse hæmorrhage again took place, and was only stopped by the application of the tourniquet above the elbow. On the 15th, the radial artery was tied; but as this did not prevent the hæmorrhage from returning, when the tourniquet was relaxed, the original wound was enlarged. The opening in the radial artery was now with some difficulty discovered; it extended two-thirds round the circumference of the vessel, leaving one-third undivided.

E. CASES SHEWING THE ORGANISATION OF THE OUTER LAYER OF
EXTRAVASATED BLOOD; REPORTED BY MR. HEWETT.

CASE XXXIX.¹ A middle-aged man received a severe injury of the chest; he lived eleven days after the accident, and during this time he never presented a single inflammatory symptom. The cavity of the left pleura was found completely filled with bloody fluid, and was subdivided into two compartments, by a portion of coloured fibrine, presenting a honey-combed appearance, which passed from the ribs to the lung. The lower compartment was itself subdivided into several others, by layers of

¹ Med.-Chir. Transactions, vol. xxviii.

coloured fibrine intersecting each other. Large portions of loosely coagulated blood were found in all the cavities; some of these clots were of a rusty colour, others approached nearer to the natural colour of the blood. The lung was compressed against the spine, and the whole surface of the pleural sac was coated by a false membrane, about two lines in thickness, formed by coagulated fibrine. The fibrine which lined the pleura pulmonalis and pleura diaphragmatica, presented on its inner surface a smooth and polished appearance, and in colour exactly resembled the yellowish fibrine found in the clots of the heart of this patient. So uniform was the coating, and so continuous was it throughout its whole extent, that it looked at first merely like thickened pleura; but this appearance was easily destroyed, by peeling off this adventitious membrane from the serous tissue, which there presented the same appearances as the pleura on the opposite side, with the exception of not being quite so smooth: there was neither thickening nor the slightest increase of vascularity in this pleura. A large rent, from which the hæmorrhage had proceeded was found in the substance of the lung.

CASE XL. A man was attacked with diffuse cellular inflammation of the inferior extremity, which terminated in two days with extensive gangrene of the skin. In the superficial and common femoral veins were extensive coagula; these did not completely fill the veins, but slightly adhered at different points to their internal coats. These clots still retained, in some places, the colouring matter of the blood, whilst at others the colourless fibrine alone remained; in both veins, the clots were enveloped in a perfectly transparent, smooth, and polished membrane, presenting the appearance of a serous tissue. In the structure of these membranes were several distinct arborescent vessels, minutely injected;¹ some of these vessels were of sufficient size to allow of the blood being made, by gentle pressure, to circulate through them; but no communication could be traced between these vessels and the coats of the veins. The membranes were easily peeled off from the surface of the clots with which they were in contact. The interior coats of the veins presented their natural colour and polished surfaces, except at the points where the slight adhesions above-mentioned existed.

ON THE MECHANISM OF TEXTURAL NUTRITION.

By RUDOLF HAAS, M.D., late Lecturer on Epidemiology in Vienna.

THE first step in the actual process of nutrition, is the attraction of the homogeneous part of the blood by the tissues.² The blood being contained in the vessels, that part only which transudes through their walls into the parenchyma of the organs can contribute to nutrition. Hence, even though a large quantity of rich healthy blood be circulating through the vessels, the textures will be imperfectly nourished, unless sufficient

¹ Mr. Gray, the Curator of the Museum of St. George's Hospital, has recently shewed me the outer layer of an effusion of blood into the arachnoid cavity, injected from the middle meningeal artery.

² VALENTIN'S Physiologie, Band 2, Sect. 2068. MUELLER'S Physiologie, p. 75.