The *Bladder* was partly filled with natural urine, which escaped by an ulcer into the vagina.

The *Uterus* was enlarged, very dark, and congested, containing an ovum of about seven or eight weeks; ovaries very large, and in one a large corpus luteus.

HEAD AND SPINE. The vessels of the scalp; dura mater, and pia mater, were much distended with dark fluid blood. Some serous effusion was present in muscles of the pia mater, upon the convolutions; none in the ventricles. The substance of both cerebrum and cerebellum was perfectly natural, except, perhaps, having more vascular points on being sliced. The medulla oblongata was firm and natural. The spinal veins were congested; the medulla spinalis itself was natural, except perhaps (but of this I am doubtful) rather softer than usual at the bulbous expansion about the seventh cervical vertebra.

At the inquest, I suggested a chemical analysis of the contents of the stomach, which I had secured; but, as there was an absence of any suspicion of the death having been occasioned by any one than herself, and, on the question being put to me, whether I could give a confident opinion as to strychnine being the cause of death, which I could, the jury, to whom the question was left by the coroner, did not consider it necessary.

Subsequently, for my own satisfaction, with the kind assistance of my friend Mr. Morley, an analysis was made on Wednesday night, one hundred and fourteen hours after death, and eighty-four after examination, when strychnine was most decidedly detected by us in considerable quantity, by the method mentioned in toxicological works. The interior of the stomach was well washed with distilled water, which, being added to the contents taken from the stomach, and acidulated with sulphuric acid, was boiled for some little time, and filtered; then neutralised with lime, and again filtered, and evaporated to dryness: a portion of this residue, on being dissolved in spirit, gave, on the addition of a small quantity of nitric acid, the characteristic brown tint of strychnine.

So much was obtained in this manner, that a solution in spirit was left at our School of Medicine, to enable Dr. P. Smith, the lecturer on medical jurisprudence, to exhibit to the students from time to time the tests for strychnine from that actually obtained from the stomach of one poisoned by it.

I was not then aware that there existed any insurmountable difficulty in detecting strychnine within a short period after death, nor do I now understand the grounds upon which such an opinion rests, unless it be meant that, when the patient lives for some considerable period after the administration of the dose, the substance may be digested, and thus all traces be removed before death; but I much doubt whether such a dose, unless frequently repeated at intervals for some time, so as to keep up its action and a constant spasm, would prove fatal, unless by exhausting the nervous power, when symptoms would be different. If the dose be sufficient to cause death within four hours, I suspect it may commonly be found; and that it does not necessarily undergo decomposition for a considerable period after death, the present case furnishes conclusive evidence.

BRONZED SKIN: DEATH: POST MORTEM EXAMINATION.

By C. M. THOMPSON, Esq., Westerham.

In connexion with the subject of bronzed skin, mentioned by Dr. Budd, I will relate a case which occurred in my own practice, and may be interesting, because it was followed by a *post mortem* examination. It occurred many years ago; but, owing to its peculiar circumstances, is indelibly impressed upon my memory.

I was called to a woman "of colour", about thirty years of age. The skin over the whole body was dark bronze; the palms of the hands were white; the conjunctivæ were perfectly clear and white.

Having no doubt of her being a woman of colour, but a little puzzled by the shape of her features, I inquired from what country she came? At this question she seemed vastly amused: assured me she was a native of my own neighbourhood, and reminded me that I had attended her on many occasions, some years before, while residing with an aunt in my town, at which time she was a fair and pretty girl. Having overcome my astonishment, I proceeded to examine her case. She stated that this discoloration had followed upon a fever, and had gradually increased. She was now suffering from severe dyspepsia, loss of appetite, frequent vomiting, emaciation, and also from violent spasm, resembling gall-stone. There was no proof, however, of bilious obstruction; the eye was clear, the urine limpid, and the motions of a bright ochre colour.

I need not detail the treatment; nothing gave her relief; and after lingering for two months, she died from an unusually severe attack of spasms. The *post mortem* examination showed all the viscera in a healthy state; but the gall-bladder was greatly distended, and its duct entirely obliterated, while the hepatic duct was pervious.

I may here mention that I have, in the course of my practice, met with four cases of what is vulgarly called "black jaundice." In all these the bronze discoloration was very great; but there was a marked difference between these cases and the former one, inasmuch as the orange eye and the obstructed bile clearly indicated the nature of the disease. In two of these I was enabled to make a *post mortem* examination, and in both I found the gall-ducts entirely obliterated; in one, from cancer of the pancreas; in the other, from simple occlusion of the common duct.

THE METEOROLOGY OF 1855.

By J. A. HINGESTON, Esq., Brighton.

COMPARED with the fatal year of 1854, the weather of 1855 was both fine and healthy. The commencement of January was warm, the temperature being as much as 12° Fahr. above the average; but it was followed by severe cold, which lasted from the 18th of January to the 21st of February. There was a good deal of snow and ice; and on the 18th of February, which was the coldest day of all, Fahrenheit's thermometer fell as low as 9°. The feathered tribes suffered severely; the thrush family sought shelter in the public thoroughfares of towns and villages, and both land and water birds were driven instinctively to the more southern parts of England. Many shrubs and perennials were killed by the frost.

From March to November the weather was fine. The summer was mild and pleasant, and the autumn warm. There were six months of continued fair weather, and the number of fine days throughout the year amounted to 196. The mornings were particularly beautiful, and the starlight nights in the autumn most enchanting.

On comparing the climate of the Crimea with that of the south coast of England, a great similarity will be observed between the two countries. They are both upon the same isothermal line. (See *Kaemtz's Meteorology*, plate vi.) Troops encamped on the open downs near Brighton, far removed from their supplies, and harassed with the slow operations of a siege, would not have suffered less during the winter here than the allied armies did on the plateau between Sebastopol and Balaklava; and the hurricane that broke upon this coast on the 25th of October last would not have proved less disastrous than that which wrecked so many of our transports against the rocky coast of the Crimea on the 14th November, 1854—particularly so large a craft as the steamship *Prince*, heavily laden, with a lee-shore close astern, and nothing but a single stream anchor to hold her head against the fury of the storm.

From the 25th of March to the 5th of May, a drought prevailed, relieved only by a few light showers. In May, 1.50 inches of rain fell, but the drought returned, and continued till the 7th of July. The grass was burnt up, or its