as well as the external wound), he gradually sank, and died about three months after the operation.

Post mortem examination showed the ileum and colon, terminating separately in the wound, the mucous membrane of the intestine being continuous with the rest of the wound, so as to shut off the cavity of the gut from that of the peritoneum.

Original Communications.

CASES IN MIDWIFERY.

By THOMAS RADFORD, M.D., F.R.C.P. Edin., etc., Consulting Physician to St. Mary's Hospital, Manchester.

SECOND SERIES.

THIS series of cases will include those which are accompanied with hæmorrhage—commencing with those of "placenta prævia".

CASE I. On June 24th, 1820, I was requested by Mr. Dick to visit a woman who resided in Greengate, Salford, who was reported to be flooding: she was in labour of her third child, and had a profuse discharge of blood. I found she had previously flooded twice, at intervals of about three weeks; but each of these attacks was comparatively trifling. There was no obvious cause for them; and on the patient being put to bed, the discharge soon began to diminish, and required no further measures besides rest in the recumbent position.

The present flow had abated when I arrived; but it was, and had been, increased on the accession of pain. On making an examination, I found the os uteri dilated to about the size of a crown-piece; it was soft and dilatable; a small portion of the placenta with the membranes was felt tightened during the pain, but hanging loosely over the os when it (the pain) subsided. The head of the child was easily touched, as the uterus acted briskly and effectively; and as the discharge of blood was now comparatively trifling, it was thought prudent not to interfere, but to leave the case to nature. In the course of two hours, the membranes ruptured; the head descended into the pelvis, pressing the separated portion of the placenta aside as it progressed, and very soon the child was born alive. The uterus contracted well; there was only an ordinary discharge; in about half an hour afterwards, the placenta was expelled. No further hæmorrhage happened. The bandage placed on before delivery was now tightened. She recovered without the slightest interruption. Those portions of the placenta which had been separated previous to labour, were readily distinguished from that (the last) which caused the flooding at this time, being more consolidated and of a different colour.

CASE II. Mrs. Buckley (on 24th March 1821) desired me to visit a hospital patient residing in Hanover Street, who was said to be in labour, and flooding. When I arrived, I found the discharge, which at first was great, considerably abated. This was her fifth pregnancy. She had felt pains four hours, which had now become severe; during each pain the flow of blood had been slightly increased from the commencement of the labour, but was trifling in amount until the time when I was sent for. Her pulse was, however, unaffected, and her countenance natural. The os uteri was dilated to the size of a half-crown, and felt thin and soft; through it, I found an edge of the placenta, which organ adhered to the right latero-posterior part of the cervix uteri; the distended membranes were readily perceived during the pain, and the head of the child was felt on its remission.

I remained with the patient some time, during which the pains increased in intensity and effect; the discharge of blood gradually ceased. I now left her in the charge of the midwife, with an express order to send for me again if the bleeding recurred. I heard nothing until the following

day, when she called, and stated the membranes ruptured in about two hours after I left; the child was born alive; the placenta was naturally expelled; there was no further loss than what usually takes place. The woman recovered well.

CASE III. On February 21st, 1822, I was requested by Mrs. Bradley to visit a hospital patient residing in Cumberland Street, who was reported to be flooding. She had now arrived at the eighth month of her second pregnancy, and had been quite well up to this period. She began flood without any known cause. Mrs. Bradley had put her to bed, and placed a bandage round her, and used cold applications. The flow of blood was said to have been profuse, but it had considerably lessened when I saw her. I examined her, and found the os uteri closed, and therefore I could not ascertain whether the placenta was situated near it or not, although I suspected it was, from feeling the cervical walls fuller and thicker than they ordinarily are at this period of pregnancy; and also, as no external cause had occurred to account for the accident.

When I left the patient, I desired Mrs. Bradley to send again for me if there was a return of the hæmorrhage. I however heard nothing more of the case until a month had elapsed, when I was called to her to extract the placenta, which I found slightly adherent. Her labour up to this point had been natural and easy. The placenta was readily detached and removed, and there was no flooding. I carefully examined this organ, and readily distinguished that portion which had been separated and caused the accident, from the rest of it; its structure had become firmly condensed, and less vascular.

CASE IV. In June 1830, I was requested to visit a lady pregnant for the third time. My opinion was sought, because in her two former labours the children were born dead. She was now nearly seven months advanced; her health was good; and she said she felt the child very lively. In a fortnight after this interview, I was sent for, as she was flooding. I found her on the bed, and looking pale; her pulse was rather quick and feeble. I was told she had lost a large quantity of blood. Although the bleeding was now comparatively trifling, yet it continued. She had no pain; and having placed my hand on the abdomen, I felt the uterus flaccid; I found the os uteri closed, and the cervix uneffaced. A bandage was placed round the body, and pinned tightly over the uterus. Cold water with vinegar was externally applied, and the apartment kept cool. An unstimulating diet was ordered, and quietude in the recumbent position strictly enjoined. Nothing further was required, as the hæmorrhage gradually ceased, and did not return. She progressively improved after this accident, and she continued to feel the child to be vigorously alive. Arrived at the eighth month, it was for a few days felt less sensibly, and then altogether ceased to move. At this time the breast lessened in size, and became more flaccid. About a week before the expiration of the ninth month, labour came on. When I visited her, the pains, now brisk and frequent, had existed for two hours. I found the os uteri considerably dilated, soft, and yielding. The membranous bag, with a portion of the placenta, was felt; but there was not the slightest discharge of blood. The os uteri continued rapidly to dilate; the membranes ruptured; and the separated (now loose) flap of the placenta was pushed aside, and passed by the descending head. The pains being very strong and forcing, the child (dead) was soon expelled. There was not the slightest hæmorrhage. The placenta was soon extruded, which, on examination, was found had lost its usual fresh vascular character; the portion which had been separated at nearly the seventh month, and then caused the flooding, was readily fixed upon.

CASE V. The following case recently occurred to my respected friend and colleague Mr. Skinner, who has kindly permitted me to cite it:—

On October 28th, 1855, 5 o'clock p.m., I was called by one of the midwives to attend Ann Maion, aged 23 years, who had been attacked, while in bed this morning, with

hæmorrhage. She is nearly at her full time, and states this is her third pregnancy, and the two previous labours were perfectly natural, and she recovered from both without a single unfavourable occurrence. Both of the children are now living.

At about the seventh month of this (the present) pregnancy, she had some slight hæmorrhage while engaged in some domestic occupation, but there was nothing in the nature of this which she could assign as a cause of the flooding, which was unaccompanied by pain. At this time she was ordered a mixture with sulphuric acid diluted, and opium, which speedily checked the dribbling discharge, which had continued. She felt the movements of the child soon afterwards. She has been free from the discharge up to my visit. On now making an examination, I found one or two small coagula of blood in the vagina; the os uteri dilated to about the size of a shilling, and I thought I detected the placenta presenting on the left latero-posterior part of this part of the uterus. She had then no pain, and, since the morning, no return of bleeding. Her pulse was tolerably good. Plumb. acet.; tinct. opii; rest; cold applications to the vulva; and the room to be kept cool.

Monday, October 29th. She has had no return of hæmorrhage; has no pain. She slept part of the night, and has taken some gruel at intervals. On an examination, I found the os uteri in the same condition as before reported. She was this evening carefully and safely conveyed (in the horizontal position) on slings to the (St. Mary's) hospital. Ordered to take an opiate draught.

October 30th. No return of hæmorrhage; she has had a comfortable night. She has no pain; she takes nourishment. Her pulse is good. Dr. Radford examined her with the stethoscope; he heard the fetal cardiac sounds; the placental souffle most distinct over the left lateral portion of the cervix uteri.

November 1st, 2 o'clock P.M. Labour suddenly commenced. The membranes ruptured, and a full grown living foetus was expelled by two or three strong pains. Mr. Runcorn (our Resident-Apothecary) was summoned up stairs. There was not the slightest flow of blood, although Mr. Runcorn had to introduce his hand to detach the remaining large portion of placenta, which was rather firmly adherent. There was only the ordinary lochial discharge. She rapidly recovered, and was discharged cured, on November 10th.

REMARKS. These cases are, in my opinion, sufficient in number to prove the fallacy of the dogmatic rule, which inculcates that active operative interference is necessary in all in which the placenta is implanted on the lower (cervical or oral) portion of the uterus.

These labours were, in the main, natural, and required only the same management as is adopted in such cases. It is true there was an unavoidable contingent—the abnormal position of the placenta; but, fortunately, the perils which frequently happen in placenta prævia, and not only threaten the woman's safety, but sometimes destroy her, were here naturally averted.

In all these cases (and in others of similar kind which have come within my observation) the mothers survived, and all the children but one were born alive; the death of this one was not traceable to any incident which happened before or during the labour, but was to be attributed to some constitutional cause which had previously destroyed the lives of two children. Examples like these are important to warn against rashness, but ought not to lead carelessly to overlook the possibility of dangers supervening, even in cases which at first promise a propitious termination. On the contrary, we should ever be watchful, and endeavour to make them lessons to teach us to adopt, artificially in other cases, the same plan which nature has wisely pointed out in these. They afford us ample evidence that several conditional circumstances must exist in order that a safe termination of labour, without artificial aid, may take place in cases of placenta prævia. A limited portion of the placenta must only be connected with the dilating portion of the uterus, whilst the great bulk of it

(the placenta) must be placed beyond the sphere of those actions which prepare for the egress of the child. The edge of the placenta, which is usually fixed to the os uteri in such cases as the above, is usually comparatively thin, and contains fewer and lesser sized vessels than those which belong to the central portions of this organ. The degree of separation of the edges of the placenta in the above cases was small, so that the natural anti-hæmorrhagic powers were equal to suppress the bleeding. The death of the child in Case iv, doubtless contributed to prevent a repetition of the flooding.

(For a more full exposition of my opinions on the subject, *vide* remarks "On the Suppression of Uterine Hæmorrhage without Artificial Assistance."—*Lancet*, vol. i, 1847, p. 296.)

The soft dilatable state of the os, and the energetic action of the body and fundus uteri, which recurred in these cases, most certainly contributed, in some measure, to a happy issue.

[To be continued.]

CHLOROFORM IN MIDWIFERY.

By EDWARD W. MURPHY, A.M., M.D., Professor of Midwifery in University College, and Obstetric Physician to University College Hospital.

[Read before the Medical Society of London, January 26th, 1856.]

HAVING already brought the use and advantage of chloroform in the practice of midwifery before the profession on three different occasions, I should not now venture to do so a fourth time were it not that on the last occasion, when the president (Dr. Snow) honoured this Society with a valuable paper on its application to surgery, there seemed to be on the part of the Fellows a strong desire to discuss its application also to midwifery. They were, however, prevented from doing so, because that question was not properly before them; and if I now introduce this subject, it is because I have been apprised of his intention to read another paper on chloroform; and I am persuaded the society will admit that it is desirable that the use of this agent should be freely examined wherever it may be employed.

I have not the slightest reason to deviate from the opinion I first expressed in 1848, of its extreme value, not only in relieving the sufferings of the parturient female, but also in promoting her more favourable recovery; but I regret to perceive that the question is even yet so enveloped in clouds of prejudice that it is scarcely capable of a fair examination.

I cannot but protest strongly against the violent and most unphilosophic anathemas which have been hurled, not only against the agent itself, but all who use it. Such a mode of treating an important subject has not added to the reputation of those who adopt it, nor of the Society where it has been thus introduced. I shall be happy, indeed, to find that this our Society—the oldest Medical Society in London—shall be distinguished for discussing the properties and value of this new agent with that moderation and gravity which befits its age.

In the use of chloroform in midwifery there are three points to which I should wish to direct the attention of this Society. *The first* is its progressive influence on the nervous system: *the second*, the difference that exists in its effects upon different constitutions: *the third* is the remarkable disparity in the amount of suffering from the pains of labour which women experience when going through what is commonly called "natural labour".

From the truth of the first I derive the safety of its application: from the second, the importance of a cautious and guarded use of it. The third proposition will, I think, prove its infinite value, not only in assuaging an amount of agony which overcomes the fortitude of women of the strongest resolution, but also in promoting their more rapid recovery.

The progressive influence of chloroform on the nervous