"One is entitled to expect that so novel and startling a fact, so dangerous if false, so revolutionary if true, would be admitted only upon proof proportioned to its improbability. Far from it. Like other marvels, the more marvellous, the more facile its reception. It is picked up in a province of a country where medical men are more numerous, more learned, more inquisitive, more addicted to inquire into the wonderful, than perhaps any other country in Europe. Nevertheless, the first and only authority for the fact, in its details, is a professed traveller, Von Tschudi; who after visiting the New World and describing its wonders, returns to retail even greater wonders in the Old. So far as I can discover . . . Von Tschudi describes merely what he was told, and personally tested nothing. He did not even see any clear skinned Styrian peasant take his frightful dose, or afterwards scale his lofty mountain. Yet upon such sorry authority, the fact appears without challenge or hesitation, for the first time, I believe, in detail in an English dress in *Chambers' Journal*. Thence it is transferred into various other public channels of information, gathering as it rolls, till at last one source of intelligence for the community apprises the world, that "it is a very common practice in some parts of the Continent—in Germany, Hungary, and the Tyrol—for persons of all classes to take arsenic in minute quantities as a tonic and stimulant!"

These strictures of Professor Christison possess great weight, as being the deliberate conclusions of one long distinguished by his intimate acquaintance with the properties of poisonous substances.

A physician in Vienna, Dr. Wurmb, has expressed similar opinions. These have been communicated to the *ASSOCIATION JOURNAL* (Nov. 1855), by Dr. George N. Balfour. Dr. Wurmb says that this statement is without foundation—*hat keinen grund*. Dr. Wurmb, practising at Vienna, may be supposed to be as capable of learning the truth of such statements as those residing at great distances.

It is, moreover, a very significant circumstance, that no countenance of this wonderful trick of poison eating is to be found in either Henke's *Zeitschrift für Staatsarzneikunde*;—nor in Casper's *Vierteljahrsschrift für Gerichtliche Medicin*—nor in the *Deutsche Zeitschrift für Gerichtliche Medicin*—nor in the *Annales d'Hygiène*—nor in the *Journal de Chimie Médicale*—all of which journals devote the whole or a portion of their pages to toxicology, and are the exponents of continental medico-legal and toxicological science. Such silence is ominous, and goes far to confirm the reasonable doubts of those who receive such wonders after transmission through so many hands.

[To be continued.]

CASES OF LITHOTRITY AND LITHOTOMY.

By DRAPER MACKINDER, M.D., Fellow of the Royal Medical and Chirurgical Society, etc., Gainsborough, Lincolnshire.

THE operations of lithotomy and lithotritry having been a good deal discussed of late, the two following cases will not probably be wholly destitute of interest, more especially the latter, which is very uncommon, if not without parallel.

CASE I. Lithotritry in a Female. M. Martin, aged 23 years, a spirited, intelligent, and active little woman, a servant in a public house, consulted me, in September last, for a pain at the lower part of the abdomen, which frequently became excruciating, and a weight and constant straining and desire to micturate. Ever since her childhood she had been subject to paroxysms of pain; but from the inattention of her mother, and, subsequently, her own false delicacy, no medical advice had been obtained until a few weeks before I saw her, when, in consequence of an augmentation of her sufferings, she sought the opinion of an elderly practitioner, who thought he could cure her with medicine. Not having, however, derived any benefit from the treatment which had been adopted, she allowed me to ex-

plore the bladder, when a calculus was discovered, of about the size of a duck's egg. She had of course tried various domestic and empirical remedies, not the least pernicious of which was brandy and water, which she had been induced to drink liberally early in the morning, for the purpose of gaining temporary relief.

On September 27th, assisted by Mr. Septimus Lowe and Mr. Jacobson (and afterwards by Dr. Baxter and Mr. Towle), I performed lithotritry, while the patient was under the influence of chloroform. The irritation that followed was not more than might have been anticipated, and was relieved by opiates. I operated again on October 12th and 20th, November 5th, December 10th and 28th, January 12th and 25th, February 4th and 25th, in all ten times, crushing the stone on an average seven times at each operation. The calculus was an alternating one, the triple phosphate forming its chief ingredient. When dried the detritus weighed fifteen drachms.

This was a very instructive case, not only on account of the great suffering which the poor young woman had endured for so long a period, with so little absolute mischief to her general health, but from the quantity of chloroform necessary to induce anæsthesia. She generally inhaled an ounce, and more than once the resulting depression was so great, that it was thought expedient to protract the periods of its administration. Twice, an ounce was consumed with no better effect than the production of talkative excitement. I have observed, indeed, in many other cases, that when the patient gets into a very talkative humour, it is almost useless to persevere with chloroform. This young woman informed me on these two occasions, after half an ounce had been inhaled, that it was useless to continue, as she felt that she could not go to sleep. Twice this operation was postponed by the appearance of the catamenia.

M. M. has been in perfect health ever since the last portion of calculus passed away, and feels herself "a new being". The external layers of the calculus were composed of the triple phosphate, and the internal, according to the analysis of Mr. John Lowe of Edinburgh, of uric acid and oxalate of lime.

CASE II. Lithotomy. William Gilby, aged 9 years, came under my care in January 1854, when I was requested to devise some means of curing him of what was considered a mere dirty habit of wetting the bed, for which he had been punished many times without avail. On perceiving an elongation of the prepuce, and hearing the boy's statement that he had a constant inclination to urinate, and often the most intense pain, which I soon had an opportunity of observing, I passed a sound into the bladder, and discovered a stone, which by its acute ring, I believed to be a mulberry calculus. It was situated above the pubis, and never changed its position, though it was struck by the sound (which it scratched), many times prior to its removal. The urethra was small, and its dilatation caused so much suffering that I was compelled to have a staff made for me, those in common use being too large.

On January 25th, assisted by Messrs. Jepson, Sharp, and Jacobson, I performed lithotomy by the lateral operation, the boy being anæsthesised by chloroform. On introducing the forceps into the bladder, I found that I could not extract the calculus, though I could feel it; which convinced me, as I had previously suggested to my medical coadjutors, that it was encysted. I then passed the forefinger of my right hand into the bladder, and with my left bore heavily on the abdomen, when I could distinctly feel the cyst or pouch, from which a portion of the calculus protruded. By a little manipulation I was fortunately enabled to dislodge the stone, which fell into the bladder, and was extracted with ease. It was a beautiful mulberry calculus, weighing exactly three drachms. The little fellow did well, and was cured in six weeks.

REMARKS. Whilst this patient was under my care, he had two attacks of spasmodic croup, one a week or two prior to, and the other about a week after, the operation. The first, I referred to the irritation pro-

duced by the calculus; the latter, to one of two causes, the presence of a second stone, or worms. I prescribed a strong dose of calomel and scammony, and had my diagnosis confirmed by the expulsion of two large lumbricoides, which were doubtless the cause of the first attack.

I do not know whether this be an isolated case of encysted calculus being removed by lithotomy, but I cannot find a similar case related by the authorities in my small library. If my memory serves me rightly, there was something said at one of the societies last session about adherent and encysted calculi. This was certainly not simply adherent, but encysted for about three-fourths its surface, the margin of the cyst feeling somewhat like the os uteri when partially dilated.

I need scarcely remark, that I considered myself exceedingly fortunate under the circumstances, though the *vox populi* is not always the *vox Dei*.

ON ECZEMA INFANTILE.

By FRASMUS WILSON, Esq., F.R.S.

[Read before the Annual Meeting of the British Medical Association, July 31st, 1856.]

INFANTS at the breast and young children are peculiarly subject to eczema, and in them it is apt to assume the severest form presented by cutaneous disease. In young infants it commences at the end of the first month or six weeks, and unless submitted to proper treatment may continue for months and years—in fact, it may lay the foundation of a cutaneous disease, which may be prolonged in a chronic form until manhood, or may hang about the patient for the remainder of his days.

Eczema infantile, like eczema adultorum, originates in malassimilation, and with good reason is commonly ascribed to a faulty secretion of milk on the part of the mother: but when once established, it is not remedied, as might be expected, by the withdrawal of the cause and the substitution of a different and less faulty food. Unsuccessful attempts to cure the disease probably carry the child on to the period of cutting the teeth; then the continuance of the disease is attributed to dentition—this time, without so good reason—and hopes are raised that when the milk-teeth are perfected, the disease will subside. The milk-teeth are all cut, but still the eczema lingers, and then a new light of prophecy beams upon the little patient: when puberty arrives, then certainly the disease will go: but puberty possesses as little of the physician's art as change of food, or completed primary dentition: and so the malady becomes perpetuated. I have seen this picture in life so frequently, that I could not refrain from sketching it.

It is remarkable how trivial an exciting cause may become the origin of this distressing malady. A lady, six weeks after her confinement, travelled by the railroad from London to the sea-coast, carrying with her her infant. She was chilled by the journey, was feverish during the night; her infant was feverish the following day and threw out an eruption of eczema; the child was brought to me some months afterwards. Recently, a neighbour brought me her infant covered with eczema from head to foot. The child was a few months old; in her confinement the mother lost her husband under painful circumstances; the distress caused by this affliction was transmitted to the offspring as an eczema rubrum. How small is the cause of malassimilation in these cases, which may be taken as the type of the whole family; how easily is the assimilative function of infants disturbed; how difficult is it often to be restored.

When cutaneous eruption attacks an infant under these circumstances, it revels in all the typical and modified forms of cutaneous disease. At the same moment, and on the same child, may be seen erythema, lichen, strophulus, eczema, impetigo, pityriasis, and psoriasis: and an obser-

vant nurse seems to take a special delight in pointing out the various diseases which pervade the flesh of the poor little sufferer. In certain parts of the body erythema is apt to prevail; but a broken or cracked state of the skin, with however small a degree of ichorous oozing, must determine the case to be eczema. On the back, lichen is apt to predominate: on the head, in the bends of the joints, and on the pudendum, eczema: on the cheeks and ears, eczema impetiginodes; all on the same skin, and in gross defiance of the orders, genera, and species, of the Plenckio-Willan method of classification.

The predominance of one or other of the typical forms of cutaneous eruption, is determined by the condition and temperament of the infant. The child may present every shade of variation of appearance, from a state difficult to distinguish from complete health, to one in which the little thing is attenuated and shriveled up, and looks like a little old man. In the former extreme, however ruddy and full the child may seem, there is evidence of an existing weakness, in the softness of its muscles; but with that exception, no trace of disorder of constitutional health can be discovered. Next to softness of muscles comes pallor in a slight degree, then an increasing whiteness of the eye, attributable to progressing anæmia: then follows emaciation; the skin shows signs of wrinkles, becomes dry and discoloured, and ultimately sordid. With these, the outward signs of the disease, malassimilation in fact, there is rarely any disturbance, or but little, of the digestive organs; the child takes its food well, and is not particularly restless or fretful. Sometimes the motions are green, sometimes mingled with an excess of mucus, and sometimes white from suspended biliary secretion: but there is nothing beyond the commonest gastro-intestinal derangement, and that in a very insignificant degree.

The eruption usually commences as a patch or blotch of slightly raised pimples; the patch is itchy, is rubbed, increases in size, becomes more inflamed, the cuticle is raised in more or less defined vesicles, which are usually broken by friction; the surface becomes excoriated, somewhat swollen, and pours out an ichorous secretion, varying, from a mere oozing, to an excess which wets through every thing that is applied to it. With the increase of irritation consequent on the excessive secretion and the congestion which gives rise to it, the patch spreads; where the eruption commenced by several blotches, they probably run into one; the ichorous discharge also increases the local disease, by irritating the parts over which it flows. The case up to this time is one of inflammatory eczema, or *eczema rubrum*; the state of *eczema simplex* has hardly existed, and is only to be seen occasionally; but the disease still runs on, its violence increases, and the morbid secretion, from being a transparent and colourless ichor, like water in appearance, becomes slightly opaque (*tinea mucosa*), milky, then yellowish and semi-purulent, and the case is transformed into *eczema impetiginodes*; or the discharge may take on a still more decidedly purulent character, while small pustules are developed on the red and tumefied skin around the patch, and then the case is one of *impetigo*. Thus the *plus* or *minus* of these pathological conditions is irrespective of the cause or essential nature of the disease; in other words, the disease being the same, it may, according to the temperament or constitution of the child, be an erythema verging upon eczema; an eczema rubrum; an eczema verging on impetigo, or eczema impetiginodes; or the pustular element being in excess, it may be an impetigo. Again, as I have before said, whatever the predominating character may be, whether erythema, lichen, eczema, or impetigo, there will always be present, in a greater or less degree, some or the whole of the other forms sprinkled over the body; a simple erythema here, an erythema with strophulus or lichen there; a few scattered vesicles of eczema in a third place, or a few congregated pyodracious pustules of impetigo in a fourth.

In this description of the general characters of eczema infantile, I suppose the eruption to be comparatively undisturbed; but that is rarely the case—the great heat, the