3. Every variety of weather contributed its share of cases; but no exact meteorological register was kept.

4. *Contagion.* Some facts would seem to favour this view; *e.g.*, a boy, aged 19, brought the disease from a village eight miles away, where many cases had occurred, to his home. There had been no prior disease in the house or in the village. A sister and brother had the disease not very severely. Two female children, aged 12 and 10, residing next door in the same block of houses, exhibited the disease severely; the youngest of the two died in two days. All these cases occurred within ten days.—A maid servant, aged 30, had the disease. Her mistress, aged 33, and two children (a boy, aged 6, and a girl of 5), showed the disease within ten days. All were recovering, when another maid servant, aged 25, also took the disease, went home on its appearance, and died within a week.

5. *Scarlatina.* Nothing occurred to prove any connection between the two diseases. Several of the children and nearly all the adults had had scarlatina prior to the occurrence of diphtheria. In one family (not quoted in the list, but personally known to me), three children had diphtheria in turn, and all died from the disease—all of them having two years previously been the subjects of undoubted scarlatina.

6. The *symptoms* attendant on the first appearance of the disease have almost uniformly been those of extreme depression, with consequently more or less of febrile reaction. External swelling of the neck has not usually been observed as a very early symptom; nor is dysphagia early in its appearance or actually necessary in the course of the disease. In one well marked case of a child, aged 4 (dying rapidly with extension of disease to the air-passages), there was never any dysphagia whatever. I am almost disposed to give as the very first local symptom (and the statement is borne out by at least two of the cases in the table) the existence of marked oedema both of the uvula and tonsils—the mucous membrane appearing to be infiltrated with serous exudation, and then studded by the characteristic lymph deposit.

7. *Exudation.* In by far the majority of cases, the exudation was present when the patient was first seen. In the few instances where the progress of the disease could be fairly watched, it came on from the first to the third day after the first complaint of illness.

8. *Microscopic examination* of the effused material in three cases gave no evidence of fungoid growth.

9. *Albumen in the urine* existed in one instance, but was not found in the other instances. Various accidental complications were noted in some of the cases.

10. *Treatment.* I have never resorted to salines, diaphoretics, or, in short, to lowering treatment of any kind; but, on the other hand, have freely given tonics, *e.g.*, tincture of sesquichloride of iron, in doses of ten to twenty minims every four hours; or bark with ammonia; stimulants, as sherry in full quantity, either with or without water, or good beer; nutriment in the shape of beef-tea, essence of meat, arrowroot.

Local Applications. I believe the best to be the strong hydrochloric acid, applied freely, in the first instance, to the parts affected, and repeated if necessary; and followed by the use of gargle of two drachms of hydrochloric acid and an ounce of glycerine with seven ounces of water.

11. *Tracheotomy.* I have no evidence on this point. One child, a girl, aged 4, died from disease invading the air-tubes; but tracheotomy was not permitted.

12. The *mortality* was 2 out of 19, or 10·52 per cent.

13. No *post mortem* examinations were made.

The disease in question is not only worthy of note in its ordinary course, for its danger to life from the exhaustion attendant on it; but also for the varied sequelæ which may show themselves, either as the disease subsides, or even after an apparent though partial recovery.

Sloughing of the cheek (interior), unconnected with the throat-affection by any direct continuity of surface, was observed in a girl aged 12; and subacute inflammation of the mucous membrane of the nose and mouth, lasting for some weeks, in the same child.

Affections in which the nerves principally suffered were very common. Aphonia, lasting some weeks, occurred in a boy aged 7. Paralysis of the portio dura, coming on directly after the subsidence of the disease, on the side most affected, was observed in a female, aged 45. Coincident with this, there were paralysis of the laryngeal and pharyngeal branches of the vagus, and numbness of the hands and feet. Loss of power over the muscles of the palate occurred in a boy aged 7.

Excessive and long continued general debility was seen in

several of the cases; coexistent with it was very complete anaemia in two of the cases—a girl aged 7, and a boy aged 5.

Tubercular disease of the lungs was called into activity in one of the adult cases.

Of skin-diseases, eczema, impetigo, etc., were observed, with formation of numerous and even large abscesses on almost every part of the body.

The whole of these results may be cited as proofs of the deficient power in the system, with, in some of the cases, local manifestations of the constitutional disorder. Whether the affections of the nerves point to any special element in the disease itself, would seem as yet a matter of uncertainty.

C A T A R R H.

By C. HANDFIELD JONES, M.B., F.R.S., Physician to St. Mary's Hospital.

DR. COPLAND says of catarrh, "Although the most common of all diseases, there are few which are less understood, or have called forth a greater diversity of opinion." He defines it as a specific irritation of the nasal and adjacent mucous surface, which may run on to catarrhal bronchitis. In another place, he writes, "It is not pure inflammation, but an irritation of a specific or peculiar kind, attended by slightly increased vascularity, afflux of the circulating fluids, and augmented secretion." He does not sanction the application of the term catarrh to all affections of mucous surfaces attended with sero-mucous discharge. Neither does he assent to the opinion that the "defluxion" is a consequence of the suppression of the cutaneous perspiration arising out of the irritation which the secretion retained in the circulation produces upon the cephalic and pulmonary mucous surfaces." He infers that the causes of catarrh affect primarily the organic nerves supplying the surface principally disordered, and through them, the system generally.

Dr. Watson simply speaks of catarrh as a "moderate degree of inflammation."

Trousseau says, "The catarrhal state and the inflammatory are different, though in some measure allied (*voisins*). The first is distinguished from the second by the circumstance that its accompanying irritations do not terminate by suppuration. Their character is an extreme superficiality, a singular mobility, and diffusion." Again, he speaks of the inflammations even in the severest forms of catarrhal fever as "quasi-inflammatory, false, or bastard." Grave pneumonias, attending such states, are rather an acute stuffing of the lungs (like that of the nasal cavities) than a real pneumonia, which can fairly be attacked by venesection. He recognises very plainly the existence of a nervous element in the disorder which crops out (so to speak) very notably in influenzal catarrh.

Neither Rokitansky nor Virchow seem to recognise catarrhal inflammation as specially different from the same process in other parts.

Catarrhal disorder, for the most part, consists of a general affection, and a local. The latter shows itself in a hyperemic state, more or less verging into downright inflammation of one or more mucous membranes, attended sooner or later with mucous, muco-purulent, or muco-watery exudation. The former consists of various febrile phenomena; it may be of brief duration, or but slightly marked. More particularly we may describe the general disorder as varying from slight chilliness, and subsequent febricitation, with some depression, to a well marked pyrexial condition, with initiatory rigors, followed by rapid, excited pulse, hot skin, thirst, and oppression; or, as is often the case in the present day, the patient chiefly complains of depression and pains in the head and limbs; feels as if beat and bruised, or sore all over, with a cool skin and languid pulse.

The picture of the local affection will of course vary according to its site. Its general features are the above mentioned phlegmasial action, together with certain morbid sensations, and induced movements, often of reflex character. We will describe these more particularly:—As to the *inflammation*. The vessels of the inflamed part are more or less filled and distended with blood, so that the surface presents, in extreme cases, a full red colour. Microscopically examined, it is seen that the congestion most affects the superficial vessels, the terminal capillary loops, which alone may appear filled with blood. Sometimes small extravasations take place, mostly on the surface; but also interstitially. Judging from what we can observe in the conjunctiva, we may believe that the blood in the vessels is not in a state of actual stasis; at least, not during the earlier period of the inflammation. A full venesection will for the time

so relieve the loaded vessels of a scarlet conjunctiva that the membrane resumes its natural paleness. What it does in this case we may suppose it would also effect in a severe bronchitis or pneumonia.

The injection of the blood vessels causes more or less tumefaction of the membrane, which is further increased by interstitial exudation, or in the compound mucous membranes (*e.g.*, gastric or intestinal), by increased formation of epithelium in the follicles. In chronic catarrh the hyperæmia may be more of a dusky red, of a venous tint, or it may lessen so much as to be almost imperceptible, being, in fact, dissipated entirely in the copious exudation.

Exudation. The exudation in the catarrh is chiefly on the free surface of the membrane; submucous effusion belongs essentially to more deeply seated and graver disorders. This distinction seems to me worth establishing as a limitation of the term catarrh; although there is no doubt that the most pure catarrhal inflammation passes into the deep seated by insensible grades. Thus, I would not name a case of purulent ophthalmia, with notable chemosis, conjunctival catarrh; nor a case of œdema of the glottis, laryngeal catarrh. It is to be borne in mind that submucous exudation is much more likely to occur when the areolar tissue in that situation is abundant or lax. Another limitation of catarrh is afforded by the change of the exudation to fibrinous matter, constituting the false membranes of croup and diphtheritis. We shall not attempt to discuss here the special causes of this change; but only point it out as a state manifestly different from that of catarrh.

The truly catarrhal exudations consist of blood-plasma, modified by its transition through a homogeneous membrane, and by the influence of the superjacent epithelial cells. The most essential constituent is the sticky, glutinous, tenacious substance which is the product of all mucous membranes, and gives its name to them. There can scarce be any doubt that this is modified albumen, though the chemist cannot reverse the process of vital chemistry and reconvert it, or obtain albumen from it. It contains less nitrogen (12.82: 15.67) and more oxygen (28.11: 22.36) than albumen, and is rich in the chlorides of alkalies. In part at least it seems probable that it is produced by the cells of the surface from which it is effused, which elaborate it in their interior. According to another view, the mucous stuff is formed from the blood-plasma as a complementary product to the epithelial cells. Catarrhal exudation always contains albumen mingled with the mucous fluid, and the same is the case with analogous productions, as synovia, and the contents of colloid cysts. The corpuscles present in morbid mucus are for the most part identical with those of pus, they are often, however, mingled with various forms of immature epithelial cells, or cells with granulous contents, or actual glomeruli. Blood-corpuscles may also be present, even in notable quantity, and sometimes minute fibrinous films. It is very worthy of notice that in the severer grades of inflammation catarrhal mucus is eminently adhesive, so much so that it can scarcely be shaken out of the porringer containing it. In this state it is also much more transparent and more scanty than any subsequent period. The transparency in a great measure depends on the absence or paucity of the multitudinous corpuscles which are crowded together in the more opaque varieties. In nasal catarrh the exudation at the outset is often remarkably watery and copious. It is not clear upon what this peculiarity depends; but it may be surmised that the large venous plexuses which underlie the mucous membrane are not unconcerned in it.

While this watery flow continues it would seem that there is rather a straining off of fluid from the blood vessels than a secretion. The latter faculty, implying an active condition of the cells covering the surface, is not restored until the flux begins to decline. The instances of the Malpighian renal tufts, and of the pulmonic air cell plexuses afford proof that simple capillary membranes in their normal state may prevent effusion of all but the saline and watery parts of the blood. When the normal filtering power is deranged by continuance of the hyperæmic stress, albuminous fluid next escapes; or even fibrinous and red corpuscles. The quantity of aqueous fluid which escapes from the renal capillaries varies very greatly we know with the state of the nervous force, being often remarkably increased when this is lowered. These remarks elucidate in some measure the peculiarity of the nasal catarrhal exudation first adverted to. They show that the filtration, taking place through homogenous capillary membranes, may vary greatly, according to circumstances with which we are as yet insufficiently acquainted; one of which, however, of prime importance, is evidently the degree of strain to which they are subjected by

the blood current. The mucus which lines the pyloric region of the stomach in gastric catarrh is often very closely adherent and tenacious, and is peculiar in not containing, so far as I have seen, any other corpuscles than detached epithelial particles. The mucus passed in dysenteric purging contains, at least in the earlier periods, numerous nuclear corpuscles, probably derived from the Lieberkühn follicles; but not pus globules. These, no doubt, are present at a later period, and in the severer cases. The mucus from the bladder and urinary passages easily becomes quite puriform, and always contains multitudes of pus corpuscles. The uterine mucus tends to be puriform in about the same degree as the bronchial. It is always notably alkaline. The vaginal secretion is remarkably acid, and when excessive in quantity is of a much more watery quality.

However, as Dr. Tyler Smith describes it, the cervical uterine "secretion" may be so profuse and watery that the traces of visciditv are nearly lost.

It is tolerably constant to find the catarrhal exudations in chronic cases attended with much debility and want of tone—profuse and watery. Perhaps the best instance of this is the flux called bronchorrhœa. Pyrosis, in my opinion, is a very analogous discharge from the gastric mucous surface. The pain preceding the gush, the copiousness of the discharge, and the absence of the disorder in mercurial and neuralgic salivation prevent me from assenting to the opinion that pyrosis results from deglutition of much saliva. In both these instances the capillary walls must be in a different state, as regards filtering power, from that existing in chronic bronchitis, with profuse muco-purulent secretion; or gastric catarrh, with tenacious mucous exudation. The state conditioning the then passive flux must be similar to that of the renal tufts in low nerve disorder.

Sensation. The sensations, the dysæsthesiæ, attending catarrh, are familiar to all. In the eye there is that of grit between the lids, with more or less stiffness and redness. In nasal catarrh, there are stuffing, heat, dryness, impairment of smell, perhaps gravative headache: in bronchial catarrh, sense of soreness or rawness, heat, and tightness in the chest: in gastric there is tenderness on pressure, and pain at the epigastrium, or between the shoulders, thirst, loss of appetite, perhaps nausea: in intestinal, abdominal uneasiness and griping: in vesical, frequent calls to micturition, "shooting, throbbing pains in the region of the bladder," increased by pressure, dysuria: in urethral, scalding in micturition, heat, pain and tenderness in the part: in uterine catarrh, pelvic uneasiness, sense of weight and bearing down, pain and aching in the loins.

Induced Movements. The induced movements are as follows: In the eye, more or less spasmodic contraction of the orbicularis palpebrarum: in the nasal cavities, sneezing, or voluntary expulsive efforts: in bronchial catarrh, cough: in gastric, vomiting: in intestinal, diarrhœa, or hurried peristaltic action: in vesical, frequent micturition.

Fever. Febrile movement often forms a part of catarrh. In the majority of cases, at the present day, it is either from the outset decidedly asthenic, or tends speedily in that direction; but it may often be of an opposite character. In influenzal catarrh the asthenic quality is strongly marked. The state of the secretions has nothing very constant in catarrh. The perspiration may be temporarily suppressed or not. In the first case its restoration in the freest manner by no means insures improvement. The urine may be high coloured and turbid if there is notable pyrexia; often it is apparently normal. The bowels are usually relaxed if the intestinal tract is affected.

Causes. The causes of catarrh are certainly manifold. Not to insist on the fact that mechanical irritation will produce catarrhal inflammation of a mucous surface, there is good ground for the statement that the disorder in its general form, and with local inflammation, may be occasioned by atmospheric influences or by exposure to damp cold after the prolonged action of relaxing heat.

Dr. Watson states that errors in diet in some persons, occasioning dyspepsia, always produce catarrh. The gouty poison will give rise to bronchial catarrh, as in a case recently under my care. So do the peculiar emanations from hay in some persons of special idiosyncrasy. The poison of measles produces commonly well marked catarrhal effects. Again, as regards nasal catarrh at least, I am satisfied it may come on without the action of cold or epidemic influence, or any discoverable cause. The disorder seems to originate *per se*. The following points seem to be well ascertained. Mere cold, however severe, does not induce catarrh, in fact, when climatic it seems to have the opposite effect, and to diminish the liability

of the system to be affected in this way. The beneficial effect of the cold shower, or douche bath, is an instance of the same kind.* The high temperature of a tropical climate is no protection against catarrh; colds and coughs, though not so common as in England, are by no means unfrequent in India. This I state on the authority of several residents whom I questioned as to the circumstance. I had lately under my care an elderly gentleman, with chronic bronchial and nasal catarrh which commenced in India. Diarrhoea, one of the most frequent of tropical diseases, in most cases is simply intestinal catarrh.† So far from affording immunity, prolonged heat evidently predisposes to catarrh by enfeebling the system, just as cold has the opposite effect. Bakers, and other workmen whose occupations expose them to vicissitudes of heat and cold, are more than usually prone to catarrh.

Dr. Copland remarks, "It should not be overlooked also, that sudden change from a low to a high temperature, or from a very dry to a very moist air, and even the being more than commonly over heated, without any very apparent chill or exposure to cold subsequently in any form, will often produce catarrh. This is especially the case if the exposure to warmth be sudden, after an impression of cold of some continuance, as the coming into an over-heated apartment out of a cold and moist atmosphere—the instantaneous transmission from a raw air of about 32° to a dry air of upwards of 70°. The production of catarrh by peculiar atmospheric influences (influenzal), independent of the effect of temperature, is a striking phenomenon. It seems natural to attribute it to the reception of some aerial poison into the system, though whether this be a material thing, as a deleterious gas, or whether it be an influence, as, e.g., some peculiar electric state which affects the nervous system in a perturbing manner, can scarcely be determined. The very sudden, wide spread invasion, and the characteristic protraction of influenza seem more like the effects of the latter kind of cause than the former.

Pathological Relations. The diseases with which catarrh appears to have decided affinity are rheumatism, malarious disorder, and some superficial cutaneous eruptions. Dr. Copland says, "Of the more pure complications of rheumatism, none are so frequent as those with catarrhal fever, or simple catarrh." Again, "Rheumatism generally proceeds from the same or very similar causes to those which produce catarrh; they are both also often present at the same time and in the same person, and the epidemic prevalence of both is not uncommon." I have recently observed the occurrence of well-marked herpes labialis in connection with pretty sharp rheumatic pains, no trace of catarrh existing.

This eruption Dr. Salter considers as positively conclusive of the presence of catarrh. I do not entirely agree with him, but there is no doubt that it very often does accompany the catarrhal state. Its occurrence in rheumatism is therefore an indication of affinity. It is generally in the less severe and marked instances of either disorder that we observe the co-existence of rheumatism and catarrh; for when one is fully developed it absorbs, so to speak, the other. This is what might have been expected in diseases alike in essence and mode of action, but differing in locality.

The affinity between catarrh and malarious disorder is indicated by the following circumstances. Dysentery and diarrhoea, both unquestionably common results of malaria, are as often almost the consequences of the causes of catarrh. (Vide the case of the army at Dettingen, quoted by Dr. Watson from Sir J. Pringle, vol. ii, p. 474.) Influenza has been observed by Huxham, Cleghorn, Petrie, and H. Greenhow, to show a decided relation to remittent or intermittent fever. The first named authority states that influenza frequently, after two or three days, ran into a quotidian, or tertian. Dr. Greenhow says, "The influenza of 1847 very often assumed an aguish character; exacerbations of the disease, preceded by anomalous pains, cough, rigors, heat and sweating, with loaded urine, recurring periodically, at tertian or quartan intervals." The pains which accompany the invasion of catarrh are truly neuralgic, indicating the especial concernment of the nervous system. The same is the import of the remarkable prostration of influenza, which in rare cases assumes the form of actual paralysis, as in a case which I have recorded in this JOURNAL,

* In a paper on Catarrh (*vide Lancet*, July 4th and 11th, 1857), I have quoted some convincing proofs of this.

† In the annual return of the European General Hospital at Bombay, out of 1093 cases of all kinds, medical and surgical, there are 7 cases of acute catarrh, 7 of asthma, 13 of acute and 8 of chronic bronchitis, 53 of acute dysentery and 7 of chronic, 37 of acute and 28 of chronic rheumatism, and 87 of diarrhoea. (*Transactions of the Medical and Physical Society of Bombay*, 1855.)

April 16th, 1859. As there is much evidence to show that malarious disorder prominently affects the nervous system, the two maladies in this respect manifest a resemblance to each other.* In ague, and in catarrhal fever there occur general febrile disorder and local hyperæmia, the latter being sometimes wanting, or very slight; and not in any case originating the former. Again, dysentery, the result of malaria, is often a substantive inflammation with excited pyrexia, just as bronchitis from catarrhal influence is with its fever.

Between catarrhal inflammations and such skin eruptions as diffuse eczema, there is a manifest resemblance. The acute or sthenic stages, and the chronic or asthenic, are quite analogous. In both there is either inflammation of, or flux from, a tegumentary surface; but in one case it is from an inner, in the other from an outer. Eczema is just such an inflammation of the skin, as catarrh is of a mucous membrane; both affections are superficial and spreading, attended with free exudation; and differ from phlegmonous or croupous inflammation. In respect of its causes, there is not often any observable connection between eczema and catarrh; but another vesicular disorder, herpes, is, as already observed, a very frequent attendant upon catarrh.

Pathology. The pathology of catarrh is of much interest to inquire into, not only on account of the frequency of the disease, its occasional severity, and its relation to other diseases, but because the advance of scientific knowledge seems to render it possible to gain a clearer and better understanding of its various phenomena. Let us take an ordinary case for examination.

A person is exposed to damp cold, and soon after begins to feel shivery and depressed, to have tightness of chest, some dyspnoea, cough; in short, has a bronchial catarrh, which proceeds, with a certain amount of pyrexia in the earlier periods, and in the latter with gradual diminution and cessation of the mucous exudation. In such a case, a morbid impression has been made on the general cutaneous sensory surface; perhaps particularly on the anterior thoracic part—the chest, as popularly termed. This, it may be presumed, is communicated through the spinal cord to the sympathetic ganglia, from whence the vaso-motor nerves of the tracheal and bronchial arteries originate. The morbid impression causes more or less paralysis of these ganglia, or, through them, of the vaso-motor nerves derived from them. A state is thus set up in the membrane lining the air passages characterised by hyperæmia, heat, and hurried nutritive actions. (Vide Bernard's description, *Leçons sur le Système nerveux, grand sympath.* vol. ii.) This lighting upon a predisposed part, gives rise to inflammation and exudation. If the individual continues to be exposed to the cause, viz., damp and chill air, the morbid impression is intensified, and the disease aggravated. Under these circumstances it may extend much beyond its original seat, and pass into catarrhal pneumonia. The same effect, i.e., the extensive spread of the disease, may take place if the morbid impression meet with a high state of predisposition in the bronchial lining membrane and that of the pulmonary air-cells. We cannot explain in what this predisposition consists. We know it means a proneness in the nutritive processes to be deranged from their normal state. That it really exists in certain persons, and at certain periods, we are sure from the very different facility with which the skin and other parts are affected by various agents in different individuals. One skin can with difficulty be inflamed by a mustard poultice or blister, another is very easily. One person's stomach is irritated by a minute dose of arsenic, another's can bear a much larger. A low degree of power in the vaso-motor nerves, and an abundance of liquor sanguinis in the blood vessels, are probably also elements of the predisposition. The kind of inflammation set up in the part will depend on the state of the tissue. If its excitability be low and weak, the inflammation will be asthenic; if it be considerable and vivacious, the morbid action will be sthenic. In the former case nerve tonics may be used with advantage, in the latter they will only increase the tissue irritation.

Pyrexia may be set up either by the original morbid impression, involving all the vaso-motor nerves of the system in a greater or less degree, or by the irritation of the inflamed part being reflected upon the same nerves and paralysing them in a way of inhibition. Increased action of the heart may be produced partly by the increased heat of the blood acting as a

* The replacement of neuralgia by ague and *vice versa*, the curative effect of mental excitants, and of quinine, whose action is undoubtedly chiefly on the nervous system, the occurrence of hemiplegia (temporary) in connection with ague attacks, as well as of cardiac paralysis, forcibly declare the especial implication of the nervous system by the malarious poison.

stimulant to its muscular tissue, partly by the paresis of its vaso-motor nerves occasioning hyperæsthesia, and hurry of its nutrition. The chilliness and shivering of the initiatory period point to a temporary excitement of the vaso-motor nerves, soon succeeded by exhaustion and debility; this excitement will induce contraction of the superficial arteries and a feeling of cold, or rather, actual lowering of temperature. If, however, the catarrhal phenomena result from the reception of a poison into the blood, or the operation of some influence which may be supposed to act at once on the whole nervous system, the process will be somewhat different. The vaso-motor nerves will be directly enfeebled, as well as the cardiac and musculo-motor, though in a less degree. The paresis of the cardiac may be sufficient to cause syncope, and that of the musculo-motor actual, if not complete, paralysis. In analogous cases it appears as if the encephalon was affected, and hemiplegia, or so-called simple apoplexy occurs. The sensory nerves are tormented with neuralgic pains, they seeming in most cases to express their enfeeblement by the cry of pain. The tissues generally feel the enfeebling influence, and the phlegmasiæ that are produced are mostly, if not always, asthenic.

This asthenic state depends upon the debility of the tissues of the affected part, which are so prostrated as to be unable to react powerfully under the hyperæmia and disturbance of their normal nutrition.

Another view that has been taken as to the pathology of cold-excited catarrh is, that the blood is poisoned by the retention of some element of suppressed perspiration. This seems to me refuted by the circumstance that the most complete diaphoresis is no certain remedy for such catarrh.

SPECIAL REMARKS. In common ophthalmic catarrh the effect of cold as an exciting cause is very marked. The popular term "a blast of cold in the eye," is probably quite correct, describing as it does a sudden morbid impression producing effects of some continuance. This impression cannot, we may be almost sure, directly touch the mucous surface that becomes inflamed, as that remains constantly applied against the globe of the eye. It is probably made on the exposed part of the membrane.

In ordinary nasal catarrh I have been able very positively to observe some circumstances influencing the flux. Fatigue has had the most marked effect in increasing the flow, rest and food in restraining it. Free ingestion of liquids did not increase it. The cold of a mountain top lessened the flow decidedly. In the outset the thin flux sometimes pours from one nostril alone; after a day or two it has ceased in the first affected, and commenced in the other. A dry, troublesome, tickling, nocturnal cough, an evident neurosis, sometimes terminates nasal catarrh.

Tonsillitis is notoriously a common result of a catarrhal stroke. It requires notice that the swelling of the part is chiefly dependent on submucous hyperæmia and exudation. The tonsil is composed, not so much of mucous follicles as of a dense layer of nuclei, holding the same relation to the buccal and faucial lining as the solitary glands and Peyer's patches do to that of the ileum. The increase and hypertrophy of this submucous deposit produces the enlargement.

In *Pharyngeal Catarrh* the follicles are apt to be affected, and to secrete tenacious muco-pus. This morbid secretion often becomes habitual, especially in tipplers.

Laryngeal and Tracheal Catarrh are very common, and give rise to severe irritating cough, which it is important not to mistake for bronchial in respect to the application of local remedies. The alteration of voice is, of course, the most diagnostic sign of the former.

Bronchial Catarrh, and Catarrhal Pneumonia have their proper auscultatory signs, which it is needless to detail. In Walshe's or Watson's works the student will find all he needs. Only I would repeat the cautions: (1) Not to estimate the severity and peril of an attack of catarrh by the loudness of the râles (the most perilous cases may produce very little râle, because the tubes are to a great extent clogged up); but by the extinction of the respiratory murmur and the blueness of the lips. (2) Never to trust merely to the auscultatory phenomena to the neglect of the general. The latter may be alone capable of giving information as to the import of the former.

Gastric Catarrh is infinitely common, quite as much so as bronchial, and proceeds from the same general causes, as well as from the irritation of indigestible food, and the like. It produces in many cases the tough, tenacious mucus before noticed, which of itself must prove a mechanical hindrance to digestion by enveloping the food and keeping it from the access of the gastric juice. It chiefly affects the papillæ and the superficial

parts of the gastric tubules; but in severer cases may probably extend more deeply, and produce alteration and separation of the glandular epithelium. The vessels that run between and surround the orifice of the tubules on the surface are always more congested; and from their rupture blood may be extravasated and mingled with the exudation. Slaty discoloration of the gastric mucous membrane proceeds also from the escape of blood, or of its colouring matter, into the tubes, or their interstices.

With regard to *duodenal catarrh*, I must join heartily with Bamberger in believing that it is a most difficult thing, in the vast majority of cases, to diagnose its existence. It is, however, always worth considering whether, in cases of icterus, the outflow of bile may not be prevented by catarrh of the duodenum implicating the bile-duct, and closing its narrow orifice by swelling of the membrane. Local tenderness on pressure is almost the only sign that can be esteemed of much value to determine this question, except the result of tentative treatment, as applying a blister to the right hypochondrium.

The diagnosis between *catarrh of the small* and of the *large intestine*, is stated by Bamberger as resting on the following points:—1. The pain in the case of the large intestine is attended with tormina and tenesmus, and is more severe than it is when the small intestine suffers: the pain then is duller, or may be quite absent. 2. The pain on external pressure is felt in the regions occupied by the colon when the large intestine is affected—i. e., the lateral and upper; in catarrh of the small intestine, it is felt in the middle and lower. 3. Vomiting occurs much more often and severely when the small intestine is affected, than when the large is. 4. General nutrition and the health suffer more in disease of the small than of the large intestine (*cæteris paribus*). 5. Collection of exuded fluid in the small intestine causes dulness and gurgling murmurs in the mid-abdominal region; in the colon, the same signs will be found in the regions it occupies. 6. In catarrh of the large intestine, the fluid stools have more the fecal character, or consist in great part of glassy mucus in small or large lumps, or of puriform fluid. The admixture of pure or recognisable blood generally indicates the occurrence of dysentery. If the upper part of the small intestine alone is affected, diarrhœa may be quite absent; if the disease extends to the ileum, the stools consist of a yellowish green, or greyish yellow green, flocculent fluid, depositing a sediment on standing.

In *vesical catarrh*, the influence of cold as an exciting cause is acknowledged by Mr. Coulson; it is frequently produced by mechanical causes, as the irritation of a calculus; and by certain drugs, as cantharides; and is also a common consequence of injury to the spinal cord, producing paralysis of its nerves. In the latter case, the inflammation is no doubt the result of vaso-motor nerve paralysis, reacted upon and aggravated by the urine, rendered alkaline by the effused mucus. The result of loss of nervous power in this instance is worth noting, with respect to the general pathology of catarrh. It seems reasonable to conclude that cold must induce a similar state of the vaso-motor nerves as results from injury to the spinal cord. Mechanical irritation, and the action of irritant poisons, as cantharides, seem, on the other hand, to operate directly on the tissue.

Urethritis, in its milder grades, may be termed urethral catarrh; it passes beyond this stage when submucous exudation takes place, as in the origination of stricture. In the male, urethral catarrh is rare, except as the result of infection. I have, however, seen a similar affection, though slight, and confined to the mucous lining of the prepuce (gonorr. ext.), in a healthy male child, where all other causes were positively excluded, except such as occasion common catarrh in other parts. The inflammation and puriform discharge have occurred on several occasions. In female children, as is well known, a catarrhal inflammation of the vulva and adjacent vagina is far from uncommon, and is quite independent of any specific disease. Gouty poison in the blood may excite urethral catarrh, just as it does bronchial.

Uterine catarrh, in the majority of cases, depends on the common influences causative of the general disorder, rather than upon any local irritation. Its treatment assimilates it wholly to catarrh of other parts.

TREATMENT. The first object in dealing with a case of catarrh, supposing it to be of sthenic character, is to arrest the morbid excitement, and stay the local attraction of blood. For this purpose, we resort to various means which seem chiefly to operate on the tissue of the part affected. One of the first to be noticed is local blood-letting, which certainly has no small power as a tissue-sedative. This is manifested not only when there is a direct vascular communication with the inflamed

part, but when a serous cavity intervenes, and widely separates the vascular plexuses of the cutaneous surface and the subjacent mucous. It is beyond all question that leeches to the epigastrium relieve an inflamed gastric membrane; but surely no one will imagine that this relief depends on any withdrawal of blood from the engorged gastric capillaries. General bleeding may be of service, if the heart's action, as evidenced by a firm pulse, is over-powerful; but it is of no much avail as a tissue-sedative alone. In some instances, we have drugs which exert a decided sedative action on the inflamed membrane. In inflammation of the air-passages, antimony or ipecacuan; in enteric catarrh, hydrargyrum cum cretâ and Dover's powder; in dysentery, ipecacuan; in tonsillitis, an emetic,—are all means of well proved efficacy. It must be confessed this list is a short one, and we should be glad to be able to enlarge and amend it. We have no reliable tissue-sedative for ophthalmic or nasal catarrh, for vesical, or urethral. In the former, we have a valuable substitute for internal remedies in the application of nitrate of silver collyria, of various degrees of strength. The arrest of severe purulent conjunctival inflammation by the application of solid lunar caustic, or Mr. Guthrie's ten-grain ointment, is a striking therapeutical fact. It is a clear proof of the vital power of an inflamed tissue being modified and altered by a means directly addressed to it. Vaginal catarrh may be dealt with sometimes in the same way; and so, according to some, may urethral. Counterirritation by blisters, turpentine stupes, mustard poultices, etc., is a valuable means of tranquillising the morbid excitement in the inflamed tissue; but it does not find its opportunity until the force of the disease is somewhat broken, or unless it be of semi-asthenic character from the outset, or is not yet fully established. The turpentine stupe is an especial favourite with me, and may be used with advantage much earlier than a blister. It may probably act to some extent as a beneficial stimulus to the cutaneous nerves, propagating through them an impression to the sympathetic ganglia and vaso-motor nerves, which obviates and nullifies the original morbid impression. A blister should seldom be applied until all fever has nearly subsided. Perpetual poultices or fomentations are often of great service in the treatment of the earlier stages of catarrh. The proceeding of incubation, as it is termed by M. Guyot (which consists in exposing diseased parts, ulcerated or inflamed, to a dry (?) heat of 96° or thereabouts), is described by him as having the effect of dissipating pain, inflammatory redness, and swelling. Heat, in fact, seems to have a general relaxing and sedative effect; it causes very positively relaxation of the contractile tissues of arteries, and it seems to act in an analogous manner upon other parts. Cold has the opposite effect, inducing reaction and elevated temperature; while heat leaves the part or the system with a diminished caloric-producing faculty.

For the purpose of allaying tissue-irritation, alkalies with their vegetable salts, as well as the so-called neutral, are remedies worthy of notice. They seem to act beneficially on all mucous surfaces when inflamed, excepting the intestinal canal, which the neutral and vegetable salts are apt to irritate. When freely administered, these agents have also a decided effect in calming the action of the heart, and in promoting the excretion of retained matters. Hydrocyanic acid is often a valuable adjunct to these remedies, and acts in the same direction. Opium, or other sedatives, may often be conjoined with antimony, mercury, or ipecacuan; and their action thereby be rendered much more safe and easy. Independently of this auxiliary action, opium in some cases by itself appears to exert power as a tissue-sedative. This effect may be produced through the nerves, but it is certainly not confined to them.

If the state of the affected part or system be one of depression, rather than of excitement—i. e., if the state be asthenic—we have recourse to various means of more or less stimulating or toning quality. These act beneficially by rousing the languid vaso-motor nerves, contracting the arteries, and so both diminishing the afflux of blood to hyperæmic parts, and quieting the hurried nutritive actions that ensue when the controlling nerves are paralysed. At the same time, they raise the often impaired power of the cerebro-spinal nervous system. Of this class we may enumerate carbonate of ammonia, the ethers, valerian, arnica, sumbul, as especially acting on the cerebro-spinal nervous system as stimulants or excitants, rather than as tonics. They may be usefully combined with salines where we wish to have but a moderate effect. On the other hand, when the signs of depression are predominant, we give them full play. Quinine, strychnine, nitric acid, the sulphates of iron and copper, the muriate and penitrate of iron, are mostly

appropriate to cases of asthenic mucous or muco-serous flux, and may be often advantageously associated with a little opium. In cases not admitting of the use of these latter tonics, the balsams, squill, turpentine, copaiba, are often very useful. Their mode of action seems to be of the same kind as that of the stronger tonics, but they are better adapted for peculiar states and special tissues. Thus copaiba has eminently an influence over the bladder, urethra, and rectum; balsam of Peru, squill, and Canadian balsam, on the bronchial membrane. A remedy of the same class is ergot, which Dr. Churchill has found efficient in some cases of uterine catarrh. The especial advantage of all these is, that they suitably stimulate and tone the delicate nervous tissue in cases where, from one cause or other, the stronger tonics would irritate, and so fail of their desired effect.

Astringents are appropriate to all cases where the exudation is mainly of the nature of passive flux; i. e., where the tissues are in a languid state, not prone to irritation. The action of astringents is probably mainly expended on the capillaries, rendering their delicate membrane firmer, and less permeable by fluids or corpuscles. I am aware that I am here touching on the oft debated question of diapedesis; but I am supported in the view, that red corpuscles do pass through the walls of capillaries and small arteries and veins, by Virchow's positive testimony, based upon actual observation. He states that openings may be seen in vessels through which red globules escape in single file, and which again close completely (*Handbuch der Speciellen Pathologie und Therapie*, p. 231); and, at p. 237, he refers the escape of the menstrual blood to a diminished cohesion of the vessels of the mucous surface. This diminished cohesion, which would of course favour the escape of mucous or serous fluid, is just what astringents correct, whether they be locally applied, or carried in the blood-currents. It is clear that it will be judicious practice in any case where the large open quality of the pulse conveys the information that the arteries are uncontracted, to administer at the same time a nerve tonic, so as, if possible, to diminish the local supply of blood, and so relieve the weak capillaries from undue strain or pressure of the current. Gallic acid, tannin and all the vegetable preparations containing it, alum, and iron alum, are our best astringent remedies. The influence of cold, also, ought not to be omitted, though this is exerted rather on the small arteries than on the capillaries. When locally applied, it will produce, of course, a local effect; but it may also operate on remote parts by the impression it makes on a sensory surface being propagated as a stimulus to the arteries of the seat of internal disorder through the medium of the nervous centres and connected vaso-motor nerves.

The foregoing exposition is presented rather as an attempt to colligate familiar facts than to educe new. I believe in the grouping of these several morbid conditions. I have strictly followed natural affinities, and it appears to me some gain to be able to regard a number of ordinary phenomena from the same point of view. He must, I hold, have a better comprehension of individual cases of disease who contemplates them always as far as possible with reference to the great pathological families to which they belong, and to their mutual affinities, than he whose attention is taken up with hair-splitting distinctions which Nature for the most part entirely ignores.

STRICTURE OF THE ŒSOPHAGUS.

By ROBERT BATTY, Esq., Liverpool.

THE subjoined case of œsophageal stricture may prove interesting more from the manner of death, and symptoms preceding it, than from any additional light which it throws upon the usual treatment and pathology of such cases. It occurred in the private practice of Mr. Bickersteth (through whose kindness I am enabled to give, briefly, the history of the patient), and was brought under my notice only two days previous to the fatal termination.

Kate W., aged 19, of apparently vigorous constitution, consulted Mr. Bickersteth about two years ago, complaining of symptoms of dysphagia. A bougie was passed with ease into the stomach. Her general health was good, and although she was not apparently of an hysterical temperament, yet her state seemed to indicate the existence of spasmodic affection of the tube dependent upon some obscure emotional cause rather than that of any organic change of structure. She herself persisted that food in a solid form did not reach the stomach; but as the bougie passed without difficulty and the food did not return, this was regarded as a morbid impression.