Original Communications.

ON THE NATURE AND ORIGIN OF RENAL DISEASE.

By J. RUSSELL, M.D., Lecturer on Pathology at Sydenham College, Birmingham; and formerly Senior Physician to the Birmingham General Dispensary.

[Read before the Birmingham and Midland Counties Branch, April 9th, 1857.]

In laying before this Branch a few remarks on the nature and origin of renal disease, I limit myself entirely to that large and important class of diseases of the kidney hitherto known under the collective name of Bright's disease; characterised as a group, though not of necessity as individuals, by the presence of albumen in the urine, by a serious diminution in the daily excretion of urinary salts, and by the passage from the kidney of tubules of certain peculiar casts; together with a tendency to the effusion of anasarcaous fluid into the cellular tissue of the limbs and trunk.

Among the many interesting questions which the phenomena of this very heterogeneous class of diseases suggest, obviously the most important is that which relates to their nature and origin; since upon the answer which is given to this question must depend the views we entertain as to treatment, both curative and preventive. I therefore make no apology for introducing a subject, which, but for its practical application, may have rather an abstract appearance.

In discussing this subject, the first inquiry must relate to those morbid changes in the kidneys which constitute the disease. These have been recently described by Dr. George Johnson, in his work on *Diseases of the Kidney*. The primary changes vary in the different forms of the disease: in one form, the epithelial cells, which perform the function of secretion, are shrivelled and broken up, and the uriniferous tubules are, in consequence, either shrunk and wasted, or, being bared of their epithelial lining, are dilated to various degrees: in another form of the disease, the place of the cells is occupied by a homogeneous material, apparently of a fibrinous character: and in a third, to which the name of Bright's disease is sometimes restricted, these epithelial cells either remain loaded with fat, or, having been ruptured by excessive distension, fill the tubules with their oily contents. These primary changes are speedily followed by others, both in the tissues of the kidney itself, and in other organs of the body; but such changes, being of a secondary nature, do not concern our present inquiry.

The structural changes I have thus briefly indicated have one important character in common, which has a special bearing upon the opinion I shall suggest as to their nature and origin; viz., that they all commence in the *secreting instruments* of the kidney, the glandular epithelial cells which line the convoluted tubules; it is the *cortical portion* of the organ which receives the stress of the disease; the straight tubes which constitute the medullary portion, and which are little besides ducts, are affected to a much less extent, and often escape altogether. Further, it is important to observe that the changes in question are degenerative in their character, and in this respect are directly opposed to that creation of additional secreting tissue which constitutes true hypertrophy of the kidney, and which occurs in the course of the natural effort made by a healthy kidney to do double work, when its fellow is disabled by injury or disease; they thus satisfactorily agree with the prominent symptom of the disease—namely, the lessened amount of salts secreted in the urine.

The explanation of these various structural changes, which is given by Dr. Johnson, and to which I now wish to direct the attention of the Branch, is the following. The morbid changes take their origin in alteration of function in the parts diseased, viz., in the epithelial cells. In consequence of exposure to unhealthy influences of various kinds, the nutrition of the blood becomes depraved, and that fluid becomes loaded with unhealthy matters, which must be eliminated from the body; the labour of effecting this is thrown, in great measure, upon the kidneys, and materials of an unhealthy character are thus being continually brought for excretion to the cells of those organs: consequently, the function of the cells becomes per-

verted; their natural food, the excreta, is changed. As a necessary result their nutrition becomes impaired, and morbid alterations of structure are induced. Such alterations may be effected gradually, almost insensibly, if the corresponding change of function be itself induced gradually; on the other hand, if the work of the organ be suddenly changed, the alteration of kidney structure will be attended by a relative commotion in the organ itself, and secondarily, in the system at large. By what singular modification in the operation of the causes the different forms of the disease are produced, we are just as ignorant as we are of the causes which determine fatty degeneration in an organ at one time, wasting at another, and calcareous transformation at another. Much, no doubt, is due to the original constitution of the patient. Thus, one variety is most frequently met with in the strumous constitution.

Dr. Johnson's explanation of the disease seems to me to be supported by the great pathological law, that in every organ and tissue nutrition follows function, is related to it, and regulated by it; and I think it derives support, not only from the history of the patient's life and habits, and from the nature of the influences which seem to stand to the disease in the relation of cause, but also from the analogy of corresponding changes in other organs of the body. I shall beg leave to submit a few arguments of each class.

First, I will refer to the arguments derived from the analogy of other organs. Until lately, inflammation has been constantly referred to as the one important source in which most of the changes of structure, with which we are familiar, take their rise: and we now most naturally look for traces of this process in explaining any disease which falls under our observation. Late researches, however, have revealed to us a very large number of organic changes, which are entirely independent of inflammation. The large class of hypertrophies, properly so called—those, namely, in which the enlargement is entirely due to increase in the natural tissue of an organ—exhibit to us instances of large increase of dimension and weight, without any inflammation having been concerned in the process. On the other hand, in the class of atrophies, there is equal change in the opposite direction; and, although inflammation, like any other influence which depraves nutrition, may produce atrophy, yet, in a vast majority of instances, the change has been entirely independent of that process.

But it may be said, there is no alteration of structure in the instances cited; nothing but augmented or diminished bulk. Look, then, at the still larger class of degenerations; here are found muscular, arterial, cartilaginous, osseous, fibrous, and other tissues transmuted entirely into fat or earthy matter, or withered into mere fibro-cellular tissue, and yet inflammation has had no part in the matter. Look again at cancerous and tubercular formations, and at the non-malignant tumours; what numberless examples of extensive structural change, independent of inflammation, do they afford. And yet there is hardly one of the changes to which I have alluded which has not afforded, at one time or other, a battle-field for the question of how far inflammation has been concerned in its production:—the contest relating to the origin of tubercle, and of the atrophy of arteries, whether or not it is connected with inflammation, is one of our own day.

Since we have discarded the inflammatory origin of these and similar changes, it is a necessary question, in what do they originate? where is their starting point? I think it will not be doubted that in a large majority of the instances I have quoted, the first step in the process of change is connected with the *function* of the organ itself. In the hypertrophied heart, the first in the series of changes is a demand made upon the organ for increased exertion, in consequence of some obstacle retarding the onward current of the blood; just as conception takes place before the natural increase of the womb in pregnancy commences, thereby creating fresh demand on the nutrition of that organ. On the other hand, in the old woman we find the ovaries atrophied, in the palsied limb we find the muscles wasted or degenerated; here again the first step has been a change in the function of the organ affected, an abrogation or serious diminution of that particular function, just as in the womb lessened requirement, consequent upon delivery, precedes the shrinking which follows that event.

Perhaps the most striking exemplification of this waiting of nutrition upon function, is afforded by the changes which are gradually effected in the tissue of every organ during the period of decline from manhood to old age; as the time for mental and bodily activity passes away, the standard of nutrition in the entire body is lowered, and every form of degenerative change may be met with in the various tissues of the

body. The great tendency of our present pathology to explain change of structure by previous alteration of function is exemplified in the hypothesis concerning the nature of cancer, now most in favour:—that the cells of which cancerous tumours are made up are designed to eliminate from the blood, in the way of secretion, some peculiar matter injurious to the system, which generates the well known cancerous cachexia. This, though only a hypothesis, indicates the present course of opinion in pathology.

One more illustration I am tempted to add, as being more immediately connected with the subject in hand; I refer to the deteriorating influence which is exerted upon the nutrition of any organ, by an *unhealthy* demand made upon it for exertion. Let the hypertrophied heart be over taxed, and nutrition fails any longer to answer the demand: let the bodily frame be subjected to a long continued and excessive demand upon its energy, and in place of the healthy increase in nutrition which work in moderation excites, there is actual wasting, a state of cachexia; and similarly, to an over-taxed brain must be referred a long list of serious nervous diseases.

In the various cases I have now cited it is to the function of the organ chiefly concerned, that we direct the largest part of our attention, in endeavouring to cure the disease: in like manner, if the same explanation be correctly applied to the degenerations in the kidney cells which exist in Bright's disease, it must be our business, whether for prevention or for cure, to pay attention first to keeping the function of the organ in a healthy state, by removing those various causes which impose excessive or unnatural labour upon its secreting cells.

But here another question presents itself; it cannot be denied that certain forms of renal disease set in with every symptom of active congestion, if not of acute inflammation, such symptoms being present from the very beginning: this is the case in what is commonly called acute dropsy; and even in the chronic form, acute congestion may be readily excited by very trifling causes. Such being the case, it becomes incumbent upon those who deny the inflammatory origin of these diseases, to explain their relation to this state of vascular excitement. This relation we are in a better position to understand at the present day, when the near connection which undoubtedly exists between inflammation and the ordinary processes of nutrition, is beginning to be perceived. Formerly, when our attention was fixed too exclusively upon the prominent symptoms of inflammation, the vascular turgescence, and the effused fluids, we were in the habit of regarding inflammation as a disease quite *sui generis*; we had neglected to observe that in the perfectly natural course of things, wonderful variations in the degree of vascular activity are being continually produced; we had forgotten that lymph, in no respect distinguishable from the product of inflammation, is poured out in the quiet process of the repair of wounds and fractures. In these and other effects we now perceive how nearly inflammation is related to nutrition, although it must be confessed that very much yet remains to be done before the exact nature of this relation is fully understood.

These remarks will find their application in the active congestions which are not unfrequently associated, at some time or other, with diseases of the kidney. The blood-vessels are the great agents in nutrition: in this respect they are second only to the proper vital action of the tissues themselves; but sound pathology requires it to be remembered that the vessels do hold the second, and not the first place, as was at one time the belief. The whole subject of cell development has placed in its correct position the relation held by the bloodvessels to every process of nutrition; for in watching the healing of a wound, for instance, we find progress effected in the transformation of the cells in the effused lymph, before any vessels have made their appearance; and in the embryo the blood is by no means the earliest tissue developed.

This secondary, but still most intimate relation of the blood-vessels to the tissues, affords means of rendering a satisfactory explanation of many cases of active congestion and inflammation, an explanation, too, which I think may be correctly applied to the case of the kidneys; for it must follow that no change can by possibility be effected in the nutrition of any organ or tissue without inducing, as a necessary consequence, a corresponding change in the circulation of the part. Determination of blood to the womb is set up as soon as the process of increased growth, or natural hypertrophy, which is instituted by conception, commences; determination to the neighbourhood of a wound is established as soon as the process of granulation has fairly begun. So long as the demand, in obedience to which this determination of blood takes place,

is within the limits of health, the excitement of the blood-vessels, though often considerable, is in perfect agreement with the normal course of nutrition; but such is not the case when the stimulus is carried beyond this point. When the infant is put to the breast, the distended veins give intimation of the increase which immediately takes place in the rush of blood, with the obvious intent of furnishing matter for the secretion of milk; but if the natural stimulus excited by the contact of the child's lips with the nipple be exaggerated by a crack or excoriation, the determination of blood runs on to inflammation: in the same manner, in any secreting organ, in the mucous membrane of the bowels, for instance, a moderate purgative increases the intestinal secretions, but a drastic purge gives rise to congestion, possibly to inflammation, with transudation of the constituent elements of the blood; or, to bring the analogy still nearer to our present subject, a large dose of a diuretic, and particularly of the stronger ones, will be followed by congestion of the kidney. Now, in all of these cases the organ is stated to have been over-stimulated; speaking more accurately, the congestion is brought on by an excessive amount of the self same stimulus, which, in a moderate degree, had simply produced healthy rush of blood, by quickening the healthy function to a state of greater activity.

There is also another influence in operation in many stages of renal disease, which equally tends to the production of congestion. We know that an intimate relation exists between the blood and the capillary vessels of every organ, although we are not able to explain this relation. Natural variations in the circulation of different organs are entirely inexplicable by the mere impulse of the heart. How otherwise could we account for the necessary determination of blood to the womb only at certain particular periods of the individual's life? How could we explain the rush to a secreting organ to supply the material of secretion? This relation is special to every organ, and upon its maintenance freedom of circulation through each organ mainly depends; therefore, when it is disturbed in any way, stagnation of blood or congestion must follow. The phenomena of asphyxia afford a very good illustration of this fact. The admission of air to the lungs being cut off, the blood in the capillaries of the lungs can no longer be aerated; consequently, the relation which this blood ought to bear to those capillaries is altered, and arrest of the capillary circulation is the consequence, as evidenced by the loaded pulmonary artery and right side of the heart, contrasted with the empty condition of the pulmonary veins.

The same cause which operates in asphyxia, impurity of blood, operates also in disease of the kidney, and must naturally tend to increase the difficulty of the circulation through their substance; and this cause has produced a very interesting secondary change, pointed out by Dr. Johnson—namely, hypertrophy of the small arteries—such hypertrophy constituting as evident a measure of retarded circulation in the particular organ, as hypertrophy of the heart does of the obstacle presented by a narrowed orifice.

From the analogies which I have now submitted to your consideration, I think I am allowed to conclude that alterations of function in any organ have a natural tendency to induce alterations of structure, varying more and more widely, according to the extent to which the function is forced to deviate from the limits of health; also that such alterations of structure may be effected either with or without vascular excitement; but that, when the vascular excitement is present, it very often exists as a secondary result. How far this conclusion extends, we are yet ignorant; we are only just beginning to understand the phenomena of inflammation, and although we can now discover their origin in the cause I have been particularly referring to, this explanation can only apply to a very limited number of cases; it is, however, if true, of the highest value, as affording a safe position from which to start in investigating those forms of inflammation which are yet so little understood.

I must, however, add, that when once this vascular excitement has been established, it may come to play a prominent part in the case; being itself the product of an unnatural state, and being, therefore, in itself unnatural, it may easily proceed to extreme lengths; it may even absorb the principal share of our attention, and require special and active treatment; but still such treatment must be conducted with a constant reference to the function, in the disorder of which the vascular excitement originated. An obvious illustration is afforded by our treatment of acute dropsy, whether after scarlatina, or exposure to cold, or from any other cause; we are often obliged to employ active antiphlogistic treatment, as cupping, leeches, counter-

irritation, etc.; but at the same time we remember the origin of the congestion in disordered function of the kidney, and we immediately endeavour to ease the labouring organ through the medium of the skin and of the bowels; indeed, the latter branch of the treatment may not unfrequently render the former either entirely unnecessary, or required to a much less extent.

It only remains that I justify the assertion, that the condition of the kidneys, in patients who are suffering from the disease in question, is such as to have involved an unnatural state of their function. Such justification must be gained from the history of the patients. I think I may safely assert, that whenever a cause has been fairly made out, it is such as must bear directly upon the function of the kidneys; thus in the forms of acute dropsy, the sudden suppression of the cutaneous, and possibly in some cases, of other secretions, makes a sudden and unnatural demand upon the kidneys. In cases of scarlatina, of cholera, of typhus, we have the element of sudden admixture of poison with the blood, in addition, in many cases at least, to unfavourable influences acting upon the skin; disorder of the urinary functions, active congestion and its attendants, are the result; and a rapid and irregular development of epithelial cells, as manifested by the urine during life, and by the kidney after death, affords visible testimony to the labour in which the organ is engaged. More frequently the cause appears of such a nature as to act gradually, but to maintain its action through a long duration of time; the gouty diathesis and habitual intemperance constitute, perhaps, the most frequent causes of chronic renal disease, of the class we are considering; here the relation to the kidneys is also distinct; the poison of gout is either itself lithic acid, or is closely allied to that substance, and alcohol also has a direct affinity for the kidney, acting as a diuretic, and being actually capable of detection in the urine.

I shall not detain the Branch whilst I enumerate other unfavourable influences which seem to act as causes: various unhealthy modes of life, different forms of cachexia, particularly the strumous, and of blood poisoning, etc.; like those to which I have already alluded, they variously derange the constitution of the blood, either by directly introducing unhealthy matters, or no doubt more frequently by altering its nutritive changes; in some of these instances, we have to depend upon analogy for explaining the operation of the supposed cause.

I ought not to omit to add, that in explaining the operation of these various causes, we have generally to take into the account those various circumstances which predispose the organs to take on disease; without these having existed previously, many of the morbid influences we have alluded to would often fail in effecting so serious a disorder of function.

It is matter of easy observation that the cells of the kidney tubes are exceedingly sensitive to all unhealthy states of the general system; in various forms of cachexia, in chronic disease, in complaints which originate in poisoned blood, such as the fevers, pyæmia, etc., the microscope readily detects very visible changes in the kidney epithelium, even though there have been no signs of disorder in the organs during life. In these appearances, which a small amount of microscopic experience will enable anyone to verify for himself, we see perhaps the very commencement of organic change, and we can readily conceive that if the unfavourable condition, in which these appearances have originated, should act more powerfully at any time, permanent changes in these delicate organs would be the result.

In conclusion, I can only make a passing reference to the analogy presented by the liver in its diseases. In certain of the structural changes to which that organ is subject, we recognise great resemblance to those which take place in the kidney; and this resemblance becomes closer in exact proportion as we discover more of the nature and operation of the causes of liver disease.

ON THE SECOND SOUND AND MURMUR OF THE HEART AND GREAT VESSELS, IN THEIR RELATION TO DIAGNOSIS.

By JOHN COCKLE, M.D., Physician to the City Dispensary, and to the Margaret Street Dispensary for Consumption, etc.

[Continued from page 338.]

I PASS now to the consideration of the second sound and murmur of the pulmonary artery. It has been already mentioned, that any change which places the artery in more immediate contact with the chest wall—contraction of the left lung, for example—allows its pulsation to be both seen and felt, and unequivocally intensifies its sounds; excavations in the supe-

rior and middle lobes of the lung; induration of its anterior border, by immediate pressure upon the artery, modifies the sounds.

The following case strikingly illustrates such change from this latter condition. Jane Barnes, aged 30, married, of short stature and florid complexion, was admitted a patient of the City Dispensary on December 11th, 1856. She attributes her illness to sleeping in a damp bed six years ago. Since that time she has suffered from dyspnoea and cough, with yellow expectoration. At present, in addition to these symptoms, she complains of pain in the shoulder blades, the right more particularly; profuse nocturnal perspiration; very slight loss of flesh; she lies most easily upon her right side. She has not menstruated during the last two years. She has never had hæmoptysis. Her mother is still living; but consumption has occurred in some members of her family.

Inspection. The chest is deficient in symmetry. The left scapula is nearer the spine, and not nearly so mobile on inspiration as the right scapula. The apex of the heart is nearly *in situ*. The fingers are clubbed. She has slight lividity of lips.

Palpation. A silvery *fremitus* is felt over the second and third left costal cartilages. Pulse weak; slightly quickened, but quite regular.

Percussion. There is marked dulness and sense of tactile resistance from below the left clavicle to the mamma. A somewhat tubular note is heard under the right clavicle.

Auscultation. Under the right clavicle are heard phenomena of cavern or dilated tubes. Under the left clavicle, bronchial breathing; the respiratory murmur is absent to the mamma. Between the second and third left costal cartilages, both sounds of the heart are entirely replaced by hoarse murmurs of nearly equal length and intensity; loudest during expiration. To the right of the sternum, the sounds gradually acquire the normal character, as also in the carotids and at the apex.

Dr. Elliotson, who examined this case, made the interesting observation that, upon placing the patient in the recumbent position, the first murmur became converted into leather creak sound.

It is stated also, upon the highest authority, that insufficiency of the mitral valve, *par excellence*, induces marked intensification of the second sound of the pulmonary artery. That such result may occasionally obtain, seems admitted by most observers; but, even at the risk of being considered thoroughly in error, I must express my total dissent from the universality of this proposition. Of course, if it be assumed that insufficiency of the left venous valve cannot exist without such accent characterising it, I have no argument to offer; as *post mortem* examinations, made of cases with the observed sign, do not very often present themselves to my notice. I can only say, I have made many examinations after death where the mitral valve had unquestionably allowed free regurgitation, and the sign before death did not exist in any marked degree; and again, in a large number of cases observed upon the living, with all the other physical signs, save increased accent, supposed to characterise such change, we are forced to the conclusion, that such signs always indicate some intraventricular change, other than regurgitation, a consequence, I fancy, opposed to our diagnostic convictions. In support of the views advanced, I think I may cite a high authority, Dr. Walshe.

In extreme stenosis of the aortic orifice, as Kurschner has most truly observed, the chambers of the heart and pulmonic system are also kept unnaturally loaded, and would, theoretically, tend equally to a more marked accent of the second sound of the pulmonary artery.

In marked aortic, as in mitral valvular disease, however, it appears to me that we often render ourselves liable to the charge of confounding relative with absolute increase of sound of the semilunar valves of the pulmonary artery. In both instances, the aorta, receiving a diminished supply of blood, would have its valvular sound of closure weakened by want of the requisite distension, leaving that of the pulmonary artery in relative excess.

I would also hazard the passing remark, that I believe slight mitral regurgitant murmur to be extremely common, and that it neither necessarily intensifies the second sound of the pulmonary artery, nor develops dilated hypertrophy of the chambers of the heart.

That congenital malformation of the valvular apparatus of the pulmonary artery, and by inference, a patent condition of the *foramen ovale*, may also cause the second sound to be entirely masked by a prolonged systolic murmur, has, I conceive, a high degree of probability in its favour. I specially refer to a