would admit. He laboured much in his breathing, with but short intervals of rest. His tongue was coated; his stomach painfully distended; the urine showed an acid deposit; the bowels were costive; the pulse 70, and feeble. The shoulders were much elevated; the countenance anxious. He was not able to lie down at night, and had a copious morning expectoration of a frothy mucus. The right lung was in an emphysematous state. He was in the habit of eating puddings and pies, much salt pork, and plenty of broth and vegetables. I ordered him some mild alterative pills and three grains of the extract of conium four times a day, and gave him the same directions as to diet as in the former case. At the end of a month, he said he was quite relieved, and wanted a more generous diet, as he wished to go to his labour on the farm again. I increased his animal diet to six ounces daily; and he now takes his place on his father's farm, and works as other men.

[To be continued.]

ON ALBUMINURIA, AND ITS FERRO-ALBUMINOUS TREATMENT.

By DAVID NELSON, M.D. Edin., formerly Physician to the Queen's Hospital, and Professor of Clinical Medicine, Queen's College, Birmingham.

IN treating the above subject, it is preferable to use the term albuminuria as a statement of fact, rather than the theoretical and uncertain name of Bright's disease, or fatty, or granular kidney, etc. This is especially desirable in all cases that have not as yet resulted in death, or been subjected to a *post mortem* examination; because it is now an established conclusion that the specific symptom of albuminous urine may arise from many and various changes occurring within the kidneys; or, on the other hand, may be accompanied with no detectable disease at all of an organic nature. Upon this latter point I had made up my mind several years ago, on the ground of personal observation, and am confirmed in such a view by the able and practical comments of Dr. Graves of Dublin, who says upon this subject, "What I wish to call your attention particularly to in this case is the state of the patient's urine. On his admission, we found that his urine was highly albuminous; when submitted to the action of heat, at a temperature of 170°, it coagulated rapidly, and showed distinct traces of the presence of a large quantity of albumen; yet, under the use of opium in moderate doses, this man's urine became, in two or three days, perfectly free from any trace of albumen, and has continued so ever since. Now this case alone would be a sufficient refutation of the opinions of those who look upon albuminous urine as a pathognomonic sign of disease of the kidneys, as described by Dr. Bright, and who are in the habit of marking such cases in the hospital as cases of 'Bright's kidney.'" He pertinently adds, "It appears rather strange that a man should have Bright's kidney to-day, and not have it the next day"; and concludes by saying, "We have had a great many instances of this kind". And so had Dr. Blackall, long prior to Dr. Bright, in his cases of serous urine, which is only another and older name for albuminuria.

I feel the more bound to insist upon this prefatory remark, in order to avoid any misconception as to the object of this paper; for, of course, it is not intended to import that the line of treatment pursued is curative or restorative of kidneys that have become disorganised, or, as one may say, disglanded; but simply to uphold and illustrate the doctrine that albuminuria, to a very great extent, and accompanied by the sundry symptoms of dropsy, etc., may exist for a length of time, and yet be subdued and staved off for indefinite periods by means of the remedies recommended.

It was in the year 1850 that I first specially bent my observation in this direction, having had several cases in hospital, most of which, under the then usual treatment by means of diuretics, alkalies, and sedatives, etc., proved fatal; while the first one to which the ferro-albuminous remedies were applied recovered beyond all expectation, and became, amongst the others, the subject of some clinical lectures at the College, which were published in this JOURNAL in the year 1852. To these particular lectures, and the cases contained in them, I would respectfully recall the attention of my brother members and the other readers of this publication on the present occasion, should they have these volumes in their libraries, and possess leisure to refer to them; but, in case it should be otherwise, it may be as well myself to refer to the views of

pathology therein broached, and their application to treatment. Thus it is observed, in reference to the varied nature of the disease, that it consists essentially in a deposit of foreign matter of low organisation within the proper tissues of the kidneys, which proper tissues thus become atrophied, and ultimately absorbed. These deposits, having no power to assume the glandular functions—although so far lying within, and having the rude outward form of a glandular apparatus—are not calculated to eliminate those excretions which ought thence to be drained, without at the same time showing a tendency to permit the most valuable reparative material in the blood (albumen) to escape along with them; and hence the common name of the complaint, from its principal symptom—albuminuria; although it must be admitted that the occasional manifestation of profuse albuminuria, in the course of other diseases tending to debilitate the nervous system and attenuate the blood, affords proof that it is possible to be sometimes a simply functional or accidental filtration of the nutrient fluids. These degenerations of structure, though all ranked under the one complimentary name of *Bright's disease*, may be of various kinds, according to the constitution of the patient and the nature of the inducing causes; and, though the most prevailing of these causes are impoverishment and debility of the body, proceeding from the scrofulous or syphilitic taint, anæmia, drunkenness, or abuse of mercury; yet another order of causes is to be looked for, in the form of local or accidental affections such as the kidneys are subject to, in common with other parts. Thus, without finding any prior deterioration of the general mass of the blood, or any evidence of a preceding local hyperæmia, we may be compelled to look for the primary source of the disorder in a simple perversion or loss of function in the renal nerves, by which the natural reactions are so modified, and the affinities so altered, that, instead of glandular or epithelial cells being formed and deposited as usual, we shall have a displacement of them by means of granular or tubercular matter, or fat. Here we would have no previous pressure, followed by absorption; but a simple death, and disappearance of the higher orders of structure, and a substitution of degenerate matters in their place. Next, the evil may proceed from a pure anæmia, in which the over-fibrinated blood may block up and ultimately obliterate the extreme vessels, thus cutting off the source of nutriment from the glandular structure, and substituting in its stead an impoverished lymph, destined, perhaps, to undergo still deeper degenerations in the course of time. Also we may have a protracted congestion, pressing upon the nervous filaments and delicate microscopic structures of the organs, thereby lowering their vitality, and preparing them to yield without resistance to the morbid precipitations which are sure to proceed from a vitiated and stagnant composite fluid like the blood. Then we may have a subacute inflammation, acting in a manner pretty similar to anæmia, so far as effects are considered, but more rapidly and expansively, in which large amounts of fibrinous exudation will be poured out, oppressing the proper structures, and leading to their destruction; while such exudation itself sinks into a lower form of organisation, and becomes the nidus of those fatty or tubercular depositions which are the last evidences of this terrible malady. Such secondary degenerations will be modified by the constitutional diathesis of each patient. Thus, in the more healthy habit, where the disease has proceeded from some accidental anæmia or local inflammation, we may only have a fibrinous or carnified kidney, without fat or tubercle; whereas, in the fatty habit, we shall have a sebaceous deposit; and in the scrofulous and mercurio-syphilitic patient, a mere granular *débris*, either generally diffused, or aggregated into larger or smaller masses, and perhaps intermixed with pus. In all cases gone to this degree, there will be found an obliteration of the vessels, so that the organ cannot be properly injected.

Now, besides the leading, though not altogether unexceptional feature of albuminuria, the symptoms that will appear in most cases are, dull pains in the loins, inclination to drowsiness, frequent but scanty micturition, nausea and epigastric tenderness, anasarca or general dropsy, and a pale puffy aspect of the countenance and flabbiness of the limbs, concomitant with a deficiency of the red particles of the blood. The facial appearance, giving a Flemish expression to persons in this country, is a very valuable one; for it will often give us a clue to the malady before its more annoying or dangerous symptoms excite attention; so much so, that the eye of the experienced physician sometimes draws its conclusion from the physiognomy alone; and I am the more satisfied of its value, because I have often found that it revealed the secret of renal disease,

as proved after death, even when there was no albumen to be detected in the urine during life.

There may, then, be albuminuria without organic degeneration, and organic degeneration without albuminuria; while the affection of the kidneys after scarlatina is a disease totally distinct from the one we are now handling—distinct in its morbo-anatomical appearances, and almost, if not quite, opposite in point of the treatment required. The leechings and the opio-antimonial appliances, so patently curative of the temporary form of disease, would certainly be quite out of the question in the chronic degeneration. This degeneration, or tendency to degeneration, I believe to be always accompanied by a deficiency of the red particles of the blood; such accompaniment being the cause of the disease when it is of constitutional origin, and the immediate and certain consequence of it when it proceeds primarily from accidental local derangement. Each case, of course, has its own peculiarities, as to the presence or absence of certain symptoms; and I have had ample opportunities of deciding that the disease may go on without dropsies in one person, without dyspepsia in another, without pain in a third, and without albuminuria in a fourth; but that the most unvarying symptom is the condition of the countenance, which in its highest degree produces a pale, puffy, pasty aspect; whilst in an ordinary way it only gives a heavy Flemish sort of expression, difficult to be described, yet easily recognised when once well observed. Yet, although such a sign is valuable, as being indicative of the probable condition, and of the precautionary measures to be adopted against it, it would not be admissible to cite each instance in which it appears as a case of kidney-disease, unless such were demonstrated after death. Medical science requires more precision; and therefore, in speaking of the cases of persons who still survive, I studiously avoid morbo-anatomical or pathological hypotheses, and strictly confine attention to the one tangible sign of albuminuria, and its partial or entire, or temporary or permanent disappearance, or its non-disappearance, under given lines of treatment.

As already observed, a very simple and unvarying formula was at one time in general vogue, arising from the frequent use of it by Dr. Bright, when he had bent his attention to this disease, and published the results. This was bicarbonate of soda, along with tincture of conium, in some stomachic and diuretic vehicle. I apprehend, however, that the treatment must be considerably varied, according to the various causes from which it may apparently spring, the varied forms which it may assume, its various stages, and the various consequences to which it may have led. Thus, when chiefly due to general debility, it will require invigorating remedies, and the usual routine of the tonic regimen, including appropriate diet, proper exercise, and free mild air. If the atrophous form of scrofula be observable, oleaginous remedies, with steel, will be required. If the degeneration be demonstrably of a fatty nature, steel will still be requisite; but rather in combination with potassa. Yet in all cases whatever, as there is essentially a great tendency to the loss of albumen from the system, and a deficiency of the blood-forming elements, so there is required, in conjunction with any other remedies employed, a very full supply of albumen in its most digestible form, along with a more than usual supply of iron, and perhaps the calcareous phosphate. And as, when the kidney is almost completely destroyed as a gland, life may yet be prolonged by furnishing the materials necessary to its operations; so, when the disease has not as yet effected such destruction, there remains the probability of restoring the natural condition of the blood, and so renovating those minute cellular and tubular structures that were on the point, as it were, of undergoing that fatal degeneration from which, if once thoroughly effected, they never could rally.

In the cases about to be cited, successful and unsuccessful, other remedies were certainly employed besides the substitutive appliances of albumen and steel, such as digestives, powerful tonics and diuretics, etc., as required; but I viewed the latter as only auxiliaries, however useful; and the former as the essential necessities for recovery; that is to say, that the ferro-albuminous supplies held out a rational hope of restoration, even without such aid; while all the forces of quinine or peppine, or what not, could hold out no chance of amendment, without the due administration of the iron and albumen as direct representatives or analogues of the losses that were being sustained by the system from day to day and from hour to hour, through periods of years, or of months at least.

CASE I. *Albuminuria complicated with Hæmorrhage and Polypus Uteri, and terminating in Death.* Mrs. C. M., married, aged 32, first consulted me on October 24th, 1853. She stated that she had formerly possessed a good colour, and been

quite robust; but, for a considerable time past, had lost that colour, and become excessively weak, which she attributed to frequent and profuse uterine hæmorrhages, which came on at uncertain periods, and rendered her quite prostrate. No cold applications nor other means had ever arrested them. Now she was blanched in countenance, and swelled under the eyes and over the face generally. She had severe dragging pain in the back; and her thighs, legs, and feet, were enormously distended with fluid. She was also troubled with constipation and continual sickness, vomiting several times in the day. She first took aperient pills, and a calmate stomachic mixture of calumba, cardamoms, and hydrocyanic acid, which had the desired effect; and when the urine was tested, it was found to be almost entirely albuminous. As the dropsical swellings were so oppressive, the first attempt made, after settling the stomach, was to draw off the fluid; and she took a drastic purgative every second or third day, with diuretics in the intervals. These she continued till the 14th of November, when the anasarca was very much reduced; and she was put under a course of steel and albumen, taking full doses of the tincture of iron in the peptic liquor, and four eggs a day. From these remedies she seemed to derive much benefit, as her colour improved, her appetite became good, and the anasarca almost entirely disappeared. One relapse she had, from some congestion of the lungs supervening, along with an increase of pain in the back, and tumefaction of the feet and eyes; but a belladonna plaister to the back and a blister over the chest subdued that attack; and, by the end of December, she looked and felt comparatively so well, while the albumen had also diminished to about a fourth of the urine passed, that she stopped visiting me, under the belief that the complaint had "made a good turn", and she might now be better without constantly taking medicine. So she went on until the 30th of March, 1854, when, after some preliminary swellings and loss of colour, she was seized with a profuse flooding at her monthly period, which rendered her again prostrate. This was arrested by means of acetate of lead and ergot; and, on examination, it was ascertained that the hæmorrhage arose from the presence of soft polypous growths; but she was in too distressed a state to submit to any operative interference on the part of the surgeon. She recommenced, however, with the steel and albumen, and diuretics, which again had the effect of removing the swellings and restoring her strength in some degree, till the 24th of April, when hæmorrhage to an alarming extent again set in, but was stopped, as before, by the lead and ergot. She did not rally, however, as before, but continued to be troubled with frequent vomitings; while her face was blanched and swollen, and her limbs distended with fluid. The albumen also was as great as ever. The vomiting was subdued by means of calumba and hydrocyanic acid, and tablespoonfuls of food at a time; but the month had not elapsed ere she had, on the 20th of May, another profuse hæmorrhage, which was stopped by ergot, with sulphuric acid and opium. Some time after this, she did submit to the operation for removal of polypus. Had her disease consisted in that alone, doubtless she might speedily have recovered; but it had been delayed too long, and the kidney disease had gone too far, and altogether she was left too destitute of constitutional stamina for any hope of recovery; and she died not long thereafter.

The case is adduced, not as an example of the beneficial effects of the remedies (though at one period they evidently were tending to good results, had they been persevered with, and the hæmorrhages effectually stopped by an earlier operation), but as instancing the connexion between this disease and the drainage of the red particles of the blood from the body, as well as the difficulty of managing such persons.

CASE II. *Albuminuria, preceded by Hæmorrhage; Relief.* Mrs. E. W. I was consulted in this instance on October 18th, 1855. The patient was a married woman, keeping a shop, and aged 54. She had had a family, had been subject to much discharge at her monthly periods and in her labours, and had lately become pale and weak, from her confinement to the house and over-devotion to business, as the family thought. She had long had an aversion to animal food, and lived principally upon tea, bread and butter, puddings, etc. For some time past, her legs had been swelled, and her face puffed; but she did not seek medical aid till the swellings had seized the thighs, and had also appeared round the throat and under the armpits, etc. Diuretics were then employed; but they did not appear to act; vomitings set in, and she became worse in all respects. When I was called, the swellings were immense; the legs were wide asunder, from their size; the eyes were scarcely visible; and the chest oppressed with congestion, in-

cluding heart and lungs. A rank urinous odour proceeded from the skin; and drowsiness, approaching to stupor, was permanent. On examination of the urine, four-fifths of it were solid albumen.

The first thing to be done was to allay the gastric irritation, and relieve the vital organs. After overcoming the sickness, purgatives were employed, and blisters were applied over the chest and behind the ears; while tonics, stimulants, and diuretics, were combined. She had the compound tincture of bark, with ammonia and squills; and in the course of a week, after recovering her natural consciousness and quickness of mind and spirits, the swellings began visibly to decrease. After another week, she could sit up; the swellings were confined to her legs and under her eyes; and she was then ordered animal food, including several eggs a day, in one shape or other; and, as she disliked liquid medicines, she took a pill of the sulphates of quinine and iron thrice a day with her food, and only the liquor pepticus in water or beef-tea when the meal should feel heavy on the stomach. Neither did she object to the nitrous ether, and accordingly had from one to two drachms of it in gin and water after dinner and at bedtime. In the course of a month, she felt much stronger; her colour was less cadaverous; and there was no anasarca, beyond a general puffiness of the limbs and face; the feet dimpling under pressure towards night, and the skin underneath the eyes being swollen in the mornings. The albumen was then only one-third of the urine.

The same remedies and regimen being continued, the symptoms, after another month, remained favourable; the cheeks were of a healthy colour; she was active, and free from swelling—in fact, said she had nothing to complain of; while the albumen amounted to not more than a sixth. The disease still existed, but was kept in check; and relapses occurred, but not with such severity.

[To be continued.]

Transactions of Branches.

EAST-ANGLIAN BRANCH.

CALCAREOUS DEPOSIT IN THE LIVER.

By RICHARD FAIRCLOTH, Esq., Newmarket.

[Read at Newmarket, June 21st, 1860.]

A. R., aged 68, having retired to bed in her usual health on the previous evening, was seized about six o'clock A.M., on June 4th, 1860, with sickness and pain in the epigastric region, becoming more and more severe, and extending over the abdomen. Some brandy and water was given, but almost immediately rejected. Castor-oil was also administered, but it returned in half-an-hour. Fomentations were applied to the body; they failed to give relief. One grain of calomel with half-a-grain of opium were given every three hours, but the patient, in spite of all treatment, gradually became collapsed, and died about eight o'clock P.M., about fourteen hours from the commencement of the illness.

The deceased had been always feeble and weakly; she had a very small appetite, and suffered much from "shortness of breath". Some ten or twelve years ago, she was under treatment for chronic dyspepsia: she had then great pains about the region of the stomach, especially after meals; and her countenance exhibited very much the appearance of a person suffering from malignant disease of the stomach or duodenum. For the last few years, she had received no medical treatment, nor had she experienced to any great extent a return of her old symptoms; but she was still "feeble and weakly".

POST MORTEM APPEARANCES twenty-four hours after death. The body was that of a thin, spare woman, about the middle height, and sufficiently well nourished; the abdominal muscles being covered with fat to the depth of half-an-inch, at least. *Chest.* The lungs were pale, but crepitant; they were small, shrunk, and collapsed. The right lung adhered very firmly to the costal pleura and to the diaphragm. The latter was much drawn up, diminishing very considerably the thoracic cavity. The left lung was less adherent, and otherwise healthy. About half-a-pint of serum was found in the left side of the chest. The pericardium also contained about four ounces of fluid. The heart was flabby, pale, soft, and natural in size; the valves

were not examined. *Abdomen.* About three pints of reddish coloured serum were found in the peritoneal cavity. The omentum was large, containing a fair quantity of fat, and lying loose over the small intestines, which, on their peritoneal surface, presented a glistening or transparent aspect and a pinkish hue. There were no adhesions betwixt the convolutions, nor effusion of lymph upon the surface of the bowels. The diaphragm was pushed up, drawing the liver with it (and almost out of sight). The gall-bladder was visible, and easily removed from its connection to the liver. It was half filled with gall-stones, varying in size from a bean to a small shot; it contained but little bile. The duct was pervious. On the posterior or under surface of the liver, near the free edge and to the left of the fissure, was a conglomeration of the surrounding structures, very difficult to discriminate, from their close adhesion to each other, and being united by a deposit, in the substance of the liver itself, of stony hardness, to the extent of three-fourths of an inch in depth, and of the circumference of half-a-crown. The spleen was normal. The stomach was pale and empty, containing only a small quantity of mucus mixed with bile. The large bowels contained feces, but were not distended. Neither kidneys, uterus, nor bladder, were examined.

REMARKS. The immediate cause of death was probably spasm of the gall-duct, during the passage of gall-stones, producing in an enfeebled constitution so much irritation and depression, as to cause the effusion spoken of into the pleural, pericardial, and peritoneal cavities. The hardened deposit in the substance of the liver was probably calcareous—the result of old mischief, perhaps abscess in the liver, which had retired or become partly absorbed, and the remainder becoming transmuted into this singular structure as hard as ivory, nay, almost as indurated as marble, and being the size of a small flattened fig. But how life could be carried on, and the body present such indisputable evidence of receiving sufficient nourishment, is indeed remarkable.

CASE OF EXTRAUTERINE PREGNATION.

By C. G. MARSHALL, Esq., Woodbridge.

[Read at Newmarket, June 21st, 1860.]

Mrs. F., a young woman, aged 28, who has been married five years and three months, but never had a child, gives the following account of her previous history.

She was sickly as a child; she menstruated first at the age of fifteen; but the last six years there has been little or no colour. Three years since she miscarried at two months, after which she suffered much for eight or nine weeks with pains in the back and abdomen, so severe that she could not breathe, except in the upright position. She continued afterwards in her usual health up to the beginning of September 1859, when everything like discharge from the uterus or vagina ceased. In October, she began to suffer excruciating pain in the abdomen and back, with twisting pains at the umbilicus; and in the latter part of the month, passed "a something like a piece of flesh" from the vagina. The body then began to increase in size, the breasts to enlarge, and all the concomitant symptoms of pregnancy made their appearance. In the latter part of December, the movements of the child were distinctly felt; but a discharge of blood twice took place from the lungs, of a bright red colour, at times corresponding to her monthly periods. She now became much emaciated, and so worn out and exhausted from pain, that she could not walk without assistance.

In April, the body began to subside, the breasts to take on their natural appearance, the movements of the child were no longer to be observed, and peritonitis set in, for which she was treated actively by calomel, opium, leeches, etc. This subsided, and took on a chronic form, with a fulness in the left iliac region. The pains in the abdomen and back continued; and added thereto was continued sickness and vomiting, so that for upwards of six weeks scarcely any food was retained on the stomach. On May 19th, she began to pass from the anus the bones before you. They were found imbedded in the faecal matter, and frequently required assistance to get them away. The woman is now much emaciated, and continues to suffer severe pain in the body and back. The sickness has ceased three weeks; the appetite is improving; but there is no return of the catamenia; and she is continuing to pass bones. The last that appeared were the body of the sphenoid, and the bones either of the hand or foot.