Original Communications.

ON GLYCOGENESIS.

By JOHN SLOANE, M.D., House-Surgeon to the Leicester Infirmary, etc.

[Read before the Leicester Medical Society, April 20th, 1858.]

SOME of you may probably desire to hear the *rationale* of the saccharine method of treatment of diabetes, which seems diametrically opposed to the old plan. I need not tell you that the cause of diabetes was formerly considered to be a derangement of digestion, by which sugar was formed in the stomach, afterwards absorbed, and passed into the blood, from which it was rapidly excreted by the kidneys. This view of the pathology of diabetes has been proved to be erroneous by the able researches of M. Claude Bernard, which are published in his *Leçons de Physiologie Expérimentale*, which, I regret, have not yet been translated into our language. I hope I shall not be considered to waste the time of this Society by reading the following extracts from these valuable lectures, describing investigations which have revealed to us many striking facts regarding what is called glycogenesis, or the formation of sugar.

Glucose, the variety of sugar found in the urine of diabetics, is generated in the livers of animals throughout the animal kingdom, almost wholly irrespective of the nature of their food. The glucose secreted by the hepatic cells passes into the hepatic veins, thence into the inferior vena cava, and through the right side of the heart to the lungs, where, being exposed to the atmosphere, it sometimes completely disappears. M. Bernard has found sugar in the livers of mammals, of birds, of reptiles, of fishes, of molluscs, and of articulated animals. He has found it in omnivorous, herbivorous, and carnivorous animals. That the secretion of sugar is independent of the nature of the food, he proves by many experiments, of which I shall mention the following. He fed dogs exclusively on flesh for six or eight months; and when they were killed, at the expiration of that period, he found as much sugar in their livers as in those of dogs fed upon a mixed diet. Owlets taken in their nests were fed exclusively on raw bullock's liver for three months, and were then killed: their livers always contained the normal quantity of sugar. Two dogs were fed solely on flesh, three on both flesh and bread, and two on amylaceous or saccharine food; they were all killed at as nearly as possible the same period of digestion, and the results of the chemical examination of their livers showed that the quantity of sugar secreted did not depend on the nature of their diet.

Rollo recommended the use of fat for diabetics. M. Thenard and Dupuytren made them eat lard. We have fed dogs with lard and axunge; and we have found this very curious fact, that, under the influence of this alimentation, the sugar diminished in the liver absolutely in the same manner as if the animal had been kept fasting. In dogs to which M. Bernard has given nothing but pure water, he has found the secretion of sugar kept diminishing, and it ceased to appear about three or four days before its death. For the first thirty-six hours, the quantity continues considerable, but during the following days it diminishes very rapidly.

A dog, having fasted thirty-six hours, had a copious repast of boiled sheep's head, and, three hours afterwards, was killed. The blood in the portal vein, previous to its entrance into the liver, contained no trace of sugar; whereas, in the blood from the hepatic veins, there was a considerable quantity. This experiment, writes Bernard, would alone suffice to cause one to admit, as a natural and necessary conclusion, that the sugar is produced in the liver; yet we have accumulated proofs of every kind about this proposition; and we have shown that the hepatic tissue constantly contained sugar, and that it was the only tissue of the body which offered this character.

In an animal fasting, the blood which arrives at the liver presents no trace of sugar; that which leaves it contains a considerable quantity. Inversely, the blood which arrives in the lung contains sugar; and that which leaves it presents no trace of this substance. The sugar in this physiological state remains hidden between the liver and the lung, and does not show itself at the exterior. This statement is true only in an

animal fasting. When the digestion commences, the quantity of sugar gradually augments; yet during the two or three hours following the ingestion of aliment, notwithstanding the increase of the saccharine secretion, all the sugar can be destroyed before it arrives at the arterial system; and it is only after the lapse of time that the production of sugar surpassing the limits of destruction becomes temporarily excessive in the organism. At this period of digestion, one finds sugar in all the vessels of the body, arterial and venous, and even in the renal arteries; but the proportion is too slight for any of the sugar to pass in the urine. Yet we shall see that, under certain physiological circumstances, the quantity of sugar can be increased to the point that it passes off in the urine without the animal being diabetic. Under the ordinary circumstances of digestion, this species of saccharine overflowing is manifested equally with animal or amylaceous diet, and it lasts about three or four hours. It is not less than six or seven hours after a meal that the excess of sugar in the blood commences to disappear, and that the equilibrium between its production and its destruction tends to re-establish itself as before digestion. This species of oscillation, which the glycogenic function presents, it is very important to know; for in the pathological state (diabetes) we find exactly the same phases, with the exaggerations we should expect in this malady. Different observers—Rayer in France, and Traube in Germany—have remarked that there are diabetics who do not pass sugar in their urine, except at the time of their digestion; and that, in the interval, their urine does not contain sugar. This phenomenon can be reconciled very naturally with the physiological fact which has been pointed out to you. There is nothing essentially different between the normal state and the pathological symptom, save the intensity of the phenomenon caused by a deviation of vital activity.

The sugar is formed from the albuminous substances; and this sugar is the result of the physiological action of the liver upon those principles, which are divided so that their oxygen, hydrogen, and carbon, are grouped so as to form sugar, whilst their azote enters into other combinations, and probably into the azotised principles of the bile. One does not know, indeed, any other origin for the saccharine matter, which cannot be produced in the intestine by digestion. Experiment has shown us that, during alimentation, by means of albuminous substances, the intestine and the blood of the portal vein never contain saccharine matter of any kind. Neither gelatine nor flesh produce saccharine matter in the intestinal tube by the known digestive processes. The amylaceous matters taken as food enter as sugar into the portal vein, and, arriving at the liver in this state, are then destroyed by this organ, and changed into another matter, which has every appearance of a fatty substance converted into an emulsion *par une matière protéique spéciale*. We have said that the sugar introduced into the intestinal tube does not augment the quantity of this matter contained in the liver, but that it is there destroyed, and causes the appearance of an emulsive substance. That the sugar introduced into the intestinal canal does not augment the quantity of this matter contained in the liver, M. Bernard shows by the following experiments. He takes two rabbits, whose urine he first finds, by testing, to be free from sugar. Into the stomach of one he injects a quantity of sugar in solution, with some ferrocyanide of potassium. Beneath the cellular tissue of the other he injects half the quantity of an exactly similar solution. He examines their urine an hour afterwards, and he finds in that of the first not the least trace of sugar, while the urine of the second presents it in considerable quantities. But you may say that this difference may be accounted for by the intestinal absorption being less rapid than the subcutaneous: but in both the ferrocyanide of potassium was readily detected in the urine. This will prove that the absorption is equally effectual in the intestine as under the skin, but that, in the first case, the solution has abandoned one of its constituents, the sugar, in traversing the liver; whereas this has not taken place in the second instance. He arrives at similar results in the following experiments. Through a small opening in the abdomen of a rabbit, he injects a quantity of the same solution into one of the branches of the portal vein; and into the jugular vein of another rabbit he injects the same quantity of the same solution. It is clear that, in this mode of operating, we cannot have any difference in the absorption, as in both cases we introduce the substances directly into the blood. Nevertheless, we obtain exactly the same result; that is to say, that in the rabbit, in which we injected by the jugular, the sugar has passed into the urine with the ferrocyanide of potassium, and with very great rapidity; whilst in the rabbit injected

by the portal vein, the ferrocyanide of potassium alone will have passed into the urine, where one cannot find the least trace of sugar. These experiments are very conclusive. Bernard proves by experiment that starch, taken as food in the intestine by the influence of the pancreatic juice, becomes converted into sugar; and this passes into the portal vein. That sugar is destroyed by the liver, receives further confirmation, he states, by the facts known in the fattening of cattle. You all know that animals fatten most by the use of food in which starch predominates; that the geese and the ducks, in which the fat livers are artificially produced, are gorged with a *pâté* of maize or other amylaceous food; that the fat formed by an animal is not in proportion with the adipose matter which it takes; that, on the contrary, the animals which only eat fat, far from becoming fat, get lean rapidly. Hereafter it is not only the biliary secretion which we shall have to look upon in the liver; it has two other functions of capital importance—one the production of sugar, which is dependent upon the aliment containing albuminous matters; the other, the production of fat, which is dependent upon the amylaceous and saccharine matters in the food.

Cane-sugar is never destroyed; it is constantly eliminated by the urine when it is injected directly into the blood; but this sugar, when in the intestine, is in part, at least, transformed into glucose. The latter, on the contrary, injected into the blood, can be destroyed in certain proportions.

When we prick the mesial line of the floor of the fourth ventricle, in the exact centre of the space between the origins of the auditory and pneumogastric nerves, we produce an exaggeration of the hepatic (saccharine) function, and of the renal secretion; if the puncture be effected a little higher, we very often only produce an augmentation in the quantity of the urine, which then frequently becomes charged with albuminous matters; while, if the puncture be below the indicated point, the discharge of sugar alone is observed, and the urine remains turbid and scanty. Hence it appears that we may distinguish two points, of which the inferior corresponds to the secretion of the liver, and the superior to that of the kidneys. As, however, these two points are very near to one another, it often happens that, if the instrument enters obliquely, they are simultaneously wounded; and the animal's urine not only becomes superabundant, but at the same time saccharine. The urine becomes saccharine in from one to two hours after the operation, but seldom continues for more than a day.

The secretion of sugar is not under the direct influence of the pneumogastric nerve; for if it be divided before irritating the floor of the fourth ventricle, sugar still appears in the urine. Bernard believes that the influence is transmitted by reflex action through the ganglia of the sympathetic.

There is a phenomenon which is manifested, for example, when, after fasting a certain time, a great quantity of sugar is taken. The intestinal absorption then proceeds with extreme rapidity. A great quantity of sugar arrives in mass in the liver; the mechanical circulation much prevails over the chemical; the sugar is poured into the general circulation in proportion much greater than occurs in the normal state; and it passes then into the urine, where its short-lived presence can be found for a certain time.

M. Bernard, after a great many experiments in reference to the subject, has proved that there is a species of election in the excretion of matters which pass out of the organism. Sugar is eliminated in two ways only—by the kidneys, and by the mucous membrane of the stomach. When sugar is injected into the blood of an animal to saturation, and puts it for a time into a state of diabetes, we do not find sugar in the saliva, in the tears, pancreatic juice, bile, nor perspiration; whilst the urine and gastric juice contain it in proportions more or less notable. These results entirely resemble those obtained in diabetic patients. Lehmann states, however, that he has obtained sugar from the saliva of a diabetic. The presence of sugar has been pointed out in the expectoration of diabetics. Bernard admits that sugar can be had in notable quantity in the expectoration. But, he writes, we must not confound the bronchial mucus which these patients, almost always phthisical, in the last stage of the disease expel in abundance, with the salivary secretion properly so called: it is the mucosities formed in the lung which contain the saccharine matter. Nevertheless, this fact is not constant; for M. Rayer has reported to the Society of Biology a case in which the expectoration of a phthisical patient examined by M. Wurtz did not contain sugar. Bernard proves by the following experiments the statements regarding the election in excretion of matters which pass out of the organism.

He takes a dog with a parotidian opening, into which he inserts a tube. Nothing flows by this tube, which proves that the secretion is not continuous. By putting in the mouth some vinegar he excites the flow of saliva, which passes out of the tube rapidly in large drops. He next injects into the jugular vein of the animal a solution containing sugar, prussiate of potash, and iodide of potassium. Immediately after this injection the salivary secretion is again excited in the same way. The saliva is received into three glasses. One is examined for sugar, and none is found. The sugar, therefore, does not pass in the saliva. The second is examined for prussiate of potash, and it is not present. The third is found to contain iodide of potassium. This substance then passes immediately into the saliva, whilst the prussiate of potash and the glucose, equally soluble, cannot be found. In the saliva extracted before the injection, none of the substances exist. In the urine of the same animal after the injection the prussiate of potash is found in considerable quantity, and the iodide of potassium in small proportion. As regards the sugar, there is none yet, but we shall find it presently. It requires an hour or more for the sugar to appear in the urine.

The urine then eliminates all these substances in a manner more or less rapid. The prussiate of potash appears first and the glucose last.

There is another secretion in which the presence of sugar can be found; this is the gastric. The passage of the sugar into the stomach has surprised most of the observers who have seen long since that when diabetics vomited, although they had eaten nothing but flesh, the vomited matters were saccharine. When it was believed that diabetes proceeded from a perversion of the digestive functions, it was considered that the flesh was changed into sugar in the stomach. But one need not now be mistaken; the flesh is not saccharine. Bernard himself has observed that, in diabetics who vomit fasting, in the vomited matters the presence of sugar could be found. But this has only occurred when the disease is at its greatest intensity; and in all those cases, even in the animals which have been rendered artificially diabetic, it is much more difficult to obtain the passage of glucose into the gastric juice than into the urine.

The sugar is formed, as we have seen, at the expense of the albuminous substances. In the healthy man it is clear that a part only of these matters is consumed for this purpose. The diabetic who forms much sugar expends a very large quantity of azotised material; the blood is impoverished; and, although the patient eats enormously, he gets thin like a man badly nourished. The liver takes in a manner the ration of the other organs, which undergo a considerable attenuation, because the albuminous elements are transformed into sugar.

M. Bouchardat has proscribed the use of amylaceous and saccharine matters in the food of diabetics. The facts which Bernard has himself witnessed in the practice of M. Rayer prove clearly the utility of azotised aliment. In the regimen of these patients, writes Bernard, vegetable aliments ought to be forbidden, as it is evident that they augment the functional activity of the liver. You know, also, that they are excitants of the kidneys; that they are much more diuretic than animal matters. Thus all the herbivora pass much more urine than carnivorous animals. In the azotised regimen diabetics have the advantage of food which is not diuretic.

I have at great length reminded you of M. Bernard's views regarding the formation of sugar in the animal economy. As some of them are of so novel a character, and so little in accordance with the notions formerly held, I have thought it advisable to mention the experiments upon which he founds his opinions. That they will, upon further investigation, be more or less modified, is not improbable; but they have been very generally received by the most distinguished physiologists and pathologists.

From M. Bernard's investigations, we learn the following facts of importance in reference to the saccharine plan of treating diabetes.

1. Sugar may be rationally administered to diabetic patients, inasmuch as the sugar found in the general circulation is almost always secreted by the liver, and as sugar introduced into the intestinal tube in its passage through the liver is there altered and converted into an emulsive substance, which serves to fatten these patients, and thus to counteract their tendency to emaciate.

2. Substances which contain glucose—such as honey and fruits, should be given to diabetics in preference to those containing cane-sugar, because the latter is not destroyed when injected into the blood, but is constantly eliminated by the

kidneys; whereas glucose can be destroyed in certain proportions.

3. Cane-sugar would be beneficial to a certain extent; as when taken into the intestine it is in part at least transformed into glucose; but if given in too large proportions to be thus completely transformed, the disease would be probably aggravated by the presence in the blood, and subsequent excretion by the kidneys, of the former variety of sugar.

4. The glucose should be given in moderate quantities at a time, and frequently, rather than in large quantities at long intervals; because, when much sugar is taken fasting, it is absorbed too quickly to admit of its complete destruction in the liver, and it passes into the general circulation, whence it is eliminated in the urine.

TOPOGRAPHY AND CLIMATE OF LLANDUDNO, NORTH WALES.

By T. C. RODEN, F.R.C.S.

LLANDUDNO, as the guide-books inform us, stands on an arm of land about four miles in length, jutting out towards the west from the mainland of North Wales at Conway. This little peninsula is terminated by a huge limestone rock, seven miles in circumference, and 750 feet in height—the Great Orme's Head—under and along the south side of which the town is built. About a couple of miles in front—that is, to the south—runs a chain of hills from the Little Orme's Head on the east towards Conway bay on the west; while a beautiful semicircular sweep of the ground between the two Ormes encloses the waters of Llandudno bay. It is chiefly this peculiar formation of the locality which impresses it with distinctive qualities, beneficial to particular classes of invalids; while the general mildness, dryness, and salubrity of the air render it an agreeable place of resort to all. From the experience of a residence of three years, I am enabled to designate the climate as mild, yet bracing, and remarkably free from humidity.

It is generally supposed by strangers that the coast of North Wales is cold and bleak; and many who visit this spot find it difficult to account for the opposite state of the case with regard to Llandudno. The shelter afforded by the Great Orme's Head from the north and north-east winds, together with the peninsular form of the site, afford the true explanation of the fact; for, being embraced on three sides by the sea, the temperature becomes more equalised. In winter, the water being warmer than the air, imparts a portion of its caloric to the atmosphere; while in summer the phenomena are reversed. The Gulf Stream, too, passing the south coast of Ireland, probably produces no unimportant share in the result.

These circumstances will make intelligible the quality of mildness, which is most striking in the winter months; so much so is this the case, that there are few points on the British coast so well fitted for the abode of invalids in winter as this. The thermometer fully bears out this statement; for my observations of it, noted regularly during two years, prove that the temperature is milder by 10° or 15° in severe weather, than in several situations in England with which I have compared it; and the contrast is the greater the more intense the cold.

It might naturally be expected, therefore, that in summer the heat would be more oppressive here than elsewhere. That it is not so, one or two facts will serve to show. The range of hills to the south, before alluded to, and the mountain on the north, bound a valley, along which the westerly winds (which generally prevail), cooled by the sea over which they sweep, pass in a grateful and refreshing current, and produce, together with the refrigerating influence of the volumes of water around, a temperature certainly not higher than that of other places, and lower than many. To substantiate this, I need only mention that from observations taken at the telegraph station on the mountain (at an altitude seven hundred feet above the level of the sea) during the hot weather of last summer, it appeared that the thermometer was usually about 5° higher than in the town below.

From what has already been said, it will excite no surprise that within the space of a square mile Llandudno should possess two or three varieties of climate. That part of the valley exposed more directly to the action of the winds is cooler in the summer, and in the winter is so bracing as to lead few to prefer it; at the foot of the Great Orme the temperature is mild at all seasons; while on the mountain side that delicious atmosphere emphatically called "mountain air" gives life and

strength to the system. The variations are not so trivial as might be imagined, but are well marked, and are severally adapted to the constitutions and requirements of different individuals. Invalids may therefore select a residence suited to their respective cases during any and every part of the year—either completely sheltered, or more or less open to bracing influences.

It is, however, to the advantages of the place as a winter residence, as not being generally understood, that I would particularly direct attention. The beauties of the scenery, the extraordinary opportunities for sea-bathing at all states of the tide, the purity and salubrity of the air, the facility of reaching all points of interest in North Wales, as well as of access to England, are already well known; but comparatively few have enabled themselves to judge of the climate from a sufficiently long residence. Visitors generally have been content to enjoy the pleasures of the scene for a time in the summer, and have probably felt a sort of compassion for those destined to spend the cold season, as they imagine, on these inclement and inhospitable shores. Far different, however, is the case of those who commit themselves to a Llandudno winter. The charms of its bright and sunny skies, clear and exhilarating air, and genial temperature—the advantages of its freedom from damp, fog, and continuous rain, and from the intense cold of most situations in northern latitudes, render it for the most part unnecessary to expatriate the weakly in health and constitution, or to seek abroad for conditions which may be ensured at home.

For the victims of dyspepsia, of rheumatism even in its severest instances, of phthisis and the various forms of pulmonary and bronchial disease, as well as for that large class of persons technically termed "delicate", the winter climate of Llandudno is admirably adapted. Several such who have, within the last few years, taken up their abode here, have experienced great benefit, and have in a remarkable degree recovered their health. Scarcely for a day throughout the year need the valetudinarian keep within doors, or refrain from that exercise in the open air so essential for the maintenance or restoration of health.

I submit, then, that these facts incontestably prove that this new watering-place is a valuable addition to the hygienic resources of the profession, as supplying a desideratum long sought for, too often perhaps on foreign shores, and certainly not hitherto realised at home.

Some meteorological and local details are reserved for a future communication.

POISONING BY ALMOND FLAVOUR: RECOVERY

By HENRY W. T. ELLIS, Esq., Crowle, Bawtry.

At four o'clock in the afternoon of last Friday, I was summoned to visit immediately the wife of Mr. Frederick Chapman, a solicitor's clerk, of this place, who was said to be in a fit. On proceeding to the house, I found my patient lying on her back across the bed, in a complete state of insensibility; her countenance was ghastly; her eyes were fixed, staring, and prominent; the pupils were dilated; the jaws firmly closed; frothy mucus exuded from the mouth; the breathing was heavy and stertorous; the pulse at the wrist was scarcely perceptible, but the carotids pulsated strongly; the extremities were relaxed. Whilst noticing these symptoms, I detected the odour of hydrocyanic acid; and, on inquiry, ascertained there had been some slight vomiting of undigested meat. This at once and unmistakeably revealed the nature of the case. Of course, not a moment was to be lost; cold affusion was therefore instantly and vigorously had recourse to, followed up by ammonia to the nostrils, and its application with camphor liniment to the chest, and with mustard to the legs. These means were persisted in for about two hours, with but slight change in the symptoms. At the end of that time, however, there were convulsive twitchings, rolling of the eyeballs, etc. In another hour, the patient became delirious, tossing herself wildly about, but gradually subsiding into sleep. Twenty leeches were in the meantime applied to the head, and warmth to the extremities (previously quite cold). Emetics and the stomach-pump were at hand, but could not be used: in fact, from the time that had elapsed before I saw her (ten to fifteen minutes after swallowing the poison), it is doubtful whether they could have been of any avail, absorption taking place so rapidly.

The case has since gone on well, the only drawback being erysipelas of the face and scalp, attributable, probably, to the leech-bites.