we exclude that insight of the understanding which is called intuition, because it is a part of its own innate energy which depends neither on memory nor the senses, though called into activity through their exercise. There can be no difficulty, then, in admitting that the disease under consideration exists in the only other set of faculties remaining; that its *exciting cause* is disease of the material organ in which they inhere; and that its *proximate cause* is a perversion of the logical understanding, and through it a perversion also of the feelings and the will. Insanity, then, may be defined as—a dangerous misconception of one's personal relations, either persistent or with lucid intervals, and with or without illusions of the senses.

It will give more completeness to this sketch if I add, that a true conception of the understanding is an idea or belief in conformity with the nature of things, and that the class of misconceptions may be tabulated as under:—

MISCONCEPTION.

1. In matters common to all men: *Error* and *Ignorance*.
2. Peculiarities gradually acquired from circumstances: *Eccentricity*.
3. Peculiarities suddenly or recently coming on from disease: some forms of *Monomania*.
4. In which liberty is dangerous to the Individual and to Society: *Insanity*.
 - a. From destruction of all mental power—*Dementia*.
 - b. From general weakness of mental power; viz., inadequate conceptions, blunted emotions, and feeble will—*Amentia*.
 - c. Strongly exciting or perverting the instincts, emotions, and will—*Mania*, *Monomania*, *Dipsomania*, *Impulsive Madness*.
 - d. Depressing the instincts, emotions, and will—*Melancholia*.

TREATMENT OF DIABETES MELLITUS.

By T. INMAN, M.D., Liverpool.

It is desirable that contributions, however small, should be made from time to time respecting any new plan of treatment proposed for severe and almost intractable diseases. As yet we have had few reports upon the practice adopted by Dr. Budd, of Bristol, in diabetic cases, although it was one eminently deserving of consideration.

I have, in my hospital experience, had four cases of diabetes mellitus under my care; two prior to the publication of Dr. Claude Bernard's researches, and the promulgation of Dr. Budd's views; and two since. Of the first two, I will only say that the patients went out of the hospital worse than when they came in, although no attempt was spared to benefit them.

The last two cases came to very different conclusions. I may, for the sake of brevity, describe both patients as being labouring men, about 40 years of age—ill for many months. The quantity of urine passed was over twenty pints daily, the specific gravity 1,045; the presence of sugar was ascertained by fermentation and other tests. Emaciation was considerable; and thirst great. Both had been under dispensary treatment before their admission. In adopting a plan of treatment I was guided by the following considerations:

1. The liver naturally produces sugar in a definite quantity. In diabetes there is an excess of sugar, and we may fairly infer that it comes from the liver. Opium has a decided effect in diminishing the bile producing or secreting function of the liver, and it is reasonable to suppose that it will reduce the sugar-forming function. Experience has long told us that no single remedy in diabetes has been so efficacious in diminishing the quantity, etc., of urine passed, as opium. Opium, therefore, should be one ingredient in the treatment.

2. Again, Bernard has shown that the liver makes sugar, no matter what is the nature of the food employed. Dr. Budd has shown that some patients, at least, may be benefited by saccharine food. But my patients did not long for sugar; and they did enjoy their ordinary food; consequently I neither restricted them to non-saccharine, or non-amylaceous diet, or prescribed unusual quantities of sugar. They were to have the ordinary full diet of the hospital, but more in quantity if they chose, either of bread, meat, or potatoes.

3. Again, it seemed to be clear that in diabetes, there was debility, implicating more or less the whole system; that there was danger of death by consumption; that the digestive

powers, notwithstanding their apparent energy, must be impaired; at any rate, that opium was liable to disorder the stomach, and that it could be tolerated in larger quantity if combined with quinine.

The result of these considerations was the following prescription for a pill:

Opium, one grain; quinine, two grains; to be taken every four hours. Full house diet, with porter daily.

The effect of this was soon apparent. The men began to improve rapidly and steadily; the urine diminished until it stood at ten pints only per day, with a specific gravity of 1,035. Commensurately with this, their strength and spirits increased, and they gained in flesh considerably. The opium never affected the head except on one occasion, when the patient, hoping to expedite his recovery, took a double dose. The bowels were habitually regular. The plan of treatment was neither varied or altered during their residence in the hospital. They remained under notice, the one about three months, the other for six weeks. Both left the house of their own accord, as they considered themselves sufficiently cured, and competent to do their ordinary work. I have seen one since he went out, and found that he continued strong, and, as he thought, well.

Of course I do not imagine that these two cases are sufficient to upset our older notions of the correct treatment of diabetes. I merely offer them as a small contribution to our general therapeutical stores.

I may just mention, as a curious fact, that one of my unsuccessful cases found that he received more benefit from a diet of raw beef than from any other thing dietetic or medicinal which he had taken; and that every new medicine did him good for about two days.

REMOVAL OF AN INVERTED UTERUS.

By J. BOWER HARRISON, M.D., Higher Broughton, Manchester.

In the *Medical Gazette* for April 1840, I published a case of the successful removal of an inverted uterus. At the period at which the case was published, only a comparatively short time had elapsed since the performance of the operation. It may be interesting, therefore, to supply, at the present time, such particulars as I am able to collect respecting the final success of the case.

I am more especially induced to offer these particulars, as the case itself seems to have escaped the notice of many who have directed their attention to this subject. In a paper, for example, which appeared in the *Medico-Chirurgical Transactions* for 1852, on the Inversion of the Uterus, by Dr. Gregory Forbes, no allusion is made to my case, although the paper professes to contain a complete record of what has been published in relation to this disease. I must acknowledge, however, that Dr. Forbes afterwards did me the favour to write me a letter in which he regretted the omission, and expressed his wish to supply the defect by a supplementary notice.

Some time subsequent to the operation, which I related in the *Medical Gazette*, my patient left Manchester, and I unfortunately lost sight of her. This I regretted, though I had every reason to suppose that she was in good health, for my friend Mr. Crosse of Norwich wrote to me in July 1844, expressing a wish to have full particulars of the case. He says, "I have for some time been attending to the subject of *inversio uteri* (having had a successful case of removal of that organ), with a view to writing a comprehensive essay thereon. By some great oversight, I was not cognisant of your excellent case detailed in the *London Medical Gazette* of April 1840. I hope you will excuse me for putting to you the subjoined queries, in further illustration of your case," etc.* Mr. Crosse happily discovered the patient himself, for, in December of the same year, he writes me word, "I do not think I have before informed you that my particular friend Mr. Samuel Smith, Senior Surgeon to the Leeds Infirmary, has found your extra-uterine patient, and sent me full particulars about her present state. She continues in good health, as do several others in this country whom I have seen. The gradual method you pursued, in removing the uterus by ligature, coincides precisely with my own ideas as to the best way to do it with success."

In a letter which Mr. Crosse wrote to me in June 1845, he says, "I was lately at Leeds, and saw your extra-uterine

* The original letters are in my possession.