pressure upon the walls of the cavity. The comparative frequency of scirrhus in India seems to be in accordance with the observations of the best pathologists, who have asserted, that in tropical climates the tendency of malignant disease is to scirrhus, and in cold climates in a much greater degree to fungus hæmatodes, and other forms of malignant tumours.

24, George Street, Hanover Square, January 1859.

ON DISEASED VESSELS; AND THE VALUE OF THE PULSE AS A SPECIAL SYMPTOM.

By HENRY DUNCALFE, Esq., M.R.C.S., West Bromwich.

[Read before the Queen's College Medico-Chirurgical Society, Birmingham.]

THE chief object of my communication is to draw attention to "special symptoms," diagnostic or nearly so of disease; and the consideration of this subject will enable me to record some cases of diseased arteries, which present points of practical importance.

At the primary investigation of a disease, a group of symptoms are presented to us; sometimes our attention will be drawn by the graphic description of the patient, to the principal evidences; at other times, the indefinite descriptions call at once upon our experience, and observation, to analyze them for ourselves; to dismiss the imaginary ones; to modify such as by increased nervous sensibility are magnified to the patients' own preceptions; to determine such as are directly connected with the organic lesion; so that we can describe the remaining dependent symptoms, which we know must follow in the train more correctly and in better order than our patients could themselves explain. Thus it is that we localise disease, and separate such signs as more properly belong to organs, which in the disturbed mechanism become secondarily involved. We do, in fact, recognise two classes of symptoms—the primary, or diagnostic, and the secondary, or the products of the disease resulting from functional disturbance in cooperating organs. We never lose the importance of minute crepitation in pneumonia, or of pectoriloquy in phthisis, or of double bruit in aneurism; but do we always elicit the full information to be obtained from the pulse, skin, secretions, etc. A very interesting "special symptom" in pericarditis has lately been announced by Dr. Barlow, namely, "restricted action of the diaphragm," and without it, the case quoted by him, in which the pericardium was filled with pus, and so the friction-sound prevented, would not have been diagnosed. Without albumen in the urine, would Dr. Bright have associated dropsy with renal disease? and without anæmia would Addison have directed special attention to diseased suprarenal capsules? While it is advantageous to determine diseases to be, by negatively proving them not to be elsewhere, it is equally necessary, and much more scientific, to arrive at once at the nature of a disease under consideration by the positive signs which that disease exhibits.

I premise my communication by these general remarks, as I intend to allude chiefly to the character of the pulse in connection with the cases about to be related.

CASE I. Aneurism of the Posterior Communicating Artery within the Head. The subject of this aneurism was a child 13 years of age. Of her history I have but little to say; for, being the child of poor parents, little notice was taken of her complaints, and it was only after an order from the coroner, that I became acquainted with it. The parents seem to have known but little of the child's previous state of health; they allow that at times she had complained of headache, had vomited on two or three occasions, and once had fainted. At the time of her death she complained of her head, staggered across the house floor and died.

On removing the brain, a large effusion of blood covered the base; and carefully tracing the arteries backwards, I came to the aneurism (about an inch and a half in circumference,) of the posterior communicating branch of the left internal carotid, which quite covered and concealed the parts contained within the circle of Willis.

The symptoms which marked the interesting case reported by Dr. afterwards Sir Gilbert Blane, are worth enumerating. His patient was a female 64 years of age, who was suddenly seized with a fit of giddiness and dimness of sight, succeeded by acute pain in the forehead, which remained for some time. The indistinctness of vision continued for six months, after which she was seized at intervals with giddiness, headache, and imperfect vision. She for some time saw objects double, and

finally showed signs of mental derangement, became maniacal, and died in five years after the first attack.

On examination, the optic nerves were found smaller than usual, and the symptoms so singular were referable to bulbs five-eighths of an inch in diameter, filling up the hollow on each side of the sella turcica. These bulbs were dilatations of the carotid arteries, and were filled with laminae of coagulated blood.

That these symptoms were partly produced by pressure seems evident to me, by analogous symptoms which prevailed in the case of a circumscribed abscess of the brain in a patient of my own, and who having had the abnormal changes in the special senses, as enumerated above, also became maniacal and died. In his case, the pulse was measured, slow, and occasionally intermitting, but otherwise regular.

The chief points of interest in the case of aneurism I have described, are, the young age of the patient, and the probability, that even with an aneurism so large, and lying against a structure so delicate, no very urgent symptoms manifested themselves. But scanty as was the information from the parents, the symptoms would have been enough to have created a strict inquiry, and some distrust as to the healthy state of brain. The value of sick-headache in children cannot be overrated; and changes in the organs of vision, as squinting, amaurosis, unequal contraction, or under contraction or dilatation of the pupil, are amongst the most important special symptoms indicative of cerebral mischief. A proof of the importance of the pulse was afforded me some short time since by a medical friend, a careful observer and investigator of disease. When called in consultation to see a little patient, "it was doubtful," he said, "at first to determine whether the child was suffering from bowel remittent fever, or whether the symptoms were produced by some obscure brain affection; but whilst investigating the case, I kept my finger on the pulse, and after two or three minutes I perceived an intermission; that circumstance guided my diagnosis; brain disease was afterwards manifested by unmistakable signs, and the child died." I have myself notes of cases where the slow deliberate pulse, occasionally intermitting, but differing from the diseased mitral pulse, in being otherwise regular, has been the chief symptom of commencing cerebral disease; and I would here call to mind the value of the hard, contracted, incompressible pulse of nervous exaltation, as in mania, etc., and the loose, easily compressible, decurtus pulse of nervous exhaustion, as in typhus, and the extremity of phthisis, and some other exhausting diseases. How important is the enfeebled circulation, with the quick small pulse, evidenced in the cupping of the fontanelles, as pointed out by Hilton, in infants when reduced by diarrhoea and other diseases, every practitioner can bear witness to!

CASE II. Cirroid Aneurism. This case, consisting of an unnatural expansion of the termination of the innominate, and the origin of the common carotid, and subclavian arteries, is an example of disease variously described by authors under the names of *aneurisma racemosum*, *aneurisma cirsoideum*, *aneurisma anastomoticum*, and *varix arterialis*. It usually occurs in arteries of the third or fourth order, such as the carotid, temporal, occipital, etc. The subject of the present notice is a lady still living, about 56 years of age, tall, muscular, and well developed, and who, during the greater portion of her life, has enjoyed the most robust health; but within the last two or three years, has been occasionally invalidated by pains in the head, giddiness, and dimness of sight. On one occasion she consulted me on account of her inability to articulate distinctly, and from her incapacity to frame an entire sentence. On a second occasion I attended her for numbness of the right upper extremity, and pricking sensations with diminished sensibility in the leg and foot. These attacks were of short duration, and readily yielded to depletory measures. Being in a comfortable position in life, she is indisposed to exercise, prone to corpulency, and fond of good living; and at the time of this illness, February 19th, I was called to arrest bleeding from the nose, which had continued for three or four hours, the blood sometimes flowing in stream. The external application of cold and astringents, together with active purgation sufficed, and there was no return of the hæmorrhage till the 23rd, when I found it necessary to plug the nares with a sponge, saturated with a solution of tannin. Directly both nostrils were closed, the blood was forced up through the lachrymal duct, and flowed *guttatim* from the right eye. Cold applications were applied, the patient was then undressed and removed to bed, the shoulders being well raised. Whilst undressing her, I noticed a pulsating tumour above the clavicle on the right side; and, observing it more minutely, I found it to be a pulsating, vermi-

form vessel, lying about an inch above the clavicle, and corresponding with the middle third of the subclavian artery. After a rest, and the subsidence of the bleeding, I made a careful examination, and found the heart in sound and rhythm, healthy; the chest clear on percussion, a systolic *bruit* not very loud being audible under the sternal articulation of the right clavicle. The innominate could be felt distinctly pulsating at the top of the sternum. The carotid was dilated and removed from its position to the anterior triangle, assuming the shape of the italic letter S. The subclavian was also dilated, and on manipulation, the first sensation was that of a globular body; but on carefully examining the tumour, the arteries could be easily separated. I was told on inquiry that she had felt for some time past a throbbing on the right side, especially if she had been excited, or after taking exercise, and for some time she has had a choking sensation, particularly when drinking fluids.

Cases of this description are usually chronic, increasing slowly; and mostly lying quiescent, are a source of discomfort, but not necessarily of death, for years. The coats of the artery are equally expanded through the whole circumference of the vessel, and, as in the case I have related, the vessels are generally tortuous, analogous to what is commonly seen in varicose veins. The treatment consists in lessening the force of the heart's action, giving support to the enlarged vessels, by pressure, and improving the plasticity of the blood and tone of the vessels by diet, steel, and exercise.

This case is interesting as affording a solution of the grave symptoms which marked the previous attacks of illness; and it was a satisfaction to me to be able to find in the diseased vessels a source from which proceeded symptoms which would have been yet more alarming if I had not been able to assign the exciting cause. Had true aneurism existed in the place of the fusiform, we might have had an exhibition of somewhat similar symptoms, but then the paralytic threatenings would have arisen from the laminae of coagulated blood diminishing the calibre of the vessels, and the brain, instead of being congested from overgorging of its vessels, would have atrophied from the diminished supply of blood.

Arteries of smaller dimensions are more generally affected; but it is not uncommon to find the aorta itself expanded, and generally in connection with insufficiency of its valves. This condition produces a character of pulse almost diagnostic in itself. When the valves from deposit are thickened and rendered more unyielding, the blood ejected from the ventricle falls back upon the heart, thereby necessitating two conditions, viz., a larger quantity of blood, and an increase of power; thus we have hypertrophy, and dilatation of the heart, and with this extraordinary increase established to overcome what is virtually an obstruction, the blood is thrown with great force into the aorta, and the latter contracting from its over dilated state, with greater resiliency produces the full, strong pulse, we easily recognise as symptomatic of hypertrophy of the heart. But it is impossible that the aorta can be constantly over dilated without losing its elasticity and tone; it becomes more yielding and less contractile, and the blood is then forced by the heart through it, as a mere passive tube, into the smaller arteries, which, after a time, follow the same degeneration as the aorta. We fail then to appreciate the interval of time between the heart's systole and the pulse; the jet of blood is instantaneous, and as the arteries lose their tone and elasticity, they are capable of a certain amount of elongation, the pulse then becomes full, but not strong, *jerking*, and *visible*, and this character alone will almost designate aortic valvular insufficiency.

CASE III. *Chronic Inflammation of the Aorta, with Ulceration, and Deposit of Calcareous Matter.* This specimen of diseased aorta was taken from a lady past the prime of life, always of a delicate constitution, and fat without power. For some years prior to her death she suffered from what she was pleased herself to call, "spasmodic asthma." Physical examination left no doubt about the state of the large arteries. From the tattered internal lining, and the roughened edges of the ulcerations, owing to albuminous and calcareous deposit, the *bruit* was loud, grating, and whirring, her pulse was small, and unsteady, and she suffered from severe pains of the chest, feelings of impending death, sudden and prolonged faintings; and from being cautioned against all excitement, and keeping within ready access ether, ammonia, and brandy, she, no doubt, warded off repeatedly the death which at last overtook her.

The heart was found dilated, slightly thickened, and soft; and the aorta and large arteries, with degeneration of the coats, thickened, and the internal lining ulcerated, in patches, and

thickened by deposit. I introduce this specimen as showing a condition from which aneurism sometimes takes its origin, and I am disappointed in not having this evening a heart and aorta similarly diseased, where, in addition, the blood had ploughed up the inner and middle coats, distending the external one into an aneurismal pouch.

When the aorta is much diseased, and the heart enlarged and softened, the pulse is generally small, and its character undecided, wavering, with now and then an attempt, as it were, to develop a right pulsation. When the heart is of normal power, or hypertrophied, from the resisting state of the arteries, the pulse is hardened and prolonged; sometimes from atheromatous deposit the arteries are rendered inelastic, and the stream of blood passing through these contracted and unyielding channels, requires a longer time for its transmission, and produces the pulse we have described as *wiry*; and when the deposit is irregular, and perhaps with more of the animal than the earthy matter, the arteries have unequal expansions and contractions, thereby producing a series of little waves of blood in quick succession, giving rise to the character of pulse described as *beady*.

I submit this paper, imperfect as it is, to your consideration, feeling the time will not be lost in calling your attention to the deep importance of "special symptoms." For myself, I can aver that, in consultation, my doubts have been cleared up, on more than one occasion, by the overwhelming evidence of some one symptom to which I had not attached its full value.

ANALYSIS OF 2157 CASES OF LABOUR.

By H. W. T. ELLIS, Esq., Crowle, Lincolnshire.

I SEND the analysis of 2157 cases of labour, which have occurred in my practice as well as in that of my brother, and of which a record has been kept.

Number of cases	2157
Twin cases	29
Total number of children { Males 1119 Females 1067 }	2186
Presentations:—	
Natural	2115
Breech	27
Foot	17
Arm	9
Funis	5
Face to pubes	13
Still-born and premature	71
Illegitimate	72
Placental presentations	7
Forceps cases	9
Acephalous fœtus	1
Hydrocephalic head (in one case breech presented)	5
Spina bifida	3
Talipes varus	5
Hare-lip (single and double), cleft palate and nose	7
Supernumerary fingers	3
Imperforate anus	1
Maternal deaths from causes connected with labour:—	
Puerperal convulsions (death on third day)	1
Puerperal fever	2
Retained placenta (death in three weeks)	1
Exhaustion from difficult labour, etc.	6
Hæmorrhage from placenta prævia (turning)	3
Typhoid fever (death in fifteen days)	1
Paraplegia and paralysis of bladder (survived one year)	1
Maternal deaths from causes unconnected with labour:—	
Phthisis	4
Erysipelas of pudendum and nates in mother (recovery)	1
Polypi uteri	1
Severe hæmorrhage (recovery)	16
Chloroform was used but once.	