cover a multitude of sins, and cloke diseases to which it has no resemblance. Let us be careful, whilst we seek to suppress quackery, we do not permit any fashionable title to be used, on *medical authority at least*, which the pretender might turn as a weapon against us, with the rebuke "tu quoque".

I would offer these remarks with much diffidence, simply adding,

"Si quid novisti rectius istis,
Candidus imperti: si non, his utere mecum."

PATHOLOGICAL CONTRIBUTIONS TO MEDICAL JURISPRUDENCE.

By WILLIAM BOYD MUSHET, M.B. Lond., late Resident Physician at St. Marylebone Infirmary.

II.—OBSCURE CEREBRAL SYMPTOMS (POSSIBILITY OF VIOLENCE OR POISON?) SATISFACTORILY INTERPRETED, ON AUTOPSY, BY NATURAL DISEASE.

F. W., aged 52, a gentleman of eminence in the scientific world, was brought to the Infirmary about 1 A.M. on October 6th, 1856, having been found insensible by the police. He was completely unconscious to external impressions; the pupils were contracted and immoveable to light; the face was pale; but the conjunctivæ were much injected. The respiration was performed at long intervals, and was of a sighing character, not stertorous. There was continual nictitation. He had no deviation of mouth, or other paralysis; no rigidity, nor convulsion. The pulse was full and laboured, but compressible; the surface was cool; the feet cold. A soft systolic murmur was heard at the heart's apex. No fracture was discoverable. There was no smell or suspicion of opium. An abrasion existed over the left malar eminence; and there was tumefaction of the upper lip, from extravasation of blood into its substance. He was ordered a drop of croton oil, sinapisms to the calves, hot bottles to the feet, and a blister to the nape; the head to be moderately raised.

I remained with him about an hour and a half, when I retired to bed, as no alteration had occurred. At half-past 8 A.M. he was breathing stertorously, which, the nurse informed me, had a short time before supervened; and there were slight convulsive twitches of the left side of the face, and mucous râles in the throat. The pulse was small and thready, and the surface cool and moist. Wine was ordered, but he could not swallow; and he died at half-past 10, nine hours and a half after admission.

The body was examined for inquiry two days after death. The weather was warm. Rigor mortis was slight in the upper limbs, more marked in the lower. The pupils were moderately dilated. The face was calm. There were marks over the left cheek and upper lip corresponding to the injury noted during life. The body was well nourished.

Head. The scalp was pale. The dura mater was natural, but a little to the right of the longitudinal sinus, about its middle, was a small osseous body (*glandula pachioni*) of the size of a small pea, which occupied a depression in the cerebral substance. There was some opacity of the arachnoid, and some subserous effusion. The substance of the brain was a little congested, and presented unusual (ramiform) injection. The section of the brain exhibited more *puncta vasculosa* than natural, and its consistence generally was below the average. Much fluid was found in the lateral ventricles. The central portions of the brain were much softer than in health. The fornix throughout its extent was very soft, almost flaccid; as was also the hippocampus major of the right side. The thalami and corpora striata were also less consistent than normal. On careful section, a small clot, not more than a drachm of effused blood, was discovered in the right anterior lobe of the hemisphere, anterior and external to the extraventricular portion of the corpus striatum, and encroaching on the grey matter at the base of the brain. At this spot, the surrounding cerebral substance was softened and disintegrated. The vessels of the brain appeared healthy. About an ounce and a half of reddish serosity was found at the base. The cerebellum and other parts were natural in appearance. There was no fracture of the calvarium, or of the base of the skull.

Chest. The heart weighed nine ounces and a half: it was apparently quite healthy. The right side was flaccid and contained dark coagula. The left side was contracted; its cavities

were empty. There was slight atheroma of the coronary arteries. No deposit was seen on the valves; and the mitral appeared to be perfectly competent. About an ounce of serum was found in the pericardium. The lungs were remarkably congested, but crepitated well. The lining membrane of the bronchi and their divisions was dusky red; and there was much frothy mucus in the air-tubes. There was slight tubercular deposit at the left apex. No pleural adhesions were present. About three or four ounces of reddish serum were present in each side of the chest. A great amount of frothy serum escaped on section of the lungs.

Abdomen. The stomach was very large, distended with dirty looking fluid, of slight sour smell. The mucous membrane was pale, not softened. The rugæ were prominent; and there was some fulness of the veins near the *cul-de-sac*. The spleen was small, pale, soft. A gall-stone was found in the gall-bladder, of about the size of a very small marble. The liver was pale at various spots; and small whitish fibrous nodules, of very firm consistence, were scattered over its substance, chiefly affecting the surface. When the left kidney was handled, a large stone was readily discernible through its walls, imparting the sensation of a solid body in a bladder of fluid. On section, a very large irregularly shaped lithic acid calculus was exposed, which was found to branch and extend firmly into the calices, which were atrophied and absorbed, so that at points the capsule alone intervened between the concretion and the surface. The structure of the organ was very pale; it weighed six ounces after extraction of the stone. The right kidney was large and pale; it weighed five ounces and a half; it was very firm, not granular or lobulated: the capsule separated well. One or two small spots of commencing cystic condition were observed. The intestines were natural. The bladder was distended. There was no stricture of the urethra.

REMARKS. In this case, the objective phenomena pointed to cerebral lesion of some obscure kind. The depression, infrequent respiration, and diminution of temperature, were suggestive of concussion; but the laboured, though compressible character of the pulse, the immobility of the pupils, and the perfect insensibility to external impressions, led me to infer that some condition must exist *plus* the concussion (if concussion were at all a contributing agent) to account for the gravity of the symptoms; as concussion *per se*, sufficiently violent to induce so strongly marked results, would express itself by greater prostration and coldness of the surface, almost total obliteration of pulse, and sympathetic action of the stomach. The symptoms, again, were unlike those of apoplexy, poisoning, or fractured skull.

I felt, therefore, inclined to regard the case as one of softening *plus* concussion; and I think it must be admitted that the *post mortem* appearances ratified the diagnosis. Some of these appearances deserve notice. The evidences of asphyxia were very marked, more marked and characteristic than I ever observed in any case of drowning or death from pure apnoea. The clot was on the opposite side to the external injury, and occupied a very unusual position. This clot seemed almost too small and circumscribed to have determined fatally; but we must bear in mind the extensive softened condition of the brain, the clot superadded, and probably concussion, which might have caused the clot; or the clot, occurring from morbid causes, might have induced vertigo, the fall, and consequent concussion.

The mitral murmur is a point of interest. It was observed in the recumbent position only, the state of the patient forbidding the upright posture. No lesion of the heart or incompetency of the bicuspid orifice was discoverable, although attention was especially addressed to this organ. Two or three similar cases have at times come under my observation, in which a systolic murmur *at apex* has existed, in absence of any appreciable pathological change. I regard the morbid sound detected in this patient to be due either to disturbed innervation, the consequence of the coma, producing irregular contraction of the fleshy columns, or, which is more probable, to deficient lentor, or altered mobility of the blood arising from the cystic condition of the kidneys, which must have greatly interfered with their depurating function.

The sudden invasion of the fatal seizure was a remarkable feature in the case. It was elicited from the relatives at the inquest, that no fit or other indication of disease of the brain had previously manifested itself, except at times a certain alteration and peculiarity of manner.