I am fully aware that many of the symptoms recorded in the above case are at variance with those described by most authors as pathognomonic symptoms of lead-poisoning. There was no wasting of the muscles; there was no lead-colic; no constipation; there was no dropping of the wrist; there was no atrophy of the ball of the thumb; the motor nerves were not the nerves affected (as they usually are in lead-poisoning), but the nerves of sensation; the blue line of the gums, if not altogether absent, was, at all events, very faint. Notwithstanding all these discrepancies, the history of the symptoms; the debility and thorough prostration, both bodily and mental, occurring in a strong and healthy man, without any assignable cause; the loss of memory; the numbness of the right arm and leg; the hair falling off in round patches; the anxiety and palpitation of the heart on the slightest exertion, the heart itself being perfectly free from organic disease; the satisfactory proof of the presence of lead in the water drank, which water the patient had been drinking for years—for nearly fifteen years; and the rapid and favourable termination of the case, simply from substituting pure water for the water impregnated with lead, without giving a particle of medicine, and without in any other particular changing the patient's diet, air, or habits; the bald places on the head being again covered with hair on the change of water,—leave on my mind no doubt that my patient's blood had been poisoned with lead—that it was a genuine case of lead-poisoning.

Luckily, the true nature of the case was found out in time; otherwise, probably, the more usual symptoms of lead-poisoning, as described by authors, might have supervened.

I think it is a recognised fact, that lead acts deleteriously on the human frame, as a blood-poison. We all know that blood-poisons (such, for instance, as scarlatina blood-poison) act in different constitutions in different ways, sometimes affecting one set of organs or nerves, at other times another.

CASES OF NERVE-DISORDER.

By C. HANDFIELD JONES, M.B., F.R.S., Physician to St. Mary's Hospital.

[Concluded from page 310.]

SERIES III.

CASE IV. M. P., aged 40, male, brushmaker, was admitted December 6th, 1858. He had been ill two months. He has no strength in either leg; the left is weakest; he drags it in walking. His feet are cold, and he has "no feeling in them". He cannot walk a quarter of a mile. He has much pain in the loins, but no tenderness is manifested on percussing the spine. He walks with the aid of a stick. The left leg jumps and twitches very much at night. The arms are strong. He has no stricture. The urine is of normal appearance. The bowels are habitually loose. The pulse is very feeble; the skin cool; the appetite very good. He has no thirst. He is not anæmic. He has no pain in the head, but is giddy at times. He has always been very subject to catarrh. He is exposed to cold in a draughty workshop. Ten years ago, he used white lead and potash for washing brushes; his hands were then contracted and numb, so that he could not write. His legs were not affected then. His hands have been well for eight years; and he has left off the use of lead for seven years, and does not now use it in any way. His illness came on gradually.

℞ Strychniæ gr. 1-20; tincturæ ferri sesquichloridi ℥xx; aquæ ʒss. M. Fiat haustus ter die sumendus.

December 13th. He walks better; the "medicine has taken the pain out of the back". The medicine was repeated, and a pitch plaister was applied to the loins.

December 23rd. He has had pain in the calves of the legs the last week; he walks much better. The left arm is numb from the elbow to the fingers along the fore part.

℞ Strychniæ gr. 1-16; tincturæ ferri sesquichloridi ℥xx; acidi hydrochlorici ℥i; ætheris chlorici ℥xv; aquæ ʒi. M. Fiat haustus ter die sumendus.

December 30th. He is much better. He has had pain in the left knee the last three or four nights. The skin is cool; the pulse very weak. There is no blue line on the gums. He was ordered to take five grains of disulphate of quinine three times a day; and to apply to the knee an embrocation of three ounces of turpentine liniment and one ounce of tincture of opium.

January 3rd, 1859. He is improving.

January 27th. The numbness has left the arm; the knees alone feel weak.

℞ Ferri et quinae citratis gra. x; misturæ nucis vomicæ ℥x; aquæ ʒi. M. Fiat haustus ter die sumendus.

February 3rd. He has been galvanised with advantage. He walked ten miles three days ago.

REMARKS. The last two cases are very analogous, showing the occurrence of both motor and sensory paralysis, and its removal, under conditions which seem to exclude the possibility of organic alteration.

CASE V. C. C., aged 10, a male, was admitted November 22nd, 1858. He has been poorly the last three weeks, and became worse yesterday. He complains of pain in his head. Last night he was very restless, and brought up about a breakfast-cupfull of blood by retching. He has no appetite; eats nothing. The epigastrium is tender; the face pallid; the tongue moist, slightly white; the bowels open; the skin cool; the pulse very weak. There is clear natural breathing all through the chest. In the consulting-room, he fell down in a fit, struggling very slightly; he scarcely lost consciousness. He was ordered to have a sinapism applied to the epigastrium every other night.

℞ Ferri et quinae citratis gra. v; tincturæ nucis vomicæ ℥v; aquæ ʒss. M. Fiat haustus ter die sumendus.

November 25th. He has lost his speech entirely; he had two fits on the 22nd. The bowels are open. The medicine was continued.

November 29th. He is better; his voice fails towards night; the pupils are very large. The mixture was repeated, and he was ordered to have a drachm of cod-liver oil three times a day.

January 27th, 1859. I saw him to-day, quite well. His mother informed me that he went to Uxbridge, and returned quite recovered. He has occasional low fits.

REMARKS. This youth appeared to have been stricken suddenly by some debilitating influence: he had, however, no actual palsy, except, for a short period, of the hypoglossal nerves. The hæmatemesis I should ascribe to a paralysis of the gastric vaso-motor nerves, rather than to an ulcer. The fits seemed to indicate a condition of cerebral disorder, not so much pure debility as debility mingled with irritation. The result forbids the idea of organic disease.

CASE VI. W. H., aged 16, male, slater, was admitted June 21st, 1858, having been ill two or three months. He feels giddy; "can't talk as easy to himself as he used." He is always very heavy and sleepy. The tongue is clean; the urine free; the bowels open; the pulse quiet. He is rather anæmic. He was ordered to take one-twentieth of a grain of strychnia three times a day.

June 28th. The giddiness is better; he has pain over the left eye. The medicine was repeated, three grains of citrate of quinine and iron being added to each dose.

July 28th. He was discharged cured.

REMARKS. This was a slight case, but nevertheless phenomena of cerebral debility and lingual paresis were distinctly marked. It is easy to see that a greater degree of enfeeblement would have produced unconsciousness and loss of speech. These slighter cases seem to me to throw considerable light on the nature of more serious ones.

CASE VII. M. S., aged 20, female, servant in a good house, was admitted November 15th, 1858, having been ill two days. She had a fit at first, and nine or ten since, but not "insensible ones". They seem to have been of convulsive character, and only the left side was affected. The tongue was bitten. One fit occurred three years ago. She has some frontal pain, which is better when she is recumbent. She has felt giddy lately, not weak. There is no rheumatism. The skin is cool; the pulse weak; the tongue natural; the catamenia regular; the bowels open; the urine clear. A blister was applied to the neck.

℞ Ammonia sesquicarbonatis gra. v; infusi cascarillæ ʒi. M. Fiat haustus ter die sumendus.

Nov. 28th. She has shaking fits in the left side, and temporary loss of use in the left arm and in the leg; "it comes all numb, and keeps jumping." The fits occur frequently, more so at night. The bowels are open. She has palpitation of the heart. The catamenia are present. The urine effervesces a little with acid; it contains a slight trace of albumen. She was ordered to drink porter, and to leave off tea.

℞ Ferri et quinae citratis gra. x; tincturæ nucis vomicæ ℥xij; aquæ ʒi. M. Fiat haustus ter die sumendus.

November 22nd. She is better, and has no fits. She likes her porter. Pulse quite weak. The arm is not convulsed. The mixture was repeated; and she was ordered to take five grains of blue pill with colocynth occasionally.

December 6th. She has had no more fits, but has a bad cough. The mixture and pills were repeated.

December 20th. She was discharged well; but was directed to continue the mixture and pills a week longer.

CASE VIII. E. P., aged 9, female, was admitted October 8th, 1857. She has been ill three months "with fainting fits"; but, during the last fourteen days, has had "struggling ones". The fits begin with drowsiness: she lies down and goes to sleep; in about ten minutes, the fit comes on with struggling and screaming; she complains that her head is coming off; she loses consciousness for three-quarters of an hour. She had two fits yesterday in the afternoon. The head is not hot; the urine is rather pale; the tongue clean; the skin cool; the pulse weak; the bowels costive. She is restless at night, talking and tossing about. Her appetite is not good; she will not take meat. She has a discharge from a scrofulous sore on the foot, connected with diseased bone. Before the fits came on, she used to have abdominal pains like cramp; these have ceased. The treatment employed was citrate of iron and quinine, and cod-liver oil, with tincture of nux vomica during the latter six weeks. Under this she improved greatly, was reported "not like the same child", and the fits ceased. A blister was once applied to the neck, but it did not seem to do any good. About six weeks before her discharge, the following note was made:—"She appears as if her intellect was affected, is unaccountably lively after the occurrence of pain in head, which comes on generally about 6 p.m., and is attended with great heat in forehead." Two years previously, this girl had been under my care with enlarged cervical glands, and she is again at the present time affected in the same way, but has no convulsions.

REMARKS. The antecedents of this case were such as to raise serious questioning whether the symptoms might not be owing to the presence of tubercle within the cranium. It seemed, however, quite possible that they might result from some unknown morbid influence on a weakly nervous system; and I was the more inclined to this more hopeful view, from experience of analogous states. These two last cases illustrate the occurrence of convulsions, as a form of nerve-disorder, in circumstances very similar to those under which paralysis or failure of power is observed; and also their cure by the same kind of treatment as has been shown to be beneficial in paralysis. If it should appear to any one that this last case was an instance of ordinary epilepsy, I must reply, that the nocturnal restlessness, the periodic head-pain, and the result of treatment, seem to me to make an essential difference. Moreover, the mere name of epilepsy, without any causal indication, explains nothing. I will admit that the convulsions were epileptoid, but I conceive that the case may be ranked with the foregoing, in which some causal influence may probably be traced.

CASE IX. W. H., aged 65, a coachman, was admitted December 7th. He was suffering with his usual winter catarrh, which he had had the last six or seven years. There was no febrile movement. On December 31st, he was suffering from diarrhoea and increased bronchitis. He improved under the treatment adopted till February 21st, when he was attacked with hemiplegia of the left side as he was sitting at home. The arm was most, and the leg but little affected. After this attack, he lost his cough and expectoration almost entirely. He was admitted into St. Mary's Hospital February 26th, and took five grains of sesquicarbonate of ammonia three times a day for twelve days, and then one-twelfth of a grain of strychnia three times a day. On March 16th, when I saw him, he had some difficulty in speaking; but the arm and leg had almost entirely recovered from the paralysis. While asleep, the right cheek was puffed out in expiration; but when he was awake, both cheeks acted well.

CASE X. Anne D., aged 68, needlewoman, was admitted March 8th, 1858. She had been ill eight days. She had had a slight fit, which took her senses away for some good while. The left arm is now paralysed in such wise that she cannot bend or extend the hand; but it does not hang motionless; she has use of the upper arm. The hand feels numb and weak; and pain extends up her arm, especially at night. Her speech was very much impaired for some days; she can put out the tongue straight now. The bowels are open. There is no febrile action, no pain, or heat about the head. The urine is thick, but not red. She is thin and feeble. She was ordered to take five grains of sesquicarbonate of ammonia three times a day, to which a drachm of tincture of cinchona was added after a week; and she took a mild aperient occasionally. By March 29th she was well enough to be discharged, having im-

proved steadily from the first. She is now again under my care, but has had no return of the paralysis.

REMARKS. The age of these two last patients would lead, *a priori*, to the belief that the cerebral disorder depended on some actual lesion of structure, probably on giving way of some small blood-vessels. Yet the treatment that was found beneficial almost forbids the idea that such was the case. The two cases, though furnishing no conclusive evidence, are yet very suggestive. What would have been the result of purging, blistering, and a course of mercury?

Gulstonian Lectures

ON

FEVER AND INFLAMMATION.

DELIVERED BEFORE THE

ROYAL COLLEGE OF PHYSICIANS, LONDON,
1859.

By WILLIAM ADDISON, M.D., F.R.S., Fellow of the College.

LECTURE I.

MR. PRESIDENT AND GENTLEMEN,—If we carefully examine into the causes which produce disease, it will be found that, in a great majority of instances, they may be resolved into some violation or neglect of the natural laws upon which the healthy actions of the animal frame depend. So also, on the other hand, the means of cure in such cases will be found to consist in aiding and adjusting natural efforts to reestablish those laws and conditions which have been outraged or infringed. Let us observe what these natural efforts are.

In the repair of injuries to the human body, certain parts are healed by the growth or production of elements identical with those of the wounded texture. On the other hand, there are organs, injuries to which are repaired, not by the growth of elements identical with those composing the injured organ, but by elements of the same kind with those which are effectual for cure in the former cases. The result is that, in all examples, and in very different parts of the body, the elements of repair appear to have a common form or nature. Now, the parts which are restored by the production of elements identical with those composing the injured texture, are such as have the most general distribution in the body, and are ministering to the purposes of life in the lowliest or simplest manner.

For example:—Fibrous and connective tissues and blood-vessels are everywhere present; no part of the body can be said to be without them; all these are speedily reproduced or repaired when injured or broken. Tendons, ligaments, and bones, are of very general distribution: these also are repaired by new growths of tendinous and osseous tissues. But when we come to investigate the healing process in more special organs, it assumes a different aspect. For, if a portion of the muscular flesh be rent away by an accident, cure is effected, not by the growth of new muscular fibres, but by a growth of common fibrous tissue. So likewise, if a part of the lung, of the brain, or of the liver or kidney, be destroyed by accident or disease, cure, if effected in these instances, is so, not by a growth of the special parenchyma, but by a common fibrous tissue.

Upon this ground, then—namely, the common form of elements of repair—we purpose to review the human body as a structure composed of three orders or classes of parts: a common tissue; the blood; and parenchymatous organs.

I.—A COMMON TISSUE: ITS SEVERAL FORMS AND USES.

The areolar tissue is a common tissue diffused throughout the whole body. It invests the exterior, and enters with the blood-vessels into the interior of every organ. Its elements everywhere incorporate with the coats of the vessels. And in all fibrous membranes the elements of the membranous tissue between the blood-vessels are of the same kind with those forming the coats of the vessels.

In natural growth, in the embryo, areolar tissue is at first composed of cells and nuclei; and the coats of the blood-vessels, and those parts of the body which are hereafter to become tendons, ligaments, and bones, consist in the embryo of cells and nuclei also. As growth proceeds, the areolar tissue becomes fibrous; and, *pari passu* with it, the coats of the