dwellings, and in crowded cities; in clean and well drained places, and the reverse. Therefore, ought we not to look to the atmosphere as producing those rapid changes, acting upon the blood, occasioning the gangrenous condition of the throat, accompanied with an extremely low and depressing type of fever, from which the patient rapidly sinks into a state of extreme and fatal exhaustion? It therefore appears consistent with pathology to administer a most powerful antiseptic remedy, to act promptly both locally and constitutionally. After taking the above view of the disease named by Bretonneau *diphtherite*, I beg to submit to my medical brethren the plan of treatment which, I am happy to say, I have found very successful.

I had under my care several severe cases in the autumn of 1857, and again in the autumn of 1858. Those in 1857 occurred in the town and neighbourhood of Croydon, and those in 1858 in a new building, an asylum for the reception of children whose parents died young: consequently, their offspring were not constitutionally the best fitted to resist so formidable a disease.

The treatment I adopted in all the cases under my care was as follows—for children:—

℞ Solutionis chlorinii ʒss; syrupi simplici ʒss; aquæ destillatæ ad ʒvj. M. Fiat gargarisma sæpe utendum.

℞ Solutionis chlorinii gtt. iv; syrupi aurantii ʒj; aquæ destillatæ ad ʒss. M. Fiat haustus 2ndâ quâque horâ sumendus.

The dose was increased according to age. Calomel was given in doses of one grain and upwards, according to age. The diet, too, consisted of concentrated jellies, strong beef-tea, wine, etc.

The same mode of treatment was adopted in adult cases; except that, instead of calomel, I gave the hydrargyrum cum cretâ.

For many years I have ordered the chlorine solution in malignant scarlet fever accompanied with a diphtheritic state of the throat, with marked success. In the *Cyclopædia of Practical Medicine*, published in the year 1835, Dr. Tweedie states that, in an epidemic sore-throat which made its appearance in Tours in 1818, and to which Bretonneau gave the name of *diphtherite*, hydrochloric acid was found most efficacious.

The orphan asylum to which I have before alluded is a new building erected on the top of a hill, open to the south-west; the rooms spacious and lofty; every attention paid to ventilation. It stands alone, in a most healthy situation; and the children, before admission, undergo a medical examination; so that, at the time of admission, they are perfectly healthy. So that does away with the supposition of its being confined to ill-drained, low, and swampy situations, and densely populated, poor localities. In the months of July and August, 1858, there were fifty cases in the above asylum, of the most malignant form. They were all placed under the treatment before named. After the third day, quinine was added to the chlorine solution. Many were covered with the scarlet eruption; eighteen had a severe form of measles; and some few were free from eruption; but all had the diphtheritic throat, and but one case out of the fifty proved fatal—a child four years old. The disease made its way through the various tissues of the throat and neck, and broke externally, giving vent to a most offensive discharge, the patient sinking from exhaustion, not being able to take sufficient stimulant and nourishment.

DIPHTHERIA.

By J. C. S. JENNINGS, Esq., Malmesbury.

THERE are a few points in the summary of cases published in the *BRITISH MEDICAL JOURNAL* of June 25th, to which I wish to advert, while completing a tabulated report of cases of diphtheria attended by me for the last two years, although I can, in truth, add but little as regards its treatment to the outline which I gave in the *JOURNAL* of March 27th, 1858.

Presence of Contagion. Of this I have had such clear proof ever since the first outbreak of diphtheria in the parish of Shipton Moyne, that I have had no hesitation in declaring its contagious character: in fact, in the first case which came under my care in that parish, the disease was clearly contracted by nursing and attendance on an adult who rapidly sank under it. Moreover, two of his brothers-in-law, residing a mile distant, contracted the disease either from a casual visit, or by attending the funeral.

Scarlatina. At the first outbreak of the disease, no cases of the above had appeared in the neighbourhood; nor were

there any until the second outbreak, during the month of January of this year, when a few cases of diphtheria occurred; but scarlatina maligna ran through several families. In those cases, however, in which the rash was well developed, and not suppressed, there was little or no throat-affection; and *vice versa*: and when the tonsils were affected, there was not the peculiar leathery exudation of diphtheria.

Treatment. The plan I have invariably adopted, regardless of sex, age, or incubation of disease, has been to give an active emetic of antimonial wine, from half an ounce to an ounce, according to age; to freely cauterise the throat with solid nitrate of silver; to have a mustard poultice applied from ear to ear; the feet and legs plunged in a hot bath; and the patient confined to bed. After the emetic action has ceased, from three to five grains of calomel with five of compound extract of colocynt were given (or, for a child, two grains of calomel with two grains of compound antimonial powder); and, four hours afterwards, the following mixture:—

℞ Quinæ disulph. ʒss; potassæ chloratis ʒj; acidi hydrochlorici diluti ʒss; aquæ ʒviij. M. Fiat mistura cujus sumatur pars sexta 4tis horis.

A gargle of chlorine solution was directed to be used frequently, prepared by impregnating water as much as can be borne with the protoxide of chlorine, generated from two parts of chlorate of potass, one of hydrochloric acid and one of water, and the fauces to be sponged out frequently with the same. The emetic I have rarely repeated more than once; but when the inflammatory stage has been severe, the fauces tense and shining, and the throat œdematous, spirit of nitrous ether and liquor of acetate of ammonia, or nitrate of potass, has been added to the mixture.

The diet has been at first farinaceous, and afterwards consisting of strong broths and jellies. Stimulants have been very rarely administered, and then only as sherry whey, alternately with the quinine, which I have trusted to as the sheet-anchor. For infants, quinine may be given in jelly, washed down with a mixture of tincture of sesquichloride of iron.

Upon this plan all my cases have been successfully treated, with the exception of one fatal case in a stout young man, where much valuable time had been allowed to elapse; the mushroom-like exudation having extended over the soft palate, completely blocking up the fauces and chink of the glottis; besides which, the emetic was not rightly administered, or failed to produce effect; neither was the purgative given as ordered; and the patient died from apnoea, suffocation being produced from spasm of the glottis when turning to lie down in bed.

Too much stress cannot be laid upon tartar emetic, quinine in large doses, and the avoidance or guarded use of alcoholic stimulants.

WOUND OF THE LIVER: PROFUSE HÆMORRHAGE: RECOVERY.

By ALBERT G. WALTER, Esq., Pittsburgh, Pennsylvania.

JOHN KOHRSEN, roofer, aged 47, of Pittsburgh, of bilious constitution, large frame, and strong muscular development, was brought to my hospital late on the evening of January 17th, 1858, having been stabbed by his wife in the scrobiculus cordis with a broad-bladed knife, entering the abdomen obliquely upwards about four inches. Withdrawing it himself, he walked some distance while bleeding profusely, and fell down exhausted upon the pavement. A surgeon summoned found a vertical wound, of an inch in length, close to the right side of the ensiform process, from which venous blood, in a rapid, continuous, and large stream, was issuing. On introducing his finger into the cut, a deep rent was discovered in the concave surface of the small lobe of the liver. This profuse flow of blood was arrested by compression through the abdominal walls upon the aorta.

Arrived at the hospital, the patient was cold, very pale, with respiration not accelerated, and slow and hardly perceptible pulse, yet restless, outward bleeding having ceased. He suffered great pain in the wound, increased by swallowing, moving, and deep inspiration; with paroxysms of cramps in the abdomen, shooting from the scrobiculus cordis to both angles of the scapula, and returning every quarter of an hour, during which the skin around the wound would swell out to the size of a pigeon's egg, as in ventral hernia. The wound was left open; the body was inclined towards the right side, to facilitate the outward flow of blood; ice in bladders was applied to the abdo-