With regard to the drachm there will, moreover, be this strange inconsistency: it will be the eighth part of an ounce avoirdupois, and will therefore be exactly twice as heavy as another weight of the same name, viz., the drachm avoirdupois, which is the sixteenth part of the same ounce.

One of the immediate consequences of the proposed change will be, that an entirely new set of brass weights will be required by every general practitioner and druggist in the kingdom; all the old weights will become valueless.

Be it observed, that the inconveniences which have thus been pointed out are not the immediate consequence of the praiseworthy aim of the Committee to get rid of the confusion between the two ancient sets of weights—the troy and avoirdupois pounds and ounces; they are not the necessary consequence of substituting these avoirdupois weights for the corresponding troy weights, but are the consequence of an endeavour to effect the division of the ounce avoirdupois into 480 integral parts. It seems to me that this object is not of so essential an importance as to justify the introducing new elements of confusion into our English system of weights.

It may, perhaps, be worth while to point out a plan by which all these inconveniences might be avoided. This may be done simply by ceasing to use the ounce in prescriptions; of course, I mean the ounce in weight, not the fluid-ounce, which is a measure, and with which it is unnecessary to interfere. Let the quantities of solid substances be prescribed in drachms, scruples, and grains, such as are now in daily use. When larger quantities are required, they can be written in drachms; or, if of the largest kind, may be expressed by the pound of 7000 grains, which is an exact multiple of the drachm.

Perhaps a still better (and the simplest) plan would be, for all quantities less than a pound, to prescribe in grains alone—in standard grains alone. The calculations between large and small quantities would always be easy, because the standard pound is a large round number (7000) of grains. We should have weights of 100, 200, 500, and 1000 grains, such as are now occasionally used in scientific investigations. There would be no difficulty in writing our prescriptions. Five hundred grains (gr. ν) is as easily written as $\mathfrak{z}$. Fifty, or a hundred, or a thousand grains, are likewise easily written, either in the ordinary figures, or as gr. ι , gr. ς , or gr. μ .

As the grain would not be changed, all the brass weights now in use for dispensing might still be used. A single set of the small weights might be made to serve for dispensing any drug not exceeding 250 grains in weight. If any new weights were needed, they would be one of 300 grains, and one of 500 grains; and these would only be required in cases in which ounce weights are now used. No other weights would be needed for dispensing and compounding on a moderate scale; and if larger weights were needed, they would be the ordinary avoirdupois pound, half-pound, and quarter, which are in common use in every shop and most private houses, and which weigh respectively 7000, 3500, and 1750 grains.

As the intentions of the *Pharmacopœia* Committee have not yet been published on authority, my remarks may appear premature. But the intentions are well known; and to wait for their authoritative publication is to wait until it might be difficult or impracticable to have them reconsidered.

ILLUSTRATIONS OF THE USE OF THE OPHTHALMOSCOPE.

By WILLIAM MARTIN, F.R.C.S., late Professor of Ophthalmic Surgery in the Calcutta Medical College.

[Continued from page 598.]

CHOROID.

Congestion of the Choroid may be diagnosed by a hyperæmic state of the chorio-capillary layer. Choroidal inflammation, in various degrees, may then be expected to ensue, and to cause atrophy, maceration of the pigment, layers of exudation, etc. These again, would, in the end, bring on disorganisation of the retina.

In the first instance, we may find an unusually pale condition of the fundus. It does not light up well, and the appearance is rather of a pale orange-yellow than the natural bright pink colour. We shall be likely, also, to find some discoloration of the fundus; instead of a uniform field, we see pigmental patches in parts. If these appearances last long the disease progresses to disorganisation. The effects on vision will be more evident according as this disorganisation—whether it be maceration of the pigment, or anything else—affects the ma-

cula lutea, or its immediate vicinity. The yellowish tint of the fundus becomes of a paler hue. We see the choroidal vessels through the retina, which before only formed part of a uniform red field. After a time we cannot distinguish the vessels.

We may see patches, of a rounded form, arising from the discoloration of the choroid, and want of pigment; at the same time the retinal vessels covering these become more tortuous, and probably varicose.

This state, which is called *maceration of the pigment*, there are strong grounds for thinking invariably leads to atrophy of the choroid and retina. According to Desmarres, the patches arising from disorganisation, the result of choroidal disease, may be distinguished from patches arising from exudation, by being of a rounded form; the exudation patches being almost always of an elongated, or oblong form. The pigment becomes collected into spots, which look dark, and have jagged edges. Between these pigmental spots are often portions, red from extravasated blood. We may have, at the same time, patches from disorganisation and the other form. See *Cases of Subacute Choroiditis*, in Desmarres, at pages 419, and 421.)

Choroidal Congestion is generally serious according to the length of time it has lasted.

Hyperæmia, of an acute character, is a concomitant of most internal ophthalmic inflammations; but is in many cases remediable, and goes off with the cessation of the primary disease. The exudation patches, which are by no means rare in choroidal affections, besides having an elongated form, like that of a fillet or band, present a colour more or less bluish, easily contrasted with the white rounded patches, indicating disorganisation, absorption of pigment, and obliteration of the vascular layer of the membrane.

Sclero-Choroiditis, which is often either a primary disease, or a concomitant of other internal affections, when it has become advanced is known by a peculiar appearance, situated in immediate contact with the optic papilla. We shall find this part surrounded, totally or partially, by a patch, which begins by being semilunar, apparently prominent, but in reality concave. This is often far from being regular in its outline; it is jagged too at the borders; and if the disorganisation has gone on to any extent, it is in turn partly surrounded by a black mark or border, arising from deficiency of the choroid, and accumulation of pigment in this part. As the disease proceeds, this white patch, or arch, extends, forming at length a complete circle, but of irregular breadth, round the papilla.

In bad cases, the papilla looks as if concealed in a large white patch, having the appearance of a double cone; and if the disorganisation proceeds, we shall have the state called *posterior, or sclerotic staphyloma*; when, in consequence of the total absence of the retina and choroid tunics, we see only the sclerotic beyond; or we may see it through a transparent retina. This is shown by our observing the retinal vessels traversing this white patch.

The entire absence of pigment epithelium, and other elements of the choroid, as well as of the retina, which has been noted in these and similar cases, by Desmarres, and other writers, we must remark is not observable in all cases, for Jäger states that in twenty-three instances he found the choroid a continuous membrane. (See Hogg *On the Ophthalmoscope*, page 80.)

Apoplexy of the Choroid. We often find by itself or in conjunction with the deep-seated inflammations, apoplectic or bloody effusions. These may be in the form of a red patch, which may evidently raise the retina; there may be several of these, of different sizes, scattered over the fundus, and they may traverse the retina and pass into the vitreous humour, where they may become diffused, or appear as distinct clots. These clots are capable of absorption, and the time occupied by this process varies much in different cases. Some are very rapidly absorbed. While undergoing this process, they lose their bright red appearance, and become more or less brown; but those which remain unabsorbed a long time, often retain their primitive vivid red colour and character. As in other local changes in the fundus, the impairment of vision altogether depends upon the situation of these deposits. If they involve the immediate vicinity of the macula lutea, vision will be very much affected; but considerable bloody deposits may exist in other parts, and be more or less permanent, without materially impairing vision. Intermixed with these collections, we often see the white patches which indicate the disappearance of pigment. (See Desmarres, vol. iii, p. 437, in which a bloody patch is seen, having its original appearance, after ten months duration.)

Atrophy of the Choroid is a concomitant or consequence of

many of the internal inflammations, and is always a grave symptom. The ophthalmoscopic appearances differ according to the particular portion of the membrane: if it affects the first layer, the fundus, instead of the usual rose colour, presents a pale orange yellow or brown tint, and perhaps we find brownish patches and streaks, giving the appearance of a scratched picture. (Desmarres.) Secondly, where the arterial layer (*chorio-capillary*) has become affected, we have obliteration of the capillary network, and we see the larger vessels beyond with unusual distinctness. Thirdly, where the venous layer has become involved, we find pigmental deficiencies and accumulations in such a manner, that the fundus has a spotted appearance, like that of a tiger's skin.

RETINA.

In cases of *congenital insensibility of the retina*, we find an anæmic state coexisting with a minute condition of the vessels. This is particularly observable in the circulation at the papilla. In certain cases, an *entire absence of blood-vessels* has been found; and this has been supposed to be congenital (Desmarres, vol. iii, p. 445). In many cases of amaurotic insensibility of the retina, we find a partial anæmic state; and the change is particularly observable in the papilla, which is unusually contracted, and presents the appearance of a pearly whiteness, with vessels few and small, etc.

Atrophy of the Retina is known by similar appearances—the pearly white contracted state of the optic papilla, etc.; a general pallor of the fundus; want of circulation, etc. In such a state, we must expect to find many other signs of severe internal disease and disorganisation, patches of exudation, patches of denudation, etc.

Varicosity of the Retinal Vessels is troublesome as a complication of the severe choroidal and retinal affections. It is said also (Desmarres) to exist in those who suffer from hæmorrhoids or habitual cerebral congestion, without producing any appreciable impairment of vision.

Retinal Patches of Exudation may be either nearly transparent, in which case we see a bluish white tint, scarcely perceptible; or in the form of thick patches, in which we find vessels of new formation, the original vessels being lost. The new vessels are said to be recognisable as being contrasted in direction with that of the normal vessels.

Hyperæmia of the Retina is frequently seen as a consequence of acute or chronic inflammation. It is recognised by the unusually brilliant colour of the fundus, and the want of the usual distinguishing clearness of the papilla, in consequence of its becoming masked by vascularity. It seems to be blended altogether, or in great part, with the rest of the fundus. Sometimes it is so red as to resemble pannus of the cornea, or it may present the appearance of fascicular redness, similar to what we see in pustular conjunctivitis, etc.

Chronic Retinitis. We find a general obscurity of the fundus, arising either from exudation of some kind, or simply congestion, or possibly alteration, of the choroid tunic beyond. This obscurity is to be distinguished from that of the vitreous humour by its not being nearly so perceptible at a distance from the optic papilla (as in the direction of the ora serrata), and from its evident existence often also, but not invariably, behind the vitreous body. The obscurity arises from light being reflected in less quantity from the fundus. We often see also the borders of the papilla irregular and ill defined; the rest of the fundus is sometimes overlaid with regular stripes of a dull red colour. The vessels, instead of pursuing a regular course to and from the papilla, sometimes disappear, and then after an interval reappear. This is a phenomenon often seen in œdema of the retina also. The vessels, at the same time, are more developed than usual, and the minutest ramifications come into view. Under this head, Mr. Hogg relates a case at page 133—a case where there was excessive congestion of the fundus, so that the papilla could not be distinguished: the retina was raised by effusion. Also, at page 134, a case is related in which, with this obscurity of the papilla, floating bodies were discerned in the vitreous humour.

Acute Retinitis. In this, as we may expect to be the case, we find various signs of plastic exudation. The fundus is more or less obscure, and looks often as if covered by a bluish grey varnish. Along the course of the vessels on both sides, we may find small white tracts. With this obscurity we may expect to find vision very defective; and it is so, particularly if the exudation or vascular disturbance extends in the direction of the macula lutea; otherwise, the disturbance of sight seems in many cases to be hardly commensurate with the changes which have taken place. See an instance in Desmarres, vol.

iii, p. 465, where, with a considerable patch of exudation (but which, however, does not extend to the vicinity of the macula lutea), the patient has very good sight. The exudation sometimes contains newly formed vessels; more often the normal vessels disappear in certain spots under the patch, to reappear at the farther edge of it, and then to pursue their regular course.

Edema of the Retina. We find true thickening round the optic papilla. The vessels present a convex appearance, and the papilla has an unnatural yellow tint, which renders it less brilliant than usual. This affection occurs as a consequence of choroidal and retinal congestions of a chronic nature, and is a frequent concomitant of syphilitic affections of the posterior tunics. It may disappear gradually, as the primary affection subsides. Desmarres describes the appearances of the œdema which exists in conjunction with syphilitic cachexia as being of a dirty rose-colour, the raised portion of the membrane encroaching upon the papilla, which is always hyperæmic. Mr. Hogg, at p. 137 of his book, relates a case in which the papilla displayed a circular grey ring within the regular margin, with mottling of the fundus; vessels large and irregular. The retina was found in an œdematous state. Also, at p. 138, another case.

This Mottling of the Fundus is a very characteristic appearance. There is a want of uniformity of the red field, which presents the appearance of a pale yellow membrane, covered over with dark blue streaks in very many places. It occurs pretty often, and seems to indicate great disturbance of the chorio-capillary circulation, and possibly obliteration of many of the minute vessels, with a very irregular and deficient distribution of pigment. It is, therefore, a concomitant of many of the more serious posterior congestions or inflammations.

Fatty Degeneration of the Retina has been found in certain cases, but the characters to be discovered by the ophthalmoscope are unsatisfactory. It may be suspected where there is albuminous urine, if at the same time we find defective vision, and the fundus generally obscured, and partly covered with yellowish patches, apparently prominent. (See Desmarres; Hogg, p. 70, etc.)

Apoplexy of the Retina is found, but not necessarily as a concomitant of special diseases, both in the plethoric and anæmic subject; the local circulation being affected in one case with positive, in the other with relative plethora. We see the fundus, perhaps more highly coloured than usual, over-spread in places with bloody clots of very various sizes. These may present the appearance of circles surrounded by other circles, and this is observable in cases of albuminuria: effusions at the central parts of the field have a radiated form. We may be able to see the point of rupture of the vessel, in the form of a dark red patch along its track; but generally we are obliged to wait until some absorption has taken place before we detect the exact point of rupture.

Sanguineous Effusions on the Retina are often observed in acute inflammations of the retina and choroid; also in many states of local and general debility, particularly in conjunction with albuminous nephritis, heart-diseases, amenorrhœa, and hæmorrhoids. In Desmarres, vol. iii, p. 474, is figured a case in which effusion of blood, with great impairment of vision, occurred from a sudden rush of blood to the head; absorption subsequently took place, and all morbid symptoms were removed.

Subretinal Dropsy and Serous Effusions on the Retina. A portion of the fundus we shall see covered over by a floating mass, of a bluish white colour, with vessels upon it or mixed up with it: this will retain its size and appearance a long time. Together with this we may see white folds, appearing as if stretched out from the optic papilla. This mass, with the liquid in it, is seen to be in a constant state of fluctuation, but rests more often at the lowest part of the fundus. Upon it are seen stretched the vivid red lines which are the retinal vessels; sometimes the mass, instead of floating freely, moving in all directions, and then, settling below, becomes fixed to a portion of the choroid. The neighbouring retinal surface having been raised into folds, these folds seem to be pasted together by the effusion. Sometimes bloody matter is mixed with the serous effusion. We may see so large a portion of the retina detached, that it presents the appearance of a funnel stretched from the ora serrata to the papilla. The most striking and characteristic appearance is the floating vessels in the posterior part of the eye. The symptoms are those of amaurosis.

Softening of the Retina. This affection has been described by Dr. Bader in the *Ophthalmic Hospital Journal* for January 1858. He gives some instances, such as—

1. In a case—W. F., aged 34—where there was little but perception of light, both optic papillæ, too, well defined, the surface of the papilla presented the appearance of blue-grey dots in a watery ground; and near its margin, on the fundus, were irregular white patches dotted with pigment; many brown pigment-islands in the field.

2. In another case—E. G., aged 55—in whom there was also only perception of light, the papilla well defined, a portion of it blue-grey, the inner portion presented a dirty white, finely dotted appearance; vessels normal, but unusually curved over a narrow ring, white, which forms its margin.

3. In another case—J. W., aged 57—vision of one eye lost; with the other he can with difficulty manage to find his way about. Optic papilla, too, well defined, and its surface blue-white, dotted on a watery ground. On the side nearest the macula lutea is a bluish grey crescent. On the fundus there are many brownish pigment-islands. Appearances of the two eyes similar.

As the disease advances, Dr. Bader describes the grey-blue appearance more marked; the circulation becomes less; and the vessels seem more bent over the margin of the papilla. Arterial pulsation is often seen. We also often see pigment-islands—that is, the hexagonal pigment layers—instead of the uniform choroidal red. He remarks it as an extraordinary circumstance, that the retina, although in this, as in many other affections, it is liable to become disintegrated, is still a transparent medium. This disease is one of the complications likely to result in acute glaucoma.

Detachment of the Retina. It has been above stated that, among other causes for this condition, may be enumerated subretinal serous effusions; and that it often presents the appearance of a funnel-shaped body, extending from the vicinity of the ora serrata to the optic papilla or other portion of the fundus. Mr. Hogg, at p. 72, relates a case of traumatic amaurosis where, with signs of exudation, the retina was becoming detached round the papilla; and notices cases of Gräfe, in which the retina was affected by various exudations, and had become detached. At p. 128 he also relates a case of Sichel's, in which, with general œdema of the retina, there was a partial detachment of it. The papilla could not be distinguished. Fundus much congested. At p. 131 he also relates a case in which the most prominent symptom was hemiopia. In both regions of the macula lutea were dark spots and pigment accumulations; in the right eye was detached retina.

Crystals of Cholesterine have been found mixed up with exudations of various kinds, as a consequence of retinal disease.

Malignant Diseases of the Retina. In certain cases we find, in an early stage of the disease, a portion, if not all of the fundus, covered by a brilliant coloured prominence, convex in shape, with blood-vessels which have become fixed to the surface, and do not fluctuate as in the partial detachment which arises from subretinal serous effusion. Sometimes, however, both conditions exist in combination (Desmarres).

Abnormal Insertion of the Optic Papilla, without any special disease, is described; but further observation is required on this point.

Anæmia of the Papilla is observed in all cases of atrophy arising from chronic retinal disorders, also as a functional and temporary change in many cases in which there is no organic alteration of the retina, but where the retinal circulation has become interrupted. It is known by the papilla being unusually distinct from the rest of the fundus, the want of blood making it look white; and, besides, we see what vessels there are unusually small and colourless. It seems sometimes to be smaller and more convex than natural, and is often somewhat irregular at its edges; at other times, it seems to be flattened out, and larger than usual. Around it the circulation sometimes has the appearance of being unusually active.

Hyperæmia of the Papilla—a state coexisting with many affections, acute or chronic, of the posterior tunics—is characterised, on the other hand, by a want of distinctness of the optic papilla; it becoming as highly coloured as the rest of the fundus, and often so very vascular that, even in cases where the entire fundus is hyperæmic, it cannot be distinguished, particularly at its margins, from the rest. The chief guide we have in that case is the course of the enlarged vessels. At the same time, we may find, particularly on compression, pulsation of the vessels which are enlarged, isochronous with the pulse at the wrist.

Aneurism of the Papilla has been spoken of by some writers. *Spontaneous Pulsation of the Vessels of the Papilla* is a sign of severe disease. It is particularly noticeable as a sign in

glaucoma, and is not always easily detected. We must observe carefully, to find the movement in the arteries; and this can generally be ascertained to be isochronous with the pulse.

Varicosities of the Vessels of the Papilla are often to be observed, in conjunction with various choroidal and retinal disorders. It is of consequence, as it is, of course, apt to lead to organic derangement; but some observers assert that considerable varicosities may exist without apparent derangement of vision.

Softening of the Papilla is a bad sign, and generally indicates approaching atrophy. The papilla looks unusually large and flattened out, with portions of it puffed; its colour a dirty yellow; borders very irregular; and there is often found, in combination, infiltration of the retina, exudations, etc.

Apoplexy of the Papilla, said to be confined to that part in some instances in which the disorder is not one of the most severe.

Atrophy of the Papilla is a sign of serious disease. It presents an unusually brilliant white appearance, the colour of mother of pearl, and is generally small and arched in the middle; the vessels sometimes small, at others rather dilated, and possibly varicose, seeming to disappear at the edges, and then reappear at another place. In cases in which the diminution of size is combined with unnatural prominence, and seems to arise from compression of the optic nerve at its entrance into the eye, the name of mushroom atrophy has been given. (See Desmarres, vol. iii, p. 515, fig. 66.) In other cases, we have a depression of the optic nerve at its entrance, or cupped atrophy (*A. en godet* of Desmarres), said to be due to retraction of the central fibres of the optic nerve. The essential disease, which must have given rise to this phenomenon, will have advanced to a considerable degree before it is observable.

Fibrinous Degeneration of the Papilla. Mr. Hogg, at p. 101, notices Gräfe's statement, that he has found a case in which the papilla appeared flattened out, with its vessels atrophied. On a *post mortem* examination, the papilla was found to consist of tendinous tissue, with here and there a few nerve-fibres interspersed.

[To be continued.]

Reviews and Notices.

EPIPHORA, OR WATERY EYE: ITS SUCCESSFUL TREATMENT BY THE NEW METHOD OF DILATATION. WITH ILLUSTRATIVE CASES. By JAMES VOSE SOLOMON, Fellow of the Royal College of Surgeons of England; Surgeon to the Birmingham and Midland Counties Eye Infirmary; etc. Pp. 32. Lond.: Churchill. Birmingham: Cornish Brothers. 1859.

THIS is a reprint, with additions, of a very valuable paper which was published some months ago in this JOURNAL by Mr. SOLOMON, and in which he described the results of his application of the plan of treatment of epiphora recommended by Mr. Bowman. The work concludes with some remarks on certain changes which affect the puncta lacrymalia, in consequence of which they may be found patulous, obliterated, contracted, or displaced. We recommend it to the favourable notice of all who have any concern with the eye as operators.

A MEMOIR ON THE TREATMENT OF THE EPIDEMIC CHOLERA. Read before the Members of the French Academy of Sciences: with their Report thereon. By JOSEPH AYRE, M.D. Pp. 44. London: J. Churchill. 1859.

THIS memoir was sent by Dr. AYRE to the French Academy of Sciences, in competition for the prize offered by the late M. Bréant for "A treatment of unquestionable efficacy, which shall cure Asiatic Cholera in the great majority of cases as certainly as cinchona cures intermittent fever." Dr. Ayre strongly advocates the claims of calomel, which he was led to employ in 1832. The author gives his own results, with those of other practitioners: he seems, however, not to have been successful in convincing the members of the section of Medicine and Surgery in the Academy of Sciences, that his treatment stands in the high rank which he claims for it.