was translucent. About half an ounce of clear serum was drawn off with a fine trocar. The fluid contained abundant albumen, contrary to what some authors have affirmed to be the case in hydrocele of the cord. The tumour has not returned.

CASE III. L. D., a gentleman of good constitution, applied to Mr. Barwell on February 8th, 1858, on account of a tumour on the back of the leg. The tumour was found to be situated in the popliteal space, at the side of the biceps tendon; it was of about the size of a large walnut, superficial, painless, rather tense, and fluctuated. It was evidently a synovial sac, probably originally connected with the sheath of the tendon above named. Various counterirritants were applied without benefit; in fact, the tumour continued to increase. Mr. Barwell punctured it, making but a very small opening, squeezed out the fluid (a thin clear serum), and applied pressure. The sac, however, filled again, and the tumour became larger than before. He examined it again most carefully, with the intention of discovering whether any communication existed between the sac and the sheath of the tendon from which he supposed it to originate, and found that such was not the case. He therefore proposed to inject the tumour with a solution of alum; the proposition was accepted, and the operation was performed with the finest trocar. The solution was of the strength of two grains to the ounce; and, after it had remained in the sac five minutes, care was taken to squeeze it out entirely. But little pain was felt at the time. Firm pressure was kept up for three days, and then removed. The tumour has not reappeared.

It may be remarked, with reference to the first and third cases, that Mr. Barwell considers alum solution preferable to iodine as an injection for synovial tumours, since it acts as an astringent, without altering the constituents of the tissues to the same degree, and therefore giving a greater chance of adhesion.

ST. GEORGE'S HOSPITAL.

I. TWO CASES OF FOREIGN BODIES IN THE NOSTRIL.

Under the care of PRESCOTT HEWETT, Esq.

Two little operations, witnessed here the other day, will be found of some practical interest, as showing the length of time during which foreign bodies may remain in the nose without exciting any mischief, and as giving a caution that, in children, we should always examine these cases minutely for foreign bodies when, as is frequently the case, a discharge goes on for any length of time from the nose without any obvious cause, and is not alleviated by the remedies employed. Both the patients were little boys, about six or seven years of age, under the care of Mr. Prescott Hewett. The first came with the history (although, as it afterwards proved, not a correct one) of a foreign body. He was said to have passed a plum-stone up his nose. This was supposed to have taken place a year ago. On examination, some foreign substance could be recognised with the probe high up in the nostril, wedged against the spongy bones. It was impossible to deal with it till chloroform had been given; and, this having been done, a long screw (more than an inch in length) was with some difficulty extracted. The child soon recovered. The other had been under the care of various practitioners for the last three years, on account of "discharge from the nose," "ozæna," etc.; but without any suspicion being entertained of a foreign body being present. On examination, Mr. Hewett recognised at the top of the nostril a black looking substance, which seemed hardly like a portion of the natural tissues. The child was therefore put under the influence of chloroform, and a small black button (such as is used on boots) was extracted. The ozæna and discharge from the nose soon disappeared.

II. FLEXION OF THE FOREARM IN WOUND OF THE PALMAR ARCH: SUCCESSFUL.

Under the care of H. C. JOHNSON, Esq.

[From Notes by F. VENNING, Esq., House-Surgeon.]

Henry Luthull, aged 6 years, a schoolboy, was admitted on August 23rd, and the following history was given by a surgeon who had previously seen him. A week before his admission, he was cutting something with a sharp knife, which slipped and cut him just over the region of the superficial palmar arch of the right hand. There was a great amount of hæmorrhage at the time, but the mother did not take him to a surgeon until

six days had elapsed after the injury. During this time there was every night severe hæmorrhage, and the mother at length becoming frightened by the bleeding, took him to a surgeon, who, finding bleeding was then going on, exposed the wound, and turned out a considerable amount of clot which was in the wound, and tried to take up the ends of the bleeding vessel, but failing to do so (on account of the parts being in a sloughy state), he brought him up to the hospital.

On admission, the wound having been bandaged up by the surgeon before his admission, and there being no bleeding, Mr. Venning merely sent him to bed, determining to watch for bleeding. On going round the wards on the night of the 23rd, about 10 p.m., Mr. Venning found hæmorrhage had again commenced; he, therefore, flexed the forearm upon the arm, and bandaged it up in that position, and the hæmorrhage immediately ceased. He did not take off the compress which had been applied before admission.

Aug. 24th. Early this morning, the night nurse had allowed the little patient to sit up, and had taken off the bandages, so that, about 12 o'clock, p.m., hæmorrhage had again commenced, and, therefore, soon afterwards, the forearm was again flexed upon the arm, and strapped up in this position; strict orders being given that the strapping was not to be removed, and that the patient was to be kept still in bed.

Aug. 25th. No more hæmorrhage had occurred; the arm was still kept flexed.

Aug. 27th. The compress was removed to-day, as there had been no more bleeding. The wound was healing nicely. Water dressing was applied, and flexion was still kept up.

Aug. 29th. There had been no return of the hæmorrhage.

Sept. 1st. The compress was removed to-day; the wound was quite healed; the strapping was removed, and the patient was allowed to get up and use the arm.

Sept. 2nd. The wound was entirely healed; the patient was much stronger, and was discharged next day.

The reader will find another successful case of the same treatment, also from St. George's, at p. 665; and another case has since occurred, and is still under treatment, where profuse bleeding from the radial artery in a case of amputation of the thumb has been hitherto controlled by somewhat similar means. All this seems to show the value of this simple and easy method of treatment.

Original Communications.

DIPHThERIA.

By WILLIAM NEWMAN, M.B.Lond., Fulbeck.

SINCE March 1st, 1858, I have met with nineteen cases of diphtheria—limiting the term strictly to those instances in which there has been present the fibrinous exudation on the surface of the tonsils, palate, or wall of the pharynx, with the coincident constitutional symptoms; not in any way referring to a much larger class of cases exhibiting sometimes specks of exudation matter, but running the course of ordinary sore-throat, or merging ultimately into well marked tonsillitis.

I will classify my remarks according to the *resumé* given in the JOURNAL of June 25th.

1. *Sex and Age.* Of the 19 cases, 11 were females, and 8 males, so that *sex* would seem to have but little, if any, influence. More than half the cases (ten out of nineteen) were under ten years of age; thus—

Under 1 year	- - - 0	Between 15 and 20	- - - 2
Between 1 and 5	- - - 4	Between 20 and 30	- - - 2
Between 5 and 10	- - - 6	Between 30 and 40	- - - 1
Between 10 and 15	- - - 2	Between 40 and 45	- - - 2

2. *Occupation* appeared to have little influence; in-door and out-door were alike attacked. Thus, in adults—one was the wife of a farmer; two were domestic servants in good houses; three were farm labourers; and one was a labourer's wife.

Residences were fairly good in certainly the majority. One death, of a female, aged 10, occurred in a house with ill placed and ill drained privy.

Food or clothing were fairly good in all; excellent in several instances.

Air. The purity or impurity appeared to exercise very little influence. The disease showed itself alike in low and damp situations, and in those high and bleak.

3. Every variety of weather contributed its share of cases; but no exact meteorological register was kept.

4. *Contagion.* Some facts would seem to favour this view; *e.g.*, a boy, aged 19, brought the disease from a village eight miles away, where many cases had occurred, to his home. There had been no prior disease in the house or in the village. A sister and brother had the disease not very severely. Two female children, aged 12 and 10, residing next door in the same block of houses, exhibited the disease severely; the youngest of the two died in two days. All these cases occurred within ten days.—A maid servant, aged 30, had the disease. Her mistress, aged 33, and two children (a boy, aged 6, and a girl of 5), showed the disease within ten days. All were recovering, when another maid servant, aged 25, also took the disease, went home on its appearance, and died within a week.

5. *Scarlatina.* Nothing occurred to prove any connection between the two diseases. Several of the children and nearly all the adults had had scarlatina prior to the occurrence of diphtheria. In one family (not quoted in the list, but personally known to me), three children had diphtheria in turn, and all died from the disease—all of them having two years previously been the subjects of undoubted scarlatina.

6. The *symptoms* attendant on the first appearance of the disease have almost uniformly been those of extreme depression, with consequently more or less of febrile reaction. External swelling of the neck has not usually been observed as a very early symptom; nor is dysphagia early in its appearance or actually necessary in the course of the disease. In one well marked case of a child, aged 4 (dying rapidly with extension of disease to the air-passages), there was never any dysphagia whatever. I am almost disposed to give as the very first local symptom (and the statement is borne out by at least two of the cases in the table) the existence of marked oedema both of the uvula and tonsils—the mucous membrane appearing to be infiltrated with serous exudation, and then studded by the characteristic lymph deposit.

7. *Exudation.* In by far the majority of cases, the exudation was present when the patient was first seen. In the few instances where the progress of the disease could be fairly watched, it came on from the first to the third day after the first complaint of illness.

8. *Microscopic examination* of the effused material in three cases gave no evidence of fungoid growth.

9. *Albumen in the urine* existed in one instance, but was not found in the other instances. Various accidental complications were noted in some of the cases.

10. *Treatment.* I have never resorted to salines, diaphoretics, or, in short, to lowering treatment of any kind; but, on the other hand, have freely given tonics, *e.g.*, tincture of sesquichloride of iron, in doses of ten to twenty minims every four hours; or bark with ammonia; stimulants, as sherry in full quantity, either with or without water, or good beer; nutriment in the shape of beef-tea, essence of meat, arrowroot.

Local Applications. I believe the best to be the strong hydrochloric acid, applied freely, in the first instance, to the parts affected, and repeated if necessary; and followed by the use of gargle of two drachms of hydrochloric acid and an ounce of glycerine with seven ounces of water.

11. *Tracheotomy.* I have no evidence on this point. One child, a girl, aged 4, died from disease invading the air-tubes; but tracheotomy was not permitted.

12. The *mortality* was 2 out of 19, or 10·52 per cent.

13. No *post mortem* examinations were made.

The disease in question is not only worthy of note in its ordinary course, for its danger to life from the exhaustion attendant on it; but also for the varied sequelæ which may show themselves, either as the disease subsides, or even after an apparent though partial recovery.

Sloughing of the cheek (interior), unconnected with the throat-affection by any direct continuity of surface, was observed in a girl aged 12; and subacute inflammation of the mucous membrane of the nose and mouth, lasting for some weeks, in the same child.

Affections in which the nerves principally suffered were very common. Aponia, lasting some weeks, occurred in a boy aged 7. Paralysis of the portio dura, coming on directly after the subsidence of the disease, on the side most affected, was observed in a female, aged 45. Coincident with this, there were paralysis of the laryngeal and pharyngeal branches of the vagus, and numbness of the hands and feet. Loss of power over the muscles of the palate occurred in a boy aged 7.

Excessive and long continued general debility was seen in

several of the cases; coexistent with it was very complete anæmia in two of the cases—a girl aged 7, and a boy aged 5.

Tubercular disease of the lungs was called into activity in one of the adult cases.

Of skin-diseases, eczema, impetigo, etc., were observed, with formation of numerous and even large abscesses on almost every part of the body.

The whole of these results may be cited as proofs of the deficient power in the system, with, in some of the cases, local manifestations of the constitutional disorder. Whether the affections of the nerves point to any special element in the disease itself, would seem as yet a matter of uncertainty.

C A T A R R H.

By C. HANDFIELD JONES, M.B., F.R.S., Physician to St. Mary's Hospital.

DR. COPLAND says of catarrh, "Although the most common of all diseases, there are few which are less understood, or have called forth a greater diversity of opinion." He defines it as a specific irritation of the nasal and adjacent mucous surface, which may run on to catarrhal bronchitis. In another place, he writes, "It is not pure inflammation, but an irritation of a specific or peculiar kind, attended by slightly increased vascularity, afflux of the circulating fluids, and augmented secretion." He does not sanction the application of the term catarrh to all affections of mucous surfaces attended with sero-mucous discharge. Neither does he assent to the opinion that the "defluxion" is a consequence of the suppression of the cutaneous perspiration arising out of the irritation which the secretion retained in the circulation produces upon the cephalic and pulmonary mucous surfaces." He infers that the causes of catarrh affect primarily the organic nerves supplying the surface principally disordered, and through them, the system generally.

Dr. Watson simply speaks of catarrh as a "moderate degree of inflammation."

Trousseau says, "The catarrhal state and the inflammatory are different, though in some measure allied (*voisins*). The first is distinguished from the second by the circumstance that its accompanying irritations do not terminate by suppuration. Their character is an extreme superficiality, a singular mobility, and diffusion." Again, he speaks of the inflammations even in the severest forms of catarrhal fever as "quasi-inflammatory, false, or bastard." Grave pneumonias, attending such states, are rather an acute stuffing of the lungs (like that of the nasal cavities) than a real pneumonia, which can fairly be attacked by venesection. He recognises very plainly the existence of a nervous element in the disorder which crops out (so to speak) very notably in influenzal catarrh.

Neither Rokitansky nor Virchow seem to recognise catarrhal inflammation as specially different from the same process in other parts.

Catarrhal disorder, for the most part, consists of a general affection, and a local. The latter shows itself in a hyperæmic state, more or less verging into downright inflammation of one or more mucous membranes, attended sooner or later with mucous, muco-purulent, or muco-watery exudation. The former consists of various febrile phenomena; it may be of brief duration, or but slightly marked. More particularly we may describe the general disorder as varying from slight chilliness, and subsequent febricitation, with some depression, to a well marked pyrexial condition, with initiatory rigors, followed by rapid, excited pulse, hot skin, thirst, and oppression; or, as is often the case in the present day, the patient chiefly complains of depression and pains in the head and limbs; feels as if beat and bruised, or sore all over, with a cool skin and languid pulse.

The picture of the local affection will of course vary according to its site. Its general features are the above mentioned phlegmasial action, together with certain morbid sensations, and induced movements, often of reflex character. We will describe these more particularly:—As to the *inflammation*. The vessels of the inflamed part are more or less filled and distended with blood, so that the surface presents, in extreme cases, a full red colour. Microscopically examined, it is seen that the congestion most affects the superficial vessels, the terminal capillary loops, which alone may appear filled with blood. Sometimes small extravasations take place, mostly on the surface; but also interstitially. Judging from what we can observe in the conjunctiva, we may believe that the blood in the vessels is not in a state of actual stasis; at least, not during the earlier period of the inflammation. A full venesection will for the time