from the ventricular wall, and, almost invariably following the course of the blood-currents, pass into the great arterial trunks leading from the ventricle. The concretion thus running along the main trunk is either tubular in character, or is entirely surrounded by blood, or grooved spirally.

There is yet another condition, in which the deposition, being more slowly carried on and more extended in its character, occurs in the form of a layer, having no one point, but a wide spread surface. In this form of deposition are produced the hollow concretions—those which line a cavity or a vessel, and enclose in their centres red blood.

Lastly, there is that concretion which undergoes *organic* development. In all instances of this nature, there is, in my opinion, a preceding change in the endocardial lining. There are injection of the subserous layer, œdema, exudation of plastic matter, with rupture of surface, and production of a point on which the fibrine of the free blood finds not only a basis, but a basis directly connected with the endocardial membrane. In process of time, vessels from the broken membrane are thrown out, and, extending through the fibrinous matter deposited by the blood, furnish it, if the patient live long enough, with the materials for a new and special organic growth, possessing a permanent and independent existence.

[To be continued.]

Original Communications.

PALLIATIVE TREATMENT OF CANCER.

By THOMAS HUNT, Esq., F.R.C.S.

My remarks apply only to those cases of true scirrhus in the breast in which there is a hard and moveable tumour, not yet advanced to the stage of ulceration.

One indication in this condition of things is as far as possible to *prevent* the occurrence of ulceration. I regard this process as chiefly the result of the *pressure* sustained by the skin and cellular membrane, from *one* hard substance within (the stony tumour), and *another* hard substance without (the patient's corset). This pressure is generally sufficiently severe not only to give rise to pain, but to effect, first, a congestion of the healthy vessels, and secondly, an absorption of the healthy structure, without being sufficient to absorb (as has been proposed by higher pressure) the cancerous tumour itself. A second indication is, to support the tumour, without undue pressure; and a third, to relieve pain.

All these purposes are easily accomplished, in most cases, by the following expedients:—The whole breast should be allowed to rest on a thick broad compress of cotton wadding. When the breast is very pendulous, an old silk handkerchief may be passed, as a sling, between the mamma and the wadding, and tied over the opposite shoulder, one tail of the handkerchief being passed over the clavicle, the other tail over the scapula and spine; and all should be included in a capacious corset, which will then become a comfortable support, instead of acting as a tormenting *vice*. Where the pain has been very severe, I have applied to the skin a belladonna plaster, spread thinly on soft thin leather. This, by encompassing not only the whole breast, but an inch or two of skin beyond it in all directions, will materially assist in giving comfortable support, and also in allaying the irritability of the nerves. By this *local* treatment I have often succeeded in relieving the patient of all pain for months together, and also in preserving the isolation of the tumour, and in many cases diminishing its size.

Together with this local treatment, great attention should also be paid to the bodily health and mental tranquillity of the patient. She should be encouraged to hope that her pains may be much relieved, if not entirely removed; that the disease may probably be checked, or even so far subdued as to become for a very considerable period comparatively harmless and benign. The health should be sustained by a generous but regulated diet; by engaging the patient as much as possible in cheerful society; and last, not least, by moderate but frequent exercise in the open air, in a locality where the atmosphere is pure and mild. This will be far, very far, better than confinement in the house, and infinitely better than confinement in the wards of a hospital containing patients with open wounds. If the health is feeble, much good will be derived from the following formula:—

R Tinct. ferri sesquichloridi 3iij; liquor arsenici chloridi 3v; aquæ destillatæ 3viij. M. Capiat minima xl ter in die ex aqua post cibum.

The dose of both the iron and the arsenic must be regulated by their respective effects. The iron is *generally* useful and necessary, the arsenic *always*; for, although I have rarely found a cancerous tumour entirely dissipated by arsenic, I have as rarely known the mineral fail to check its onward course. It most assuredly exerts, when discreetly administered, a certain amount of specific influence over the disease. I wish I could say more than this in favour of arsenical treatment; but truth requires that I should stop here. That there is no medicine which is so uniform and potent in its controlling power over this disease, is, however, most evident; and in this opinion I am supported by many high authorities in surgery.

As illustrative of the good effects of the proposed treatment, I have selected, out of many, the two following cases.

CASE I. A female servant, aged 40, single, consulted me in the autumn of 1858, on account of a tumour in the left mamma, which had been observed to exist upwards of a year, and which was becoming painful, and was beginning seriously to affect her health.

Sept. 16th, 1858. The tumour was about the size of a nutmeg, irregular, moveable, and of petrous hardness, having little sensibility, and little apparent connexion with the surrounding parts. It occupied a space a little below the nipple, which was not contracted, nor was the skin puckered or discoloured. The absorbent glands were unaffected; but the complexion was sallow and dusky; the appetite nearly gone; the nights were disturbed; and she complained of feeling "very weak and sinking". The alvine and uterine secretions were normal. A soft cushion of cotton wadding was placed under the whole breast, resting on a corset of ample size. She was directed to live well; to take every opportunity of exercise out of doors; and the chloride of arsenic and iron was exhibited thrice a day, in doses of twelve minims and a half of the solution of chloride of arsenic, and seven minims and a half of the tincture of sesquichloride of iron.

Nov. 1st. The pain was much relieved, and she had better nights; the appetite was improving, and the patient felt and looked better. The tumour was diminished in size. The treatment was continued.

Jan. 25th, 1859. She had no pain whatever. The tumour was still wasting. The health was improving. The bowels being inactive, a compound rhubarb pill was directed to be taken every night, and the arsenic and iron persisted in.

July 22nd. She had persevered most regularly in the treatment up to this time, and had had no pain whatever for several months. She looked well and plump, and had a good appetite. She considered herself quite well; but I advised her to persevere for at least another month, and to take especial care to use the cotton wadding as before.

This patient had been advised to have the tumour removed; but she strongly objected to the operation, and had resigned herself to despair. When I last saw her, her spirits were so good that I suppose no one could have convinced her that the tumour was malignant.

CASE II. Miss —, aged 24, residing on a healthy eminence in the country, had observed for several months a tumour on the right mamma, below and to the right of the nipple. She was a fine handsome girl; and nothing but the extreme hardness of the tumour, coupled with failing health, could have convinced me that she was the subject of cancer.

The tumour was of the size of a bantam's egg, of oval figure, but irregular surface, very hard. The mamma was swollen and tender; the axillary glands were not sensibly enlarged, but very tender, and somewhat painful. The pain in the tumour was often intense and lancing. The nipple was normal; the skin was slightly puckered over the tumour, which, although buried in a highly developed mamma, was in some degree adherent to the integuments. The health was considerably impaired. The bowels were very much constipated, and the catamenia very irregular, sometimes profuse, and sometimes scanty. There was also severe leucorrhœa, with pelvic pains and general uterine distress; as well as a failing appetite and a coated tongue. The pulse was rapid and feeble. The patient had wasted considerably during the last three months, and her spirits were "wretched".

March 12th, 1858. The breast was enveloped in a belladonna plaster, supported by cotton wadding and a sling. She was directed to regulate the bowels by pills of colocynth, aloes, and rhubarb; and to take the chlorides of iron and arsenic, as

in the former case. She was ordered a full diet, with stout or porter; directed to take exercise in the open air; and encouraged to hope for a speedy amendment. In this she was not disappointed, for in less than a fortnight her health had considerably improved.

May 5th. She was much more free from pain, and better every way in health and spirits. The leucorrhœal discharge was much reduced; the catamenia had become regular and normal. The tumour felt more loose and moveable, and less irregular on its surface; and the puckering of the skin was less obvious. The pain and tenderness in the axilla were quite gone. She complained, however, of headache, and thought the iron did not quite agree with her. The arsenic and aperient pills were continued, without the iron; and the local treatment as before. Fowler's solution, in doses of four minims, was afterwards substituted for the chloride, which began to nauseate a little.

July 19th. She was better every way; appetite good; was gaining flesh, and was in high spirits. The tumour was decidedly reduced in size, and almost free from pain. The treatment was continued.

From this period I saw nothing of the patient for five months; and, as I had never revealed to her the malignant character of the tumour, she expected she was recovering altogether. Accordingly, she visited some friends at a distance, neglected her medicine, and took no heed to the support of the breast.

Dec. 12th. The tumour was larger and more painful; the catamenia were too frequent; leucorrhœa was constant; and there had been a discharge of blood from the bowels, which were constipated. The appetite was fickle and capricious, and the patient looked haggard and dejected. The arsenic, iron, and purgatives, were ordered to be resumed.

I have not seen the patient since the last date; but the relapse of all the bad symptoms, on her neglecting the treatment, is quite as instructive as the benefit previously derived from it.

I am quite aware that it has been suggested by a microscopic pathologist of repute, whose name I forget, that inasmuch as a cancer is a parasitic growth, the application of warm coverings is objectionable, as tending to the more rapid development and growth of the parasite; but a few such facts as the above surely afford a sufficient refutation of this theory. Variations of temperature are great hindrances to healthy action in local disease of every kind; and besides that the cotton wadding serves as a soft cushion, it probably exercises a salutary influence by regulating the temperature of the parts.

Alfred Place, Bedford Square, January 1860.

CASE OF TRAUMATIC EMPYEMA OF SIXTEEN MONTHS STANDING, WITH FISTULÆ: TREATED SUCCESSFULLY.

By ALBERT G. WALTER, Surgeon, Pittsburgh, Pennsylvania, United States.

JOHN O'NEAL, aged 32 years, of decidedly strumous diathesis, with fair skin, red hair, by occupation a river-man, had enjoyed excellent health up to the period of his receipt of the following injury. Whilst in discharge of his duties as mate on a Mississippi river steamer, in a dispute with one of the crew, he was stabbed with a broad bladed knife under the left arm, the knife entering the chest between the eighth and ninth ribs, an inch posterior to the median line. Profuse external bleeding from the wounded intercostal artery followed, which was arrested by closing the wound with stitches. Dyspnœa, cough, with bloody sputa and high inflammatory fever, ensued, terminating in empyema. On the ninth day after the injury, an abscess pointed below and anterior to the original wound (which then was closed). This was opened by incision, giving vent to a large quantity of pus. A month later, the wound itself opened spontaneously; from both openings, purulent matter was freely discharged. After six weeks, he had so improved as to be able to go about; but the dyspnœa, chills, and night-sweats, with loss of appetite and strength, too plainly indicated the severe irritation under which his system was suffering. Some months later, considerable improvement had appeared, as the openings, ceasing to discharge, became closed. This state was, however, of short duration. With the recurrence of pleuritic pains, dyspnœa, and hectic, the original wound again opened, giving exit to a still greater quantity of highly offensive greenish

matter. Discouraged by the fruitless efforts of his medical adviser, his means exhausted in the vain endeavour to find relief, he returned to his home near this city, and applied for admission into my hospital.

His appearance was that of a person in confirmed phthisis; the shoulders stooped; respiration was short and hurried; the pulse was feeble and rapid; he had frequent chills, during which the countenance assumed a purple hue; and there were irregular paroxysms of fever, with night sweats. The hands and feet were moist or clammy; emaciation was extreme; he had a short teasing cough, with purulent sputa, loss of appetite, disturbed sleep, occasional diarrhœa, with œdema of the ankles. The chest was much sunken in the infraclavicular region on the left side. The opening, surrounded by a papilla of fungous granulation, gave exit to a stream of pus on forced and prolonged inspiration, and admitted a probe downwards and forwards to the extent of four inches and a half, and posteriorly to over six inches. Percussion gave a dull sound over the entire left lung. Auscultation indicated a total absence of respiratory murmur, with the exception of a small space anteriorly, where it could be feebly perceived. In addition to the chest affection, there existed on the left side of the anus, an inch from its orifice, a fistula communicating with an internal opening above the sphincter, which had appeared some months ago.

Under all the circumstances, especially the condition of the pleural sac, whence issued daily fully two pints of offensive purulent discharge, local treatment directed to the obliteration of the cavity, aided by generous aliment, alone promised to afford relief. With this intention, the cavity was daily injected with a weak solution of tincture of iodine in water during some weeks, with but slight and temporary improvement. Concluding that retention of coagulated blood from the wounded artery, or a carious state of the ribs, was the cause of such prolonged and profuse suppuration, I determined on enlarging the opening, and clearing the cavity of the offending *fons mali*. With the patient's consent, on December 8th, 1857, the fistula was enlarged, anæsthesia having been procured by chloroform; but the ribs were found so closely approximated that a free discharge could not be effected. A free crucial incision was made through the skin, fascia, and muscles; the flaps were separated, and an inch of the eighth rib was resected by pliers. This was hardly accomplished before a quantity of very offensive matter escaped, followed by masses much resembling putrefied lung, which were removed by the forceps, but which proved to be the fibrine of coagulated blood, which, acting as a foreign body, had kept up the suppuration. The pleural sac was washed out with water, and carefully explored with the finger, which, however, could not reach the pleura pulmonalis, as the lung was forced upwards and backwards by the long continued compressing force of effused blood and pus. The portion of rib removed exhibited increased ossification, with considerable hypertrophy, as the result of the prolonged inflammatory action in the pleura. But little blood was lost during the operation; the exhaustion was, however, very great—the after effect of chloroform requiring a free use of stimulants. Reaction was slight. The relief to the patient was well marked; his breathing became more free and less frequent. The wound was kept open by pledgets of lint; anodynes at night, with a supporting diet, were administered; and he was directed to make deep and long inspirations frequently, for the purpose of inflating the compressed air-cells, hoping thus to expand the lung, and eventually bring the surfaces of the pleura into contact, and obliterate the sac by adhesion of the costal and pulmonary pleura.

For a time the improvement was well marked. Hectic disappeared; the appetite and strength returned; respiration had become deeper and more free from oppression; the left side of the thorax had sunk in front and laterally; the discharge had gradually diminished in quantity, and improved in quality. Still the probe entered the cavity freely. Tincture of iodine injections were again resumed, after previously washing out the sac; but these had to be discontinued, as they caused chills and irritative fever.

The condition of the patient, having remained stationary for some time, suddenly and without any evident exterior cause assumed an unfavourable aspect. An increased discharge of a more offensive matter appeared, with fever, night-sweats, and diarrhœa. Though the empyematous cavity had evidently become much diminished in size, yet, the original cause appearing to continue, on February 11th, 1858, a freer opening in the chest was made by resecting two inches of the eighth and ninth ribs. The interior of the abscess being thus