The question may therefore fairly be asked, whether the loudness of the second sound does stand in a direct causal relation to the strength of the ventricular systole and the reaction of the arterial walls?

I have observed in typhus fever, fatty heart, spanæmic women, and cases of general debility, that when the first sound was decidedly weak, the second was still loud and clear. In the so called accentuation of the second sound of the pulmonary artery where it would be said to exist, it has appeared to me usually rather a forcible diastolic recoil of the artery than an accent properly so called.

Taking the second aortic sound, or its replacement by murmur, as a basis for differential diagnosis, the following points have presented themselves to my notice.

A preternatural accent of the second sound in the direct course of the aorta, to the right of the sternum, has been observed and specially indicated in some cases of aortic aneurism. I have recently had an opportunity of watching throughout a case of aneurism of the ascending aorta. The earliest symptoms were those which apparently indicated an inflammatory condition of the aorta: at this period, the only persistent physical sign appreciable was marked intensification of the second sound over the aorta. Some months later, pulsating tumour was seen and felt about the second and third right costal cartilages; the second sound still being preternaturally loud. In the early stage, it occurred to me that the loud second sound might be explained on the hypothesis that the inflammation of the aorta had induced unusual contractility of this vessel, and thus, by rendering it more rigid, making it a better conductor of sound. At a later period, the extensive thickening of the coats, by excessive deposit, might have rendered the tube a better conductor; at all events, the intensified sound could not be explained upon the hypothesis of increased recoil of the vessel, since here the elastic reaction of the vessel was impaired or even destroyed by extreme degeneration. A further argument against the loudness of the second sound being supposed to result from the injecting force of the heart, may be borrowed from Dr. Stokes. When treating of softening of the heart in typhus fever, he states, that in his 41st and 56th cases, the second sound was preternaturally loud, and in the latter case attended with impulse, while the first sound was extinct, and the impulse not appreciable by palpation. I again submit, therefore, that the hypothesis of the causal relation between the loudness of the second sound and the propulsive force of the ventricle, requires reconsideration.

It is hardly necessary to mention induration of the pulmonary tissue, excavations therein, contraction of the lung from phthisis, permitting the pulmonary artery to come in immediate proximity with the chest wall—a pathologic condition first indicated, I believe, by Dr. Sibson—enlargement of the liver, etc., as sources of intensification of the sounds. I have, however, some reason to believe that a marked intensification of the sound in the post clavicular space may, now and then, assist in the diagnosis of deposit at the apices of the lungs.

In insufficiency of the aortic orifice, the second sound is more or less perfectly replaced by murmur, soft, harsh, or musical. Such murmur, however, does not nor cannot measure the amount of patency. I have particularly observed that in some cases, in which, judging from the extraordinary collapse of the arteries, the patency was extreme, that the murmur in its physical characters differed in no respect from other less advanced cases in which no collapse was observed.

I believe enlarged experience scarcely supports the notion of Dr. Hope, that aortic may be distinguished from other murmurs by the pitch. The pitch of a murmur at any orifice infinitely varies with the physical condition engendering it. An aortic regurgitant murmur may be of so high a pitch as to approach sibilation in point of intensity.

I may add, that this distinguished observer was also decidedly in error when he stated that a mitral murmur was never audible above the level of the aortic valves. Murmurs at the orifices are not so restricted as many suppose.

There are one or two points, however, with regard to aortic diastolic murmur, which I will take leave to mention. The first is that, contrary to the opinion of some observers, I have not unfrequently found it attended with well marked diastolic fremitus even at the apex, and this in cases unattended with any excessive degree of spanæmia; and, indeed, there appears no reason why a partially rigid valve should not be susceptible of being thrown into tactile as well as sonorous vibration by the reflux current.

It is usually taught that the differential diagnosis of aortic regurgitant murmur may be, at least, in part established by

the clear click of the valves of the pulmonary artery auscultated to the left. I am far from asserting this is not occasionally correct; but so far as my opportunities of observation extend, I should, in the majority of cases, deny the validity of this proposition.

When the diastolic murmur is at all intense, I believe the second sound of the pulmonary artery is always more or less obscured, and in musical murmur almost invariably so. Indeed, if we bear in mind the anatomical condition existent, and consider how readily vibrations are excited in contiguous and homogeneous structures, the difficulty of conceiving the reverse is obvious.

The limits of these murmurs again, and particularly the musical, have been too much restricted; they extend over a very wide area; at times, all over the anterior and posterior surfaces of the chest, from the epigastrium to the pubis; and posteriorly, from the vertex nearly down to the lower end of the sacrum. From their site of origin, they may, I am quite sure, from comparison of cases examined after death, extend across the apex and round the axilla to the posterior scapular region; and this without coexistent disease of the mitral valve.

Excluding from consideration some rare cases of aortic aneurism, described by Gendrin as being liable to be confounded with aortic patency, is there any other condition which gives rise to similar physical phenomena? I believe there is.

A few months ago, I exhibited, at the Pathological Society, a specimen of diseased suprarenal capsules; they were removed from a female advanced in life, who had exhibited well marked symptoms of cardiac disease: a harsh double murmur was heard, with its maximum near the base, and extending with great intensity up the aortic, and downwards to the apex, axilla, and posterior scapular region. The heart's action was somewhat heaving, and its rhythm irregular. These phenomena were persistent, and scarcely underwent any change when examined on several occasions by Mr. Robinson (whose patient she was) and myself. We considered there might possibly exist both aortic and mitral disease. After death, hypertrophy of the heart was found. The aortic and mitral valves were apparently intact; but the aorta at its arch was nearly converted into a long tube; and in its ascending portion, rough, calcified portions projected, the lining membrane being entirely destroyed.

In another case, of Dr. Hawksley's, similar physical phenomena were constantly observed, and the same irregular rhythm. After death, enlargement of the heart was found; rough, calcified projections studded the ascending aorta and its arch; a slight calcification was certainly found at the base of one of the aortic valves, but not preventing tolerably fair closure; and, to the best of my judgment, quite inadequate to the production of the intense double murmur audible over base, apex, and axilla.

[To be continued.]

ON THE OCCURRENCE OF AMAUROSIS IN PATIENTS SUFFERING FROM BRIGHT'S DISEASE OF THE KIDNEY.

By HOLMES COOTE, F.R.C.S., Assistant-Surgeon to St. Bartholomew's Hospital, etc.

THE connexion between amblyopia, or amaurosis, and Bright's disease of the kidney, was, I believe, first pointed out in 1849 by Landouzy of Rheims, who published, in *L'Union Médicale*, a paper containing several cases in illustration of the fact, occurring in patients in whom nephritis had been produced by the use of alcoholic drinks, by exposure to damp, or by want of the common necessities of life. It is a matter of surprise that, with the knowledge of the frequency with which extravasations of blood and other accidents occur within the cranium in persons the subject of albuminuria, the idea should not have more readily suggested itself of examining in such cases the condition of the nervous contents of the orbit, constituting, as they do, an integral part of the cerebro-spinal centre. We might in many instances reasonably expect to find there morbid changes of a similar character to those noticed on the convolutions, or in the substance of the brain, provided there were any means of illuminating the retina and the choroid behind the transparent media of the eye. I need scarcely remark, that this end is readily attained by the use of the ophthalmoscope.

When the kidneys are so diseased that the proper urinary

excretion is impaired, and the effect is felt in the general constitution of the blood, the walls, not only of the larger arteries, but likewise of the capillaries, undergo a molecular or granular degeneration, by which they lose their elasticity and strength, and are very liable to give way. In the examination of the body of persons who have died under these circumstances, I have found in the brain four or even more small extravasations of blood in the various stages, between recent effusion and complete repair. The same morbid changes occur, I believe, in the arteries of the choroid, retina, and vitreous humour, where they give rise to changes which impair or destroy vision, and may be recognised during life.

I had occasion to make an examination of the eye, a few days ago, in the following case.

Anne P., aged 50, an unhealthy looking woman, with puffy countenance and sallow complexion, was under the care of an eminent physician, a friend of my own, for a complication of diseases, any one of which was sufficient to shorten life. She had had chronic bronchitis for many years; the right cavities of the heart were dilated, and there was general anasarca; she suffered from pain in the loins; the urinary secretion was diminished in quantity, and albuminous; there was hard cancer of the right mammary gland, with similar disease of the absorbent glands extending into the neck. Three weeks ago, while under treatment, she became suddenly blind of the right eye, and continues so up to the present time. Upon examining the affected organ (April 10th) with the ophthalmoscope, I found the transparent media of normal consistence, but the posterior surface of the eye wanted the usual brilliant hue; the vessels of the retina appeared to me smaller than natural, and to cease more abruptly; the vessels of the choroid membrane were not made out with common ease. By throwing the light to the right of the entrance of the optic nerve, I brought into view three smaller and one larger extravasations of blood, the former towards the lower part of the globe, the latter directly in the axis of the pupil: they seemed to be either in the structure of the retina, or between that membrane and the vitreous humour; the usual colour of blood was apparent. In another part of the retina, there was a yet darker spot, apparently of pigment, and of uncertain character.

In such a case, it is obviously impossible to direct, for the treatment of the diseased eye, the administration of any remedies which might interfere with the management of the more serious affections of the kidneys or thoracic viscera. In all probability, this patient will have ceased to live before the completion of any process of repair in the eye; and we know by experience that absorption of blood or other fluids takes place very slowly in a diseased organ, however rapid may be the same process among tissues perfectly healthy. In such a case, the administration of mercury would be inadmissible, on account of the general condition of the patient's health; and, even if it were otherwise, I am yet to learn that mercury "promotes absorption" otherwise than indirectly; namely, by controlling the action of the smaller vessels in an inflamed part, and preventing further effusion. Such circumstances do not exist in the case now under consideration.

Although Blödig found only three cases of this combination of amaurosis and Bright's disease of the kidney during the five years he studied in the Ophthalmic Clinic of Vienna, I believe it will be found of more general occurrence—not perhaps always manifested by sudden loss of vision from the rupture of diseased capillaries and the effusion of blood, but by changes dependent on the disturbed circulation through vessels which have lost their elasticity and power of regulating the flow of blood through tissues perhaps the most delicate in the human body—changes which, though slow, are not the less sure to terminate in the disorganisation of those parts upon which depends the susceptibility to the impressions of light. Thus Dr. Heymann of Dresden found, in one case of similar nature, turbidity of the vitreous humour; in another, a sort of serous swelling of the optic nerve papilla; in another case, the retina was covered "as if by a cicatrix". I found, in a case recently examined, loss of brilliancy in the choroid—absence of defined arrangement of the arterial capillaries. The vessels brought into view appeared to be venous, and were tortuous and varicose.

The case here related illustrates the importance of studying diseases of the eye in conjunction with other branches of medicine; for, by attending to the condition of organs which may be remote, and not long ago would have been considered wholly unconnected with the ophthalmic affection, can we alone hope to be of service to our patient.

THE ANATOMY, PHYSIOLOGY, AND DISEASES OF THE MEMBRANA PUPILLARIS.

By AUGUSTIN PRICHARD, Esq., Surgeon to the Bristol Royal Infirmary.

[Read before the Bath and Bristol Branch, February 19th, 1857.]

I HAVE ventured to introduce this subject to the notice of the members of the Society, for the purpose of pointing out one or two anatomical errors which, in my opinion, have hitherto existed in the descriptions of the membrana pupillaris and its vessels, and for the sake of bringing forward an opinion which I hold as to the office it serves in foetal life, and also because I wish to narrate a few cases of disease of the eye, which I refer to the continued existence of this organ.

The membrana pupillaris, a clear transparent membrane, closing the pupil during foetal life, was first discovered by Wachendorf, in 1738. Afterwards, Albinus, Haller, Zinn, Wrisberg, and Dr. William Hunter, described it about the same time, but no one of them with complete accuracy. It is a simple membrane, continuous with the anterior surface of the iris, quite transparent, and found at every period of foetal life. The centre of the posterior surface is in contact with the anterior capsule of the lens. Dr. Jacob was the first British author who pointed out that the membrane exists until the ninth month of foetal life; for it had been considered previously not to exist after the seventh—an error originating from the fact that the vessels cease to be permeable at that time; and consequently the membrane is almost invisible, and can only be demonstrated by great care. Tiedemann, Retzius, Arnold, and M. Portal, were also of the opinion that the membrane remained till birth.

I must call to your remembrance one or two facts respecting the iris, which it is necessary to keep in mind in order to understand the subject, and which, like other anatomical minutiae, are apt to slip away from the recollection, unless there is something to induce us to dwell upon them.

In the fully developed and healthy iris, the vessels are posterior, and are covered in behind by the pigment or uvea: the vessels are tortuous and anastomose tolerably freely, the general direction being towards the pupil, where there is a more free anastomosis, miscalled the "circulus arteriosus iridis minor". The human iris, freed from pigment, is almost transparent. In the very dark eye, particularly in those of the coloured tribes, there are pigment-cells scattered upon the anterior surface, in addition to the uvea behind; but, in the blue or light coloured iris, there is no pigment in front: the colour is due to the reflection of the light through the transparent iris from the uvea, the light being decomposed, and colour thus developed by the finely divided surface of the membrane; and when the pigment has been removed by washing, there is no appreciable difference in colour between an iris that was originally blue and one that was grey, or green, or light brown.

In the human foetus, there are vessels upon both surfaces of the iris. The posterior set passes as far as the pupillary margin, and not beyond; whilst the anterior run onwards in a straight course upon the membrana pupillaris, and, turning round at a little distance from the pupil, pass back to gain another part of the membrane, but still upon the anterior surface.

In an injected foetus of about six months, the membrana pupillaris is seen in perfection. The iris is narrow, and not fully developed; the pupil is large; and the membrane is thin, transparent, and covered with fine vessels, visible by the naked eye, particularly if it is stretched upon a white surface. (Fig. 1.) These vessels form a series of loops upon the membrane, and occasionally one traverses the centre. Cloquet, in his memoir on the subject, says that there are thirty or forty distributed in this way; and Henle says there are from ninety to a hundred. The number no doubt varies in different specimens, for I have counted upwards of one hundred and twenty upon an injected foetal iris, passing forwards to supply the membrane. I have also seen that a few vessels run from the posterior vessels of the iris, and inosculate with the arteries on the membrane. As the foetus advances towards the termination of intrauterine life, whilst the membrane is still entire, the vessels are gradually obliterated, like the vessels of the posterior capsule of the lens, which become incapable of carrying red blood, and gradually disappear, leaving the organ perfectly transparent. I have found the membrane entire and transparent in the eyes of a foetus of nine months; and I have once seen, in a specimen of