plied, it follows, as a matter of necessity, that when they are found wanting or deficient they ought to be supplied, if possible. It may happen, in some cases, that the very organisation itself is incapable of appropriating such pabulum after being furnished with it, but these are the exceptions, and not the rule, unless such organisation be worn out by protracted disease or old age. In the special matter of this albuminuria, as we see that in most cases the dreaded symptom seems to be preceded by hæmorrhage or other sources of drainage of the red particles, and is, when it takes place, itself a drainage of the albumen, so an extra supply of steel and albumen appears the readiest method of meeting these evils, and the following deductions may therefore be advanced with safety.

1. When the premonitory symptoms of debility and anæmia have not as yet produced actual renal disorganisation, the supplies of the deficient elements afford a fair hope of averting the threatened malady of albuminuria.

2. If the renal disorganisation be commencing, the improvement of the mass of the blood by such means may yet give the constitution the power of arresting the dire changes that may precede, accompany, or supervene upon the albuminuria.

3. Even if the disorganisation be considerably advanced, such supplies may postpone the fatal issue for an indefinite period.

The cases of absolute and complete disorganisation or disglanding are, of course, as already observed, beyond the scope of this or any other therapeutic paper, and belong exclusively to the province of morbid anatomy.

MINOR QUERIES IN MEDICAL SCIENCE.

By W. HINDS, M.D., Lecturer on Botanical Science at the Birmingham Midland Institute.

II.—CHOREA AND ACUTE PERICARDITIS: WHAT IS THE NATURE OF THEIR RELATION?

CASE III. A little girl, aged 10 or 11 years, was attacked with fever and pains in the joints, especially of the shoulders and knees, and had a most distressing pain referred to the region of the heart. The attack began by slight indisposition for about a week before I saw her, and there was also some gradually increasing cardiac pain. On April 22nd, a physical examination revealed a to-and-fro or friction-sound, with a kind of bellows-murmur apparently intermingled. There was some cough; no dulness, nor crepitation over the chest. By the 25th, the to-and-fro sound had become very distinct indeed and louder. The countenance was anxious, and expressed distress.

From the first, the case was treated by leeching over the region of the heart, and by calomel in repeated doses; also by a blister, which was dressed with very much diluted mercurial ointment.

April 28th. The friction-sound became less rough; the gums were tender from the mercury. The breathing was less short, and the pulse fuller. There was no delirium, but a little jactitation and restlessness.

May 1st. The tongue was found blistered on the right side; the friction-sound still continued; and chorea had completely supervened. The arms were constantly tossing about involuntarily. Half words were continually uttered. There was perfect consciousness, and the mind was apparently unobscured in the slightest degree. The jerking movements were so severe, that the patient was near falling out of bed sometimes. She had had no sleep for two nights; she answered questions in an abrupt and jerking manner. The skin was cool, and there was no fever. The patient sank down in the bed, and could not be kept with the head upon the pillow. She declared herself free from pain everywhere, except in the arms. The mucous membrane of the tongue was much ulcerated, probably by the mercury. She took half a grain of opium each of the last two nights without any effect. The calomel was ordered to be omitted.

R. Ætheris chlor. ʒiss; tincturæ opii ʒxl; aquæ ʒiv. M. Capiat cochleare medium secundâ quâque horâ. She was also ordered to take the carbonate of iron in six-grain doses thrice daily; and to have free nourishment.

May 4th. She was much better; had slept many times, an hour at once. The movements were much less severe. The skin was quite cool. She had more command of speech.

May 7th. No further improvement had taken place. There was insomnia, even though she took last night a grain of powdered opium. Her general aspect was about as before. She was ordered to have one-sixth of a grain of morphia and two grains of extract of hyoscyamus at bed-time, and to change the carbonate of iron for tincture of the sesquichloride. The friction-sounds were still evident.

May 14th. She had slept well the last night or two for the first time since the attack. The choreal movements were gone; there was not the least trace left. She was dressed, and able to sit up. There still remained the friction-sound, though not so distinct.

May 21st. The patient appeared well, except some general weakness. At the same time, the friction-sound, to some extent present at the last examination, had entirely ceased. There was a very soft bellows-murmur only with the first sound.

REMARKS. The last statement shows that active or calomel treatment omitted fourteen days before, must not necessarily be pursued because the to-and-fro sound exists—the treatment being regulated by the whole circumstances. It is evident that the inflammation of the pericardium from which the exudation of lymph proceeds, requires special treatment only so long as it lasts, and that the exuded lymph which makes the two surfaces rub together, so as to produce audible sounds, will or can only gradually disappear, and will do so with even a tonic treatment. I am aware that this contains an assumption at variance with the views of some high authorities in medicine.

The view here alluded to is, that the lymph on the surface of the heart and pericardium never does clear away; and that when the friction-sound ceases, it can only be because the two surfaces become glued together. I believe this view to be unsubstantiated; but reserve this special point for a future paper, when I shall give other cases in illustration. To discuss it now, would be wandering from my present object—which is to inquire into the relation which the disease under notice holds to chorea.

The connexion of chorea with acute rheumatism was probably first specially noticed and commented on by Dr. Bright, in the *Medico-Chirurgical Transactions*. It has been also mentioned by Dr. Copland; and Dr. Watson says:—"I certainly have seen jactitations like those of chorea in a few instances of rheumatic carditis." Dr. Watson, however, does "not perceive any obvious or direct connexion between the cardiac disorder and the nervous disorder"; but he conjectures some relation to the fibrous structures supposed to be implicated in both, or else some morbid condition of the membranes of the spinal canal.

It is scarcely, I think, desirable to adopt either of these views of that relation. The history of the case given above, shows the supervention of clonic spasm to have occurred exactly at that point at which the nervous system became exhausted, and its complete disappearance at the exact point when the exhaustion may be regarded as removed; and that there is manifestly a more direct or intimate connexion here than simply with the disease itself, for the cardiac affection not only began a considerable time before clonic spasms began, but must be regarded as not entirely removed even after their subsidence. Remotely, however, no doubt, there is ground for an indirect connexion. Nervous symptoms unquestionably do often occur in even chronic affections of the heart.

In carditis, the nervous symptoms are shown in the "care-worn" and in the "anxious expression" of countenance, and the "distress" so generally noted. Then circulatory disturbance tends also to insomnia; and when the heart is inflamed, and its movements obstructed, an impression will of necessity be stamped upon the nervous system and brain. It may be simple distress or delirium, or clonic spasm—the two former being the most common.

The state of exhaustion being induced, and the nervous system being also in this manner excited, and that in a delicate or nervous subject, will sufficiently, I believe, account for the choreal affection. The power of the brain and of the will are nearly prostrated. In the alimentary canal are present stimulating and vitiated secretions and other morbid matters, which often alone will give rise to spasm, as we see in cholera, in tetania, and many other cases.

The doctrine of spinal irritation in even idiopathic chorea is a very questionable one. If we look to the causes of chorea, enumerated by the best authorities as unquestionably ascertained, we shall find these to be of two classes; namely, *cerebral* or *centric*, and *peripheral*. Amongst the most prominent of these causes are sudden shock or fright, blows on the head

of the first class, and gastric irritation from irritating matters and worms, and painful menstruation on the other.

For spinal irritation, therefore, a term which should probably be expunged altogether from medical etiology, we may, I think, substitute *peripheral* irritation of one kind or other—a really intelligible and meaning term—in some if not many of the cases sought to be explained by the former.

Then we have *paralysis agitans*, in some persons with semi-worn-out cerebral power, evidently one phase of the choreal or clonic state of the muscles.

From a careful examination of this subject, I find it difficult to conclude that any more direct relation exists between chorea and carditis than the one now explained; and I do not think it at all necessary to this inquiry, even if it were possible to attempt anything in the way of explanation as to that mysterious connexion between the cardiac and other ganglia of the sympathetic and the cerebral and true spinal system, nor of the *modus operandi* of the influence which an inflamed heart would exercise through this relation upon the brain and nervous system.

CASES ILLUSTRATIVE OF THE INFLUENCE OF BELLADONNA.

By RICHARD HUGHES, M.R.C.S., L.R.C.P.Ed.(Exam.), Surgeon to the Brighton Orthopædic Hospital.

In two papers—one contained in the *BRITISH MEDICAL JOURNAL* for June 2nd, 1860, and the other in the *London Medical Review* for August 1860—I have discussed at some length the physiological influence of belladonna. The conclusions to which I have there come are, that this drug is an excitant of the sympathetic nervous system, and a depressant (mainly by local action) of the cerebro-spinal system.

In the following paper, I propose to exhibit some results of the practical application of this theory, and to point out the farther indications for practice thereby afforded. And I will begin with cases where the action of this drug upon the sympathetic system appears to be the *rationale* of its curative influence.

CASE I. Enuresis. July 1859. George M., aged 14, had long suffered from nocturnal incontinence of urine. There was nothing particular about the general appearance of the patient, or that of his urine. I ordered small doses of tincture of cantharides, which relieved him temporarily; but immediate relapse followed its discontinuance. I now prescribed one-third of a grain of extract of belladonna twice a day, in cinnamon-water. In three days his complaint left him; a slight relapse, about a week subsequently, was immediately removed by a few more doses of the medicine; and he has since been entirely free of his ailment.

REMARKS. Incontinence of urine may obviously result from two immediate causes. 1. From over-action of the longitudinal fibres of the muscular coat of the bladder (detrusor urinae). This is usually excited by some irritating quality of the urine, and is best removed by alkaline medicines and opiate suppositories. 2. From weakness or paralysis of its sphincter. This second and far more common cause of enuresis, usually met with in delicate children, requires for its cure some agent capable of toning the sphincter muscle, either immediately, or through the medium of its nerves. Such an agent we have, according to my views, in belladonna; and, accordingly, the accounts of the influence of this drug over the disease from all parts of Great Britain and the Continent are highly encouraging. It rarely fails to overcome it, and in its use is subject to none of the evils occasionally produced by cantharides.

In prescribing this drug for children, the discovery of Dr. Fuller as to its comparatively slight effects on young subjects must be borne in mind; and what would seem to be large doses must be unhesitatingly used.

CASE II. Milk-Abscess. July 1860. Emily H., aged 24, nine weeks after confinement caught cold. When I saw her, I found at the upper part of the left breast a hard swelling elevated above the surface, of about the size of a hen's egg; it was very painful, red, hot, and tender to the touch. There was also pain in the right side, constipation, headache, high fever; and, in the course of the preceding night, there had been some delirium. The fever had somewhat abated since vomiting a quantity of pure bile early in the morning. These symptoms pointed to concomitant derangement of the liver, and soon yielded to a mercurial pill and saline draught. The state of the breast I treated by suspending the mamma by a

handkerchief tied over the shoulder, and keeping the swelling covered with the unguentum belladonnae of the *Pharmacopœia*. In twelve hours, all pain had ceased, and there had been no increase of swelling. Next day, the redness and heat had gone, some tenderness only remaining. The swelling remained stationary and slightly tender for a few days, then slowly disappeared. The infant was not put to the breast during the mother's illness, and the general fever was sufficient to diminish the secretion of milk; but, as soon as she became convalescent, the flow of milk in both breasts returned as freely as before.

REMARKS. The localised inflammation of the breast to which (from its great tendency to suppuration) this term "milk-abscess" is commonly applied, has now been frequently treated by belladonna. This drug has been ascertained to possess galactifuge properties; and accordingly, being applied in the form of extract or ointment around the nipple in these cases, it speedily checks the secretion of milk, and with it the inflammation (*BRITISH MEDICAL JOURNAL*, Feb. 6, May 29, June 12, 1858, etc.).

The galactifuge properties of belladonna are easily explicable upon the theory of its action which I have advanced. Cl. Bernard has found that there are two ways of diminishing the secreting power of a gland—1, galvanising the sympathetic nerves which supply its blood-vessels; 2, dividing the cerebro-spinal nerves which supply its secreting cells. Belladonna, according to my view of it, both excites the sympathetic (as by galvanism), and depresses the cerebro-spinal influence (as by section) of the parts which it affects. The result of its application to a gland like the breast will therefore be to diminish alike its secreting power and the supply of the material on which that power is exercised. And this galactifuge property of the drug will be found invaluable in cases of still-birth, premature weaning, and others, where the check of secretion of milk is loudly called for.

But it will be noticed that, in the case I have recorded, there was no check of the secretion of milk. Not to speak of the superior practical advantages of a plan of treatment which avoids this unpleasant concomitant, it is obvious that the *rationale* of the cure must be a different one. It seems to me to resolve itself into the simple action of belladonna in exciting the vaso-motor nerves, and thereby contracting the capillaries. The essence of the inflammatory condition, to my thinking, is a morbidly dilated state of these vessels. The exudation from them, on which Professor Hughes Bennett and his school lay so much stress, is surely a consequence only of their preternatural dilatation and fulness. A substance which will contract them will save them the necessity of relieving themselves by exudation at all, and, even when this has taken place, will most surely promote its absorption. And this was the view of John Hunter. "If," he wrote in 1794, "we could discover an agent which would cause the contraction of the extreme vessels, we should have in this the true remedy for inflammation."

This case, then, of cure of threatened milk-abscess by the local application of belladonna, appears to open a wide field for the treatment of inflammation in general. The three indications which inflammation gives us for treatment are—1. To moderate the force and frequency of the heart's action; 2. To diminish the superfibrination of the blood; 3. To reduce the hyperæmia of the part affected. The old mode of effecting the first two objects—in other words, of relieving the symptomatic inflammatory fever—was to bleed, and to give mercury; which may fairly be described as robbing a patient of a portion of his life, and poisoning the remainder. A more enlightened practice fulfils the former indication by antimony, aconite, and the other substances which have this physiological effect upon the heart; and the latter by low diet, diluents, and salines.* The third indication—the reduction of the hyperæmia of the part—is commonly met by local bleeding and cold. But these are measures only applicable in their fulness to external inflammations, and even there are open to much improvement. Local bleeding, however efficacious, wastes blood, and leaves the capillaries weak and still morbidly dilated; and cold must be continuously applied, to be of avail. We should gain immensely, could we find a remedy capable of acting, whether locally or through the general system, upon the nerves of the inflamed part, so that its dilated capillaries might be aided to that contraction for which Nature is striving, and thus the hyperæmia of the part reduced without the loss of a drop of blood, or the inconveniences inseparable from leeches, cupping, and the continuous application of cold.

* I find nitre, which combines both these actions, an admirable general remedy for the mild fevers and inflammations of childhood.