

## Mortality at 30 days after hip fracture for different patient subgroups

NHFD admissions 2013	Total	Hospital stay 0-10 days			Hospital stay >10 days		
		Alive at 30 days	Died	(%)	Alive at 30 days	Died	(%)
All patients	65,535	8,350	1,890	18.5%	51,932	3,058	5.6%
Discharged alive	60,163	8,258	226	2.7%	50,637	741	1.4%
<b>Admitted from own home</b>	<b>46,414</b>	<b>6,347</b>	<b>83</b>	<b>1.3%</b>	<b>39,433</b>	<b>318</b>	<b>0.8%</b>
Discharged to own home	29,937	4,763	20	0.4%	24,983	163	0.6%
Discharged to care home	3,920	135	10	6.9%	3,741	34	0.9%
Discharged to 'hospital/rehab.'	11,046	1,310	30	2.2%	9,616	88	0.9%
Discharged to 'other'	1,052	106	20	15.9%	903	23	2.5%
<b>Admitted from a care home</b>	<b>10,519</b>	<b>1,616</b>	<b>122</b>	<b>7.0%</b>	<b>8,376</b>	<b>356</b>	<b>4.1%</b>
Discharged to a care home	9,347	1,472	114	7.2%	7,419	315	4.1%